

SCEPTER® 70 DG herbicide

Supplemental Labeling

EPA Reg. No. 241-306

FOR USE IN SOYBEANS
THIS LABEL EXPIRES APRIL 30, 1998

ROTATIONAL CROP RESTRICTIONS IN USE REGION 3

OBSERVE ALL PRECAUTIONARY STATEMENTS AND LIMITATIONS IN THE SCEPTER 70 DG LABEL BEFORE USING. SEE THE SCEPTER 70 DG LABEL FOR PRECAUTIONS, WORKER PROTECTION STANDARD REQUIREMENTS, APPLICATION INFORMATION AND USES WITH OTHER PRODUCTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

A single preplant incorporated, preemergence, or postemergence application of SCEPTER 70 DG may be made at the 2.8 ounces per acre rate.

ROTATIONAL CROP RESTRICTIONS

The following rotational crops may be planted in Use Region 3, as defined in the SCEPTER 70 DG label, which includes Nebraska (east of U.S. 81, north of U.S. 34 and also that area east of U.S. 283, south of U.S. 30, and west of U.S. 81), South Dakota (east of U.S. 81), Illinois (north of S.R. 116 west of Peoria; north of U.S. 24, east of Peoria), Wisconsin, Iowa (counties other than those listed in Use Region 2), and Minnesota (south of S.R. 210), following a soil application of SCEPTER 70 DG at 2.8 ounces per acre.

Soybeans	No restrictions
Wheat	18 months ¹
Field Corn (non IMI-CORN®)	18 months ^{1,2}
IMI-CORN	9.5 months ³
Barley & Oats	18 months
Edible Beans, Grain Sorghum & Tobacco	11 months
Sugar Beets & Red Table Beets	40 months
Other Crops	18 months ⁴

ACCEPTED
with COMMENTS
In EPA Letter Dated
APR 14 1997
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
241-306

¹ In Nebraska, east of U.S. 283, south of U.S. 30, and west of U.S. 81, wheat may be planted 4 months after a SCEPTER 70 DG application. In this geography, only field corn hybrids (IMI-CORN) which possess tolerance or resistance to SCEPTER 70 DG and other imidazolinone herbicides may be planted the spring of the year following an application of SCEPTER 70 DG (SEE NOTE 3).

(See other side)

² For USE REGION 3 of the SCEPTER 70 DG label, field corn (non IMI-CORN) may be planted as a rotational crop 18 months following the application of SCEPTER 70 DG unless extreme drought conditions develop (less than 15 inches of rainfall or irrigation is received from two weeks prior to the date of SCEPTER 70 DG application through November 15 of the same year). If the minimum rainfall requirement is not met, field corn (non IMI-CORN) may not be planted the spring of the year following the 18 month crop rotation period.


If the minimum rainfall requirement is not met, only field corn hybrids (IMI-CORN) which possess tolerance or resistance to SCEPTER 70 DG and other imidazolinone herbicides may be planted the spring of the year following the 18 month crop rotation period (SEE NOTE 3).

³ Prior to the application of SCEPTER 70 DG in Use Region 3, it is recommended that users contact their chemical dealer, seed supplier, or American Cyanamid to obtain information regarding the availability of imidazolinone tolerant field corn hybrids (IMI-CORN) which are adapted to your area.

⁴ For USE REGION 3 as defined in the USE AREA section of the SCEPTER 70 DG label; canola, strawberries, cabbage, tomatoes, potatoes, carrots, celery, cole crops, garlic, onions, spinach, asparagus, cauliflower, and broccoli may be planted 26 months after a SCEPTER 70 DG application. Other crops may be planted 18 months after a SCEPTER 70 DG application.

Consult the SCEPTER 70 DG label for spraying and application information, weeds controlled, and use rates. When tank mixing, always follow the more restrictive label.

Use of SCEPTER 70 DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risk associated with the use of SCEPTER 70 DG and, therefore, rotational crop injury is always possible.



PM 25 241-306 4/14/97 page 172
SCEPTER®
70 DG herbicide

Supplemental Labeling

EPA Reg. No. 241-306

FOR USE IN SOYBEANS
THIS LABEL EXPIRES APRIL 30, 1998

ROTATIONAL CROP RESTRICTIONS IN USE REGION 3

OBSERVE ALL PRECAUTIONARY STATEMENTS AND LIMITATIONS IN THE SCEPTER 70 DG LABEL BEFORE USING. SEE THE SCEPTER 70 DG LABEL FOR PRECAUTIONS, WORKER PROTECTION STANDARD REQUIREMENTS, APPLICATION INFORMATION AND USES WITH OTHER PRODUCTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

A single preplant incorporated, preemergence, or postemergence application of SCEPTER 70 DG may be made at the 2.8 ounces per acre rate.

ROTATIONAL CROP RESTRICTIONS

The following rotational crops may be planted in Use Region 3, as defined in the SCEPTER 70 DG label, which includes Nebraska (east of U.S. 81, north of U.S. 34 and also that area east of U.S. 283, south of U.S. 30, and west of U.S. 81), South Dakota (east of U.S. 81), Illinois (north of S.R. 116 west of Peoria; north of U.S. 24, east of Peoria), Wisconsin, Iowa (counties other than those listed in Use Region 2), and Minnesota (south of S.R. 210), following a soil application of SCEPTER 70 DG at 2.8 ounces per acre.

Soybeans	No restrictions
Wheat	18 months ¹
Field Corn (non IMI-CORN®)	18 months ^{1,2}
IMI-CORN	9.5 months ³
Barley & Oats	18 months
Edible Beans, Grain Sorghum & Tobacco	11 months
Sugar Beets & Red Table Beets	40 months
Other Crops	18 months ⁴

ACCEPTED
with COMMENTS
in EPA Letter Dated

APR 14 1997

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

241-306

¹ In Nebraska, east of U.S. 283, south of U.S. 30, and west of U.S. 81, wheat may be planted 4 months after a SCEPTER 70 DG application. In this geography, only field corn hybrids (IMI-CORN) which possess tolerance or resistance to SCEPTER 70 DG and other imidazolinone herbicides may be planted the spring of the year following an application of SCEPTER 70 DG (SEE NOTE 3).

(See other side)

102

292

² For USE REGION 3 of the SCEPTER 70 DG label, field corn (non IMI-CORN) may be planted as a rotational crop 18 months following the application of SCEPTER 70 DG unless extreme drought conditions develop (less than 15 inches of rainfall or irrigation is received from two weeks prior to the date of SCEPTER 70 DG application through November 15 of the same year). If the minimum rainfall requirement is not met, field corn (non IMI-CORN) may not be planted the spring of the year following the 18 month crop rotation period.

If the minimum rainfall requirement is not met, only field corn hybrids (IMI-CORN) which possess tolerance or resistance to SCEPTER 70 DG and other imidazolinone herbicides may be planted the spring of the year following the 18 month crop rotation period (SEE NOTE 3).

³ Prior to the application of SCEPTER 70 DG in Use Region 3, it is recommended that users contact their chemical dealer, seed supplier, or American Cyanamid to obtain information regarding the availability of imidazolinone tolerant field corn hybrids (IMI-CORN) which are adapted to your area.

⁴ For USE REGION 3 as defined in the USE AREA section of the SCEPTER 70 DG label; canola, strawberries, cabbage, tomatoes, potatoes, carrots, celery, cole crops, garlic, onions, spinach, asparagus, cauliflower, and broccoli may be planted 26 months after a SCEPTER 70 DG application. Other crops may be planted 18 months after a SCEPTER 70 DG application.

Consult the SCEPTER 70 DG label for spraying and application information, weeds controlled, and use rates. When tank mixing, always follow the more restrictive label.

Use of SCEPTER 70 DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risk associated with the use of SCEPTER 70 DG and, therefore, rotational crop injury is always possible.



292