



000241-00306-122997

Printed: 09:48:54 Thursday, 24 Jun, 1999 # 43 / 3443

Systems Integration Group, Inc.

PM 25

241-306

12/29/97

Page 182

Apr. 20, 1998

3:13PM

PESTICIDE MGMT DIV

No. 9516

E. 3/7



Supplemental Labeling

EPA Reg. No. 241-306

FOR USE IN EASTERN COTTONWOOD (*Populus deltoides*) AND HYBRID POPLAR CLONES INCLUDING CROSSES BETWEEN *Populus deltoides*, *Populus maximowiczii*, *Populus nigra* and *Populus trichocarpa* GROWN ON FIBER FARMS

NOT FOR USE IN CALIFORNIA

THIS LABEL EXPIRES, APRIL 30, 1999

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE SCEPTER 70 DG HERBICIDE LABELING BEFORE USING. SEE THE SCEPTER 70 DG HERBICIDE LEAFLET LABEL FOR PRECAUTIONS, WORKER PROTECTION STANDARD REQUIREMENTS, APPLICATION INFORMATION, WEEDS CONTROLLED, USES WITH OTHER PRODUCTS AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE


It is violation of Federal law to use this product in a manner inconsistent with its labeling. The use of SCEPTER 70 DG herbicide not consistent with this labeling may result in injury to crops. **DO NOT** apply this product through any type of irrigation system.

This label must be in the possession of the user at the time of herbicide application.

GENERAL INFORMATION

SCEPTER 70 DG herbicide applied preemergence or postemergence provides early season weed control and residual activity on broadleaf weeds and grass weeds in eastern cottonwood and hybrid poplars, grown on fiber farms or tree plantations.

When tank-mixing herbicides, always follow the most restrictive label.



APPLICATION INSTRUCTIONS

Apply 2.8 to 5.6 oz. SCEPTER 70 DG herbicide per acre after planting and before the trees break dormancy (bud swell stage). SCEPTER 70 DG herbicide may be tank mixed with 1.2 to 4.8 quarts PENDULUM® 3.3 EC herbicide to improve preemergence control of grasses. SCEPTER 70 DG herbicide may also be applied as a sequential preemergence or postemergence treatment following application of PENDULUM 3.3 EC herbicide.

Applications of 2.8 to 5.6 oz. per acre SCEPTER 70 DG herbicide may also be made over the top of actively growing trees. Clones may vary in their tolerance to SCEPTER. Therefore, before making over the top applications on actively growing trees, spray a few trees and observe for plant damage over a period of six weeks prior to full scale application. For best crop safety on actively growing trees, do not include a surfactant. Do not tank mix with PENDULUM 3.3 EC herbicide for applications over the top of actively growing trees.

Refer to the SCEPTER 70 DG herbicide label for additional information on application instructions.

ROTATIONAL CROP INFORMATION

Refer to rotational crop restrictions listed in the SCEPTER 70 DG herbicide label.

Use of SCEPTER 70 DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

**CYANAMID**

American Cyanamid Company
North America Agricultural Products Division
Specialty Products Department
One Campus Drive
Parlappany, NJ 07054

©1997

7350

• Registered Trademarks, American Cyanamid Company

PE - XXXXXX

12/12/97