



000241-00306-123098

Printed: 09:49:45 Thursday, 24 Jun, 1999 # 47 / 3447

Systems Integration Group, Inc.



Supplemental Labeling

EPA Reg. No. 241-306

FOR USE IN EASTERN COTTONWOOD (*Populus deltoides*) AND HYBRID POPLAR CLONES INCLUDING CROSSES BETWEEN *Populus deltoides*, *Populus maximowiczii*, *Populus nigra* and *Populus trichocarpa* GROWN ON FIBER FARMS

ONLY FOR DISTRIBUTION AND USE IN THE FOLLOWING STATES


Alabama	Kansas	Nebraska	South Dakota
Arkansas	Kentucky	New Jersey	Tennessee
Florida	Louisiana	North Carolina	Texas
Georgia	Maryland	Ohio	Virginia
Idaho	Michigan	Oklahoma	West Virginia
Illinois	Minnesota	Oregon	Washington
Indiana	Mississippi	Pennsylvania	Wisconsin
Iowa	Missouri	South Carolina	

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE SCEPTER 70 DG HERBICIDE LABELING BEFORE USING. SEE THE SCEPTER 70 DG HERBICIDE LEAFLET LABEL FOR PRECAUTIONS, WORKER PROTECTION STANDARD REQUIREMENTS, APPLICATION INFORMATION, WEEDS CONTROLLED, USES WITH OTHER PRODUCTS AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is violation of Federal law to use this product in a manner inconsistent with its labeling. The use of SCEPTER 70 DG herbicide not consistent with this labeling may result in injury to crops. **DO NOT** apply this product through any type of irrigation system.

This labeling must be in the possession of the user at the time of herbicide application.



GENERAL INFORMATION

SCEPTER 70 DG herbicide applied preplant incorporated, preemergence or postemergence provides early season weed control and residual activity on broadleaf weeds and grass weeds in eastern cottonwood and hybrid poplars, grown on fiber farms or tree plantations.

When tank-mixing herbicides, always follow the most restrictive label.

APPLICATION INSTRUCTIONS

Applications prior to planting trees: Apply 2.8 to 5.6 oz SCEPTER 70 DG herbicide per acre preplant incorporated, preemergence or postemergence to control labeled weeds. For any of these methods tank mix with 1.2 to 4.8 quarts PENDULUM® 3.3 EC herbicide to improve preemergence control of weeds listed on the PENDULUM 3.3 EC label.

For preplant incorporated applications, SCEPTER 70 DG may be tank mixed with trifluralin. Refer to the trifluralin label for specific use recommendations, rates and weeds controlled.

For postemergence applications, glyphosate may be used with SCEPTER 70 DG alone or in combination with PENDULUM 3.3 EC. Refer to the glyphosate label for specific use recommendations, rates and weeds controlled.

Postplant applications prior to trees breaking dormancy (bud swell stage): Apply 2.8 to 5.6 oz SCEPTER 70 DG herbicide per acre preemergence or postemergence to control labeled weeds. Tank mix with 1.2 to 4.8 quarts PENDULUM 3.3 EC herbicide to improve preemergence control of weeds listed on the PENDULUM 3.3 EC label.

Postplant applications on actively growing trees: Apply 2.8 to 5.6 oz SCEPTER 70 DG herbicide per acre preemergence or postemergence to control labeled weeds. Different clonal lines may vary in their tolerance to SCEPTER 70 DG. Therefore, before making over the top applications on actively growing trees, spray a few trees and observe for plant damage over a period of six weeks prior to full scale application. For best crop safety on actively growing trees, do not include a surfactant. Do not tank mix with PENDULUM 3.3 EC herbicide.

Do not apply more than 5.6 oz SCEPTER 70 DG herbicide per acre per year.

Refer to the SCEPTER 70 DG herbicide label for additional information on application instructions.



WEEDS CONTROLLED

When applied as directed, SCEPTER 70 DG herbicide will provide postemergence control of the following weed species. Refer to SCEPTER 70 DG herbicide label for weeds controlled by soil applications.

5.6 oz. SCEPTER 70 DG per acre

<u>Common Name</u>	<u>Scientific Name</u>
Broadleaf Weeds	
Carpeweed	<i>Mollugo verticillata</i>
Cocklebur, common	<i>Xanthium strumarium</i>
Eclipta	<i>Eclipta prostrata</i>
Jimsonweed	<i>Datura stramonium</i>
Mustard, wild	<i>Brassica kaber</i>
Nightshade, eastern black	<i>Solanum ptycanthum</i>
Poinsettia, wild	<i>Euphorbia heterophylla</i>
Prickly sida	<i>Sida spinosa</i>
Smartweed, ladysthumb	<i>Polygonum persicaria</i>
Smartweed, Pennsylvania	<i>Polygonum pennsylvanicum</i>
Sunflower, common	<i>Helianthus annuus</i>
Pigweed	
Palmer	<i>Amaranthus palmeri</i>
Redroot	<i>Amaranthus retroflexus</i>
Smooth	<i>Amaranthus hybridus</i>
Spiny	<i>Amaranthus spinosus</i>
Venice mallow	<i>Hibiscus trionum</i>
Grasses	
Corn, volunteer*	<i>Zea mays</i>
Foxtail, giant	<i>Setaria faberi</i>
Foxtail, yellow	<i>Setaria glauca</i>
Panicum, fall	<i>Panicum dichotomiflorum</i>

* A postemergence application of SCEPTER 70 DG herbicide will control only those field corn hybrids which DO NOT possess tolerance or resistance to SCEPTER 70 DG herbicide or other imidazolinone herbicides.



® Registered Trademarks, American Cyanamid Company
 PE - 12331 Rev. 2 12/7/98

