

E
7 30
2-307

EPA Reg. No. 241-307

FOR USE IN SOYBEANS

USE AREA ADDITIONS

OBSERVE ALL PRECAUTIONARY STATEMENTS AND LIMITATIONS ON THE TRI-SCEPT LABEL. See the TRI-SCEPT label for mixing, spraying, application information, weed species controlled and follow crop restrictions.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This label should be in the possession of the user at the time of pesticide application.

Observe all cautions and limitations in this leaflet. The use of TRI-SCEPT not consistent with this label can result in injury to crops, animals, or persons. Keep container closed to avoid spills and contamination.

DO NOT apply this product through any type of irrigation system.

USE AREAS

In addition to those areas stated on the TRI-SCEPT leaflet label, TRI-SCEPT herbicide may also be applied in the following areas: All of Michigan and Wisconsin, Southern Minnesota, South Dakota, east of U.S. 81, and Kansas, (east of U.S. 81) and the counties of Cloud, Ellsworth, Harvey, Jewell, Lincoln, Mitchell, McPherson, Ottawa, Republic, Saline, Sedgewick, and Sumner.

*Registered Trademark of American Cyanamid Company



BEST AVAILABLE COPY

ACTIVE INGREDIENTS:

Ammonium salt of imazaquin (2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-quinolinecarboxylic acid)*.....	5.00%
trifluralin: 1,1,1-trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine	28.90%
INERT INGREDIENTS	65.92%
TOTAL	100.00%

Contains petroleum distillates

*Equivalent to 4.83%, 2-[4,5-dihydro-4-methyl-4-(1-methylethyl)5-oxo-1H-imidazol-2-yl]-3-quinolinecarboxylic acid.

TRI-SCEPT contains 3.0 pounds of active ingredients per gallon (0.43 pounds active ingredient of imazaquin and 2.57 pounds active ingredient of trifluralin per gallon).

EPA Reg. No. 241-307

EPA Est. No. 32761-MO-3


KEEP OUT OF THE REACH OF CHILDREN
WARNING!/AVISO!

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

In case of emergency endangering life or property involving this product, call collect, day or night, area code 201-835-3100

See Next Page For Additional Precautionary Statements.

AMERICAN CYANAMID COMPANY
AGRICULTURAL DIVISION
CROP PROTECTION CHEMICALS DEPARTMENT
WAYNE, NEW JERSEY 07470



®Registered Trademark of American Cyanamid Company

D43
12/88

Net Contents: 2.5 gallons

STATEMENT OF PRACTICAL TREATMENT

If in eyes: hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention if irritation persists.

If on skin: wash with plenty of soap and water. Get medical attention if irritation persists.

If inhaled: remove victim to fresh air.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

WARNING!

Causes substantial but temporary eye injury.

HARMFUL IF ABSORBED THROUGH THE SKIN OR INHALED

DO NOT get in eyes or on clothing. Wear goggles or face shield. Avoid breathing spray mist. Avoid contact with skin. Wash skin thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. DO NOT apply directly to water or wetlands (swamps, bogs, or marshes). Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. DO NOT contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of pesticide application.

Observe all cautions and limitations in this leaflet. The use of TRI-SCEPT not consistent with this label can result in injury to crops, animals, or persons. Keep container closed to avoid spills and contamination.

DO NOT apply this product through any type of irrigation system.

STORAGE AND DISPOSAL

STORAGE: KEEP FROM FREEZING. DO NOT STORE BELOW 32°F. SHAKE WELL BEFORE USING. DO NOT contaminate water, food, or feed by storage or disposal.

DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

BEST AVAILABLE COPY

DISCLAIMER

The label instructions for the use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the use or application of the product contrary to label instructions, all of which are beyond the control of American Cyanamid Company. All such risks shall be assumed by the user.

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use.

Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

American Cyanamid Company makes no other express or implied warranty, including any other express or implied warranty of FITNESS or MERCHANTABILITY. USER assumes the risk of any use contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable by American Cyanamid company.

GENERAL INFORMATION

For control of grass and broadleaf weeds, apply TRI-SCEPT as a preplant incorporated treatment.

After TRI-SCEPT is applied, some susceptible weeds emerge, growth stops, and then the weeds either die or are not competitive with the crop.

Some dwarfing of soybean plants may occur with TRI-SCEPT applications. This is the result of shortened plant internodes and has no effect on soybean yields.

There should be an interval of 90 days between the TRI-SCEPT application and soybean harvest.

NOTE: DO NOT apply TRI-SCEPT postemergence to soybeans as crop injury may occur.

Crops other than soybeans, such as cotton, corn and vegetables, may be injured by spray drift or other indirect contact with TRI-SCEPT.

To avoid injury to sensitive crops from spray drift, follow all use directions and precautions in Spraying Instructions on page 8.

To avoid injury to sensitive crops, spray equipment used for TRI-SCEPT applications must be drained and thoroughly cleaned with water before being used to apply other products to these crops.

See page 13 for Rotational Crop Restrictions.

Replanting: If replanting is necessary in field previously treated with TRI-SCEPT, the field may be replanted to soybeans. Rework the soil no deeper than the treated zone. DO NOT apply a second treatment of TRI-SCEPT.

BEST AVAILABLE COPY

MIXING INSTRUCTIONS

Fill the spray tank one-half to three-quarters full with clean water. While agitating add the required amount of product and then fill the remainder of the tank with water. Maintain agitation while spraying to ensure a uniform spray mixture.

SPRAYING INSTRUCTIONS

NOTE: DO NOT make applications when spray may be carried by wind to sensitive crops.

Avoid overlaps when spraying.

GROUND APPLICATIONS:

Uniformly apply with properly calibrated ground equipment in 10 to 40 gallons of water per acre. A spray pressure of 20 to 40 psi is recommended.

DO NOT apply with ground equipment when wind velocity is greater than 10 mph.

AERIAL APPLICATIONS:

Uniformly apply with properly calibrated aerial equipment in 5 or more gallons of water per acre.


DO NOT ALLOW TRI-SCEPT TO DRIFT OUTSIDE OF THE INTENDED TREATED FIELD.

DO NOT SPRAY WHEN ENVIRONMENTAL CONDITIONS COULD CAUSE DRIFT.

Applicator is responsible for any loss or damage which results from spraying TRI-SCEPT in a manner other than recommended in this label. In addition, applicator must follow all applicable state and local regulations and ordinances in regard to spraying.

USE RATE

Apply TRI-SCEPT at a rate of 2.33 pints per acre.



APPLICATION INSTRUCTIONS

PREPLANT INCORPORATED APPLICATIONS: Apply TRI-SCEPT before planting and thoroughly incorporate into the top 2 inches of soil.

- (a) Field cultivator with 3 to 4 rows of sweeps (Do not use chisel points), spaced at intervals of 7 inches or less and staggered so that no soil is left unturned, set to cut 3 inches deep, and operated two times at more than 5 mph with the second pass made at an angle to the first.
- (b) PTO driven equipment should be set to cut 2 inches deep and operate one time at 4 mph or less.
- (c) Rolling cultivator set to cut 2 to 3 inches deep and operated two times at 6 to 8 mph. Use only on coarse textured soils.
- (d) Lely Roterra² set and operated according to the manufacturer's directions to thoroughly incorporate TRI-SCEPT to a depth of 2 inches.
- (e) Do-All set and operated according to the manufacturer's directions to thoroughly incorporate TRI-SCEPT to a depth of 2 inches.

(f) Disk harrow set to cut 3 to 4 inches deep and operated in two different directions at 4 to 6 mph.

Incorporate into the soil within 24 hours after application. TRI-SCEPT may be applied during land preparation up to 45 days prior to planting or immediately prior to planting. Adequate moisture is required for activation of TRI-SCEPT.

If soybeans are planted on beds, apply and incorporate after bed formation.

²Lely Roterra is a Trademark of Lely.

WEEDS CONTROLLED



The following grass and broadleaf weeds are controlled by recommended treatments of TRI-SCEPT:

GRASSES

- Barnyardgrass (Echinochloa crus-galli)
- Crabgrass (Digitaria spp.)
- Crowfootgrass (Dactyloctenium aegyptium)
- Cupgrass, Woolly (Eriochloa villosa)
- Foxtail
 - Giant (Setaria faberi)
 - Green (Setaria viridis)
 - Robust (Setaria sp.)
 - Yellow (Setaria glauca)
- Goosegrass (Fleusine indica)
- Johnsongrass, seedling (Sorghum halepense)
- Panicum
 - Fall* (Panicum dichotomiflorum)
 - Texas (Panicum texanum)
- Sandbur, Field (Cenchrus incertus)
- Shattercane* (Sorghum bicolor)
- Signalgrass, Broadleaf (Brachiaria platyphylla)

BROADLEAF WEEDS

- Bristly Starbur (Acanthospermum hispidum)
- Burcucumber (Sicyos angulatus)
- Carpetweed (Mollugo verticillata)
- Cocklebur, Common (Xanthium strumarium)
- Jimsonweed (Datura stramonium)
- Kochia (Kocnia scoparia)
- Lambsquarters, Common (Chenopodium album)
- Mallow, Venice (Hibiscus trionum)
- Morningglory
 - Palm Leaf (Ipomoea wrightii)
 - Pitted (Ipomoea lacunosa)
 - Smallflower (Jacquemontia tamnifolia)
- Mustard Species (Brassica spp.)
- Nightshade, Eastern Black (Solanum ptycanthum)
- Pigweed
 - Palmer (Amaranthus palmeri)
 - Redroot (Amaranthus retroflexus)
 - Smooth (Amaranthus hybridus)
 - Spiny (Amaranthus spinosus)
 - Waterhemp, Tall (Amaranthus tuberculatus)
- Poinsettia, Wild (Euphorbia heterophylla)
- Puncturevine (Tribulus terrestris)
- Purslane (Portulaca oleracea)
- Pusley, Florida (Richardia scabra)



Ragweed	
Common	(Ambrosia artemisiifolia)
giant	(Ambrosia trifida)
Redweed	(Melochia corchorifolia)
Sida, Prickly (Teaweed)	(Sida spinosa)
Smartweed,	
Ladysthumb	(Polygonum persicaria)
Pennsylvania	(Polygonum pennsylvanicum)
Sunflower, Common	(Helianthus annuus)
Velvetleaf	(Abutilon theophrasti)

TRI-SCEPT will aid in the control and reduce competition from:

Morningglory	
Entireleaf	(Ipomoea hederacea var. intergriuscula)
Ivyleaf	(Ipomoea hederacea)
Tall	(Ipomoea purpurea)
Nutsedge, Yellow	(Cyperus esculentus)
Spurge, Spotted	(Euphorbia maculata)

*See TRI-SCEPT plus trifluralin or PROWL® herbicide Tank Mixture section below for use directions.

®Registered Trademark Of American Cyanamid Company

TRI-SCEPT
ROTATIONAL CROP RESTRICTIONS

The following rotational crops may be planted after applying TRI-SCEPT in soybeans.

1. Wheat may be planted in the fall, four months after TRI-SCEPT application in soybeans. Rice may be planted in the spring of the year following TRI-SCEPT application (See Northern Use Area Restrictions*).
2. Eleven months after TRI-SCEPT application:
 - ✓ Barley (See Northern Use Area Restrictions*)
 - Oats (See Northern Use Area Restrictions*)
 - . Field corn (See Northern Use Area* or Southern Use Area Restrictions**)
 - . Edible beans
 - Grain sorghum
 - . Peanuts
 - . Tobacco

BEST AVAILABLE COPY

*NORTHERN USE AREA RESTRICTIONS.

For the Northern Use Area as defined in the USE AREA section of the label, DO NOT plant field corn, wheat, barley, or oats as a rotational crop in the fall or in the spring of the year following TRI-SCEPT application.

**SOUTHERN USE AREA RESTRICTIONS:

For the Southern Use Area, as defined in the USE AREA section of this label; field corn may be planted in the spring of the year following TRI-SCEPT application unless extreme drought conditions develop and less than 15 inches of rainfall or irrigation is received from the date of application through October of the year of application. For eastern Oklahoma, Arkansas, the bootheel of Missouri, Tennessee, North Carolina and states south: field corn may be planted in the spring of the year following TRI-SCEPT application unless extreme drought conditions develop and less than 15 inches of rainfall or irrigation is received within 6 months following the date of application.

ONLY rotational crops harvested at maturity may be used for feed or food.

DO NOT plant other rotational crops prior to 18 months after last TRI-SCEPT application.

DO NOT apply products containing chlorimuron ethyl (e.g. Classic, Canopy, Gemini, Lorox Plus, Preview¹, etc.), imazaquin (e.g. SCEPTER[®] [refer to the Scepter label for exceptions], Squadron[®], etc.), or imazethapyr (e.g. PURSUIT[®], PURSUIT[®] PLUS, etc.) the same year as TRI-SCEPT or injury to follow crops may occur.

DO NOT graze or feed treated soybean forage, hay, or straw to livestock.

¹Classic, Canopy, Gemini, Lorox Plus, and Preview are trademarks of E.I. duPont de Nemours and Company.

TRI-SCEPT Plus trifluralin or PROWL[®] Tank Mixture For Shattercane and Fall Panicum Infestations in Soybeans

Trifluralin or PROWL must be tank mixed with TRI-SCEPT and applied preplant incorporated if infestations of shattercane or fall panicum are anticipated.

For shattercane control, add trifluralin 4 EC or PROWL at a rate of 0.5 pint per acre on medium textured soils or 1.0 pint per acre on fine textured soils. For control of fall panicum, add 0.5 pint per acre of trifluralin or PROWL for all soil types.

Observe all precautions and limitations on the trifluralin label.

Observe all precautions and limitations on the PROWL label.

TRI-SCEPT PLUS Command² 4 EC Preplant Incorporated Tank Mixture For Use When Heavy Infestations of Velvetleaf are Anticipated

Observe all precautionary statements on the Command label before using. Use sprayers equipped with nozzles that provide accurate and uniform application. NOTE: The use of an agriculturally approved drift reducing agent is required at spray volumes of 10 to 15 gallons per acre.

Apply TRI-SCEPT plus Command 4 EC to the soil surface and uniformly incorporate to a depth of 2 inches. Incorporate within 3 hours of application. DO NOT apply aerially or through irrigation systems. DO NOT apply this tank mixture to overly moist or wet soils.

SPECIAL PRECAUTION

Command 4 EC is a volatile compound. Off-site movement of spray drift or vapors of Command can cause foliar whitening or yellowing of some plants. Prior to using Command, read and strictly follow all precautions and application instructions on the Command 4 EC label.

WEED SPECIES CONTROLLED

The TRI-SCEPT plus Command tank mixture controls heavy infestations of velvetleaf as well as the weeds listed in the WEEDS CONTROLLED section of the label.

[®]Registered Trademarks: American Cyanamid Company.



471-3

541-307

37

USE RATES

Apply TRI-SCEPT at a rate of 2 1/3 pints per acre in tank mixture with Command 4 EC at 1/4 to 1/2 pints per acre.

ROTATIONAL CROP RESTRICTIONS


Follow all rotational crop restrictions on the TRI-SCEPT and Command labels. Always follow the most restrictive label.

In the event of a crop loss due to weather conditions, soybeans can be replanted. DO NOT work the soil deeper than 2 inches.

2 Command is a trademark of FMC Corporation

USE AREA

TRI-SCEPT can be applied only in the states or parts shaded in the following map:




XXXXX - Northern Use Area (includes: Nebraska, east of U.S. 81, north of U.S. 34; South Dakota, east of U.S. 81; Illinois, north of S. R. 116 for area west of Peoria, north of U.S. 24 for area east of Peoria; Indiana, north of U.S. 24 for area west of Huntington, north of U.S. 224 for area east of Huntington; Ohio, north of U.S. 224; and all other states or portions of states shaded in the Northern Use Area map).

///// - Southern Use Area (includes: Kansas (east of U.S. 81) and the counties of Cloud, Ellsworth, Harvey, Jewell, Lincoln, Mitchell, McPherson, Ottawa, Republic, Saline, Sedgewick, and Sumner; southern portions of Illinois, Indiana, Ohio and eastern Nebraska not covered by Northern Use Area description; and all other states or portions of states shaded in the Southern Use Area map).

NOTE: See ROTATIONAL CROP RESTRICTIONS section for recommendations applying to the Northern vs. Southern Use Areas.

BEST AVAILABLE COPY



EPA Reg. No. 241-307
FOR USE IN SOYBEANS
USE AREA ADDITIONS

OBSERVE ALL PRECAUTIONARY STATEMENTS AND LIMITATIONS ON THE TRI-SCEPT LABEL. See the TRI-SCEPT label for mixing, spraying, application information, weed species controlled and follow crop restrictions.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.


This label should be in the possession of the user at the time of pesticide application.

Observe all cautions and limitations in this leaflet. The use of TRI-SCEPT not consistent with this label can result in injury to crops, animals, or persons. Keep container closed to avoid spills and contamination.


DO NOT apply this product through any type of irrigation system

USE AREAS

In addition to those areas stated on the TRI-SCEPT leaflet label, TRI-SCEPT herbicide may also be applied in the following areas: All of Michigan and Wisconsin, Southern Minnesota, South Dakota, east of U.S. 81, and Kansas, (east of U.S. 81) and the counties of Cloud, Ellsworth, Harvey, Jewell, Lincoln, Mitchell, McPherson, Ottawa, Republic, Saline, Sedgewick, and Sumner.



Registered Trademark of American Cyanamid Company



TENNESSEE, EASTERN TEXAS, VIRGINIA, WEST VIRGINIA AND WISCONSIN.

ACTIVE INGREDIENTS:

Ammonium salt of imazaquin (2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-quinolinecarboxylic acid)*..... 5.09%

trifluralin: $\text{C}_{12}\text{H}_{17}\text{F}_3\text{N}_2\text{O}_2$ -trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine 28.99%

INERT INGREDIENTS 65.92%

TOTAL 100.00%

Contains petroleum distillates

*Equivalent to 4.83%, 2-[4,5-dihydro-4-methyl-4-(1-methylethyl)5-oxo-1H-imidazol-2-yl]-3-quinolinecarboxylic acid.

TRI-SCEPT contains 3.0 pounds of active ingredients per gallon (0.43 pounds active ingredient of imazaquin and 2.57 pounds active ingredient of trifluralin per gallon).

EPA Reg. No. 241-307

EPA Est. No. 32761-MO-3

KEEP OUT OF THE REACH OF CHILDREN
WARNING!/AVISO!

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

In case of emergency endangering life or property involving this product, call collect, day or night, area code 201-835-3100

See Next Page For Additional Precautionary Statements.

AMERICAN CYANAMID COMPANY
AGRICULTURAL DIVISION
CROP PROTECTION CHEMICALS DEPARTMENT
WAYNE, NEW JERSEY 07470

®Registered Trademark of American Cyanamid Company

Net Contents: 2.5 gallons

D43
12/88

BEST AVAILABLE COPY

STATEMENT OF PRACTICAL TREATMENT

If in eyes: hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention if irritation persists.
If on skin: wash with plenty of soap and water. Get medical attention if irritation persists.
If inhaled: remove victim to fresh air.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
WARNING!**

Causes substantial but temporary eye injury.

HARMFUL IF ABSORBED THROUGH THE SKIN OR INHALED

DO NOT get in eyes or on clothing. Wear goggles or face shield. Avoid breathing spray mist. Avoid contact with skin. Wash skin thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. DO NOT apply directly to water or wetlands (swamps, bogs, or marshes). Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. DO NOT contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of pesticide application.

Observe all cautions and limitations in this leaflet. The use of TRI-SCEPT not consistent with this label can result in injury to crops, animals, or persons. Keep container closed to avoid spills and contamination.

DO NOT apply this product through any type of irrigation system.

STORAGE AND DISPOSAL

STORAGE: KEEP FROM FREEZING. DO NOT STORE BELOW 32°F. SHAKE WELL BEFORE USING. DO NOT contaminate water, food, or feed by storage or disposal.

DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

BEST AVAILABLE COPY

DISCLAIMER

The label instructions for the use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the use or application of the product contrary to label instructions, all of which are beyond the control of American Cyanamid Company. All such risks shall be assumed by the user.

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use.

Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

American Cyanamid Company makes no other express or implied warranty, including any other express or implied warranty of FITNESS or MERCHANTABILITY. USER assumes the risk of any use contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable by American Cyanamid company.

GENERAL INFORMATION

For control of grass and broadleaf weeds, apply TRI-SCEPT as a preplant incorporated treatment.

After TRI-SCEPT is applied, some susceptible weeds emerge, growth stops, and then the weeds either die or are not competitive with the crop.

Some dwarfing of soybean plants may occur with TRI-SCEPT applications. This is the result of shortened plant internodes and has no effect on soybean yields.

There should be an interval of 90 days between the TRI-SCEPT application and soybean harvest.

NOTE: DO NOT apply TRI-SCEPT postemergence to soybeans as crop injury may occur.

Crops other than soybeans, such as cotton, corn and vegetables, may be injured by spray drift or other indirect contact with TRI-SCEPT.

To avoid injury to sensitive crops from spray drift, follow all use directions and precautions in Spraying Instructions on page 8.

To avoid injury to sensitive crops, spray equipment used for TRI-SCEPT applications must be drained and thoroughly cleaned with water before being used to apply other products to these crops.

See page 13 for Rotational Crop Restrictions.

Replanting: If replanting is necessary in field previously treated with TRI-SCEPT, the field may be replanted to soybeans. Rework the soil no deeper than the treated zone. DO NOT apply a second treatment of TRI-SCEPT.

MIXING INSTRUCTIONS

Fill the spray tank one-half to three-quarters full with clean water. While agitating add the required amount of product and then fill the remainder of the tank with water. Maintain agitation while spraying to ensure a uniform spray mixture.

SPRAYING INSTRUCTIONS

NOTE: DO NOT make applications when spray may be carried by wind to sensitive crops.

Avoid overlaps when spraying.

GROUND APPLICATIONS:

Uniformly apply with properly calibrated ground equipment in 10 to 40 gallons of water per acre. A spray pressure of 20 to 40 psi is recommended.

DO NOT apply with ground equipment when wind velocity is greater than 10 mph.

AERIAL APPLICATIONS:

Uniformly apply with properly calibrated aerial equipment in 5 or more gallons of water per acre.

DO NOT ALLOW TRI-SCEPT TO DRIFT OUTSIDE OF THE INTENDED TREATED FIELD.

DO NOT SPRAY WHEN ENVIRONMENTAL CONDITIONS COULD CAUSE DRIFT.

Applicator is responsible for any loss or damage which results from spraying TRI-SCEPT in a manner other than recommended in this label. In addition, applicator must follow all applicable state and local regulations and ordinances in regard to spraying.

USE RATE

Apply TRI-SCEPT at a rate of 2.33 pints per acre.

BEST AVAILABLE COPY

APPLICATION INSTRUCTIONS

PREPLANT INCORPORATED APPLICATIONS: Apply TRI-SCEPT before planting and thoroughly incorporate into the top 2 inches of soil.

- (a) Field cultivator with 3 to 4 rows of sweeps (Do not use chisel points), spaced at intervals of 7 inches or less and staggered so that no soil is left unturned, set to cut 3 inches deep, and operated two times at more than 5 mph with the second pass made at an angle to the first.
- (b) PTO driven equipment should be set to cut 2 inches deep and operated one time at 4 mph or less.
- (c) Rolling cultivator set to cut 2 to 3 inches deep and operated two times at 6 to 8 mph. Use only on coarse textured soils.
- (d) Lely Roterra² set and operated according to the manufacturer's directions to thoroughly incorporate TRI-SCEPT to a depth of 2 inches.
- (e) Do-All set and operated according to the manufacturer's directions to thoroughly incorporate TRI-SCEPT to a depth of 2 inches.

(f) Disk harrow set to cut 3 to 4 inches deep and operated in two different directions at 4 to 6 mph.

Incorporate into the soil within 24 hours after application. TRI-SCEPT may be applied during land preparation up to 45 days prior to planting or immediately prior to planting. Adequate moisture is required for activation of TRI-SCEPT.

If soybeans are planted on beds, apply and incorporate after bed formation.

²Lely Roterra is a Trademark of Lely.

WEEDS CONTROLLED


The following grass and broadleaf weeds are controlled by recommended treatments of TRI-SCEPT:

GRASSES

- Barnyardgrass (Echinochloa crus-galli)
- Crabgrass (Digitaria spp.)
- Crowfootgrass (Dactyloctenium aegyptium)
- Cupgrass, Woolly (Eriochloa villosa)
- Foxtail
 - Giant (Setaria faberi)
 - Green (Setaria viridis)
 - Robust (Setaria sp.)
 - Yellow (Setaria glauca)
- Goosegrass (Eleusine indica)
- Johnsongrass, seedling (Sorghum halepense)
- Panicum
 - Fall* (Panicum dichotomiflorum)
 - Texas (Panicum texanum)
- Sandbur, Field (Cenchrus incertus)
- Shattercane* (Sorghum bicolor)
- Signalgrass, Broadleaf (Brachiaria platyphylla)

BROADLEAF WEEDS

- Bristly Starbur (Acanthospermum hispidum)
- Burcucumber (Sicyos angulatus)
- Carpetweed (Mollugo verticillata)
- Cocklebur, Common (Xanthium strumarium)
- Jimsonweed (Datura stramonium)
- Kochia (Kochia scoparia)
- Lambsquarters, Common (Chenopodium album)
- Mallow, Venice (Hibiscus trionum)
- Morningglory
 - Palm Leaf (Ipomoea wrightii)
 - Pitted (Ipomoea lacunosa)
 - Smallflower (Jacquemontia tamnifolia)
- Mustard Species (Brassica spp.)
- Nightshade, Eastern Black (Solanum ptycanthum)
- Pigweed
 - Palmer (Amaranthus palmeri)
 - Redroot (Amaranthus retroflexus)
 - Smooth (Amaranthus hybridus)
 - Spiny (Amaranthus spinosus)
 - Waterhemp, Tall (Amaranthus tuberculatus)
- Poinsettia, Wild (Euphorbia heterophylla)
- Puncturevine (Tribulus terrestris)
- Purslane (Portulaca oleracea)
- Pusley, Florida (Richardia scabra)



Ragweed	
Common	(Ambrosia artemisiifolia)
giant	(Ambrosia trifida)
Redweed	(Melochia corchorifolia)
Sida, Prickly (Teaweed)	(Sida spinosa)
Smartweed,	
Ladysthumb	(Polygonum persicaria)
Pennsylvania	(Polygonum pennsylvanicum)
Sunflower, Common	(Helianthus annuus)
Velvetleaf	(Abutilon theophrasti)

TRI-SCEPT will aid in the control and reduce competition from:

Morningglory	
Entireleaf	(Ipomoea hederacea var. intergriuscula)
Ivyleaf	(Ipomoea hederacea)
Tall	(Ipomoea purpurea)
Nutsedge, Yellow	(Cyperus esculentus)
Spurge, Spotted	(Euphorbia maculata)

*See TRI-SCEPT plus trifluralin or PROWL® herbicide Tank Mixture section below for use directions.

®Registered Trademark Of American Cyanamid Company

TRI-SCEPT
ROTATIONAL CROP RESTRICTIONS

The following rotational crops may be planted after applying TRI-SCEPT in soybeans:

1. Wheat may be planted in the fall, four months after TRI-SCEPT application in soybeans. Rice may be planted in the spring of the year following TRI-SCEPT application (See Northern Use Area Restrictions*).
2. Eleven months after TRI-SCEPT application:
 - ✓ Barley (See Northern Use Area Restrictions*)
 - Oats (See Northern Use Area Restrictions*)
 - Field corn (See Northern Use Area* or Southern Use Area Restrictions**)
 - Edible beans
 - Grain sorghum
 - Peanuts
 - Tobacco


BEST AVAILABLE COPY

*NORTHERN USE AREA RESTRICTIONS:

For the Northern Use Area as defined in the USE AREA section of the label, DO NOT plant field corn, wheat, barley, or oats as a rotational crop in the fall or in the spring of the year following TRI-SCEPT application.

**SOUTHERN USE AREA RESTRICTIONS:

For the Southern Use Area, as defined in the USE AREA section of this label; field corn may be planted in the spring of the year following TRI-SCEPT application unless extreme drought conditions develop and less than 15 inches of rainfall or irrigation is received from the date of application through October of the year of application. For eastern Oklahoma, Arkansas, the bootheel of Missouri, Tennessee, North Carolina and states south: field corn may be planted in the spring of the year following TRI-SCEPT application unless extreme drought conditions develop and less than 15 inches of rainfall or irrigation is received within 6 months following the date of application.



ONLY rotational crops harvested at maturity may be used for feed or food.

DO NOT plant other rotational crops prior to 18 months after last TRI-SCEPT application.

DO NOT apply products containing chlorimuron ethyl (e.g. Classic, Canopy, Gemini, Lorox Plus, Preview¹, etc.), imazaquin (e.g. SCEPTER[®] [refer to the Scepter label for exceptions], Squadron[®], etc.), or imazethapyr (e.g. PURSUIT[®], PURSUIT[®] PLUS, etc.) the same year as TRI-SCEPT or injury to follow crops may occur.

DO NOT graze or feed treated soybean forage, hay, or straw to livestock.

¹Classic, Canopy, Gemini, Lorox Plus, and Preview are trademarks of E.I. duPont de Nemours and Company.

TRI-SCEPT Plus trifluralin or PROWL[®] Tank Mixture For Shattercane and Fall Panicum Infestations in Soybeans

Trifluralin or PROWL must be tank mixed with TRI-SCEPT and applied preplant incorporated if infestations of shattercane or fall panicum are anticipated.

For shattercane control, add trifluralin 4 EC or PROWL at a rate of 0.5 pint per acre on medium textured soils or 1.0 pint per acre on fine textured soils. For control of fall panicum, add 0.5 pint per acre of trifluralin or PROWL for all soil types.

Observe all precautions and limitations on the trifluralin label.

Observe all precautions and limitations on the PROWL label.

TRI-SCEPT PLUS Command² 4 EC Preplant Incorporated Tank Mixture For Use When Heavy Infestations of Velvetleaf are Anticipated

Observe all precautionary statements on the Command label before using. Use sprayers equipped with nozzles that provide accurate and uniform application. NOTE: The use of an agriculturally approved drift reducing agent is required at spray volumes of 10 to 15 gallons per acre.

Apply TRI-SCEPT plus Command 4 EC to the soil surface and uniformly incorporate to a depth of 2 inches. Incorporate within 3 hours of application. DO NOT apply aerially or through irrigation systems. DO NOT apply this tank mixture to overly moist or wet soils.

SPECIAL PRECAUTION

Command 4 EC is a volatile compound. Off-site movement of spray drift or vapors of Command can cause foliar whitening or yellowing of some plants. Prior to using Command, read and strictly follow all precautions and application instructions on the Command 4 EC label.

WEED SPECIES CONTROLLED

The TRI-SCEPT plus Command tank mixture controls heavy infestations of velvetleaf as well as the weeds listed in the WEEDS CONTROLLED section of the label.

[®]Registered Trademarks Of American Cyanamid Company.



USE RATES ✓

Apply TRI-SCEPT at a rate of 2 1/3 pints per acre in tank mixture with Command 4 EC at 1/4 to 1/2 pints per acre.

ROTATIONAL CROP RESTRICTIONS


Follow all rotational crop restrictions on the TRI-SCEPT and Command labels. Always follow the most restrictive label.

In the event of a crop loss due to weather conditions, soybeans can be replanted. DO NOT work the soil deeper than 2 inches.

² Command is a trademark of FMC Corporation

USE AREA

TRI-SCEPT can be applied only in the states or parts shaded in the following map:



XXXXX - Northern Use Area (includes: Nebraska, east of U.S. 81, north of U.S. 34; South Dakota, east of U.S. 31; Illinois, north of S. R. 116 for area west of Peoria, north of U.S. 24 for area east of Peoria; Indiana, north of U.S. 24 for area west of Huntington, north of U.S. 224 for area east of Huntington; Ohio, north of U.S. 224; and all other states or portions of states shaded in the Northern Use Area map).

///// - Southern Use Area (includes: Kansas (east of U.S. 81) and the counties of Cloud, Ellsworth, Harvey, Jewell, Lincoln, Mitchell, McPherson, Ottawa, Republic, Saline, Sedgewick, and Sumner; southern portions of Illinois, Indiana, Ohio and eastern Nebraska not covered by Northern Use Area description; and all other states or portions of states shaded in the Southern Use Area map).

NOTE: See ROTATIONAL CROP RESTRICTIONS section for recommendations applying to the Northern vs. Southern Use Areas.

BEST AVAILABLE COPY