

EPA Reg. No. 241-307

## FOR USE IN SOYBEANS

# TRI-SCEPT FOLLOWED BY CLASSIC<sup>1</sup> POSTEMERGENCE FOR CONTROL OF SICKLEPOD

FOR DISTRIBUTION AND USE ONLY IN ALABAMA, ARKANSAS, FLORIDA, GEORGIA,  
LOUISIANA, MISSISSIPPI, NORTH CAROLINA, SOUTH CAROLINA, TENNESSEE AND  
TEXAS

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE TRI-SCEPT AND CLASSIC LABELS BEFORE USING. SEE THE TRI-SCEPT AND CLASSIC LABELS FOR SPRAYING AND APPLICATION INFORMATION. SEE THE TRI-SCEPT LABEL FOR TRI-SCEPT USE RATES. ALWAYS FOLLOW THE MORE RESTRICTIVE LABEL.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product through any type of irrigation system.

A single postemergence application of Classic at a rate of 1/2 to 3/4 ounce per acre may follow registered TRI-SCEPT applications in soybeans. This treatment will control sicklepod in addition to the Weeds Controlled on the TRI-SCEPT label. Refer to the Classic label for additional weeds controlled by Classic.

Refer to the Classic label for weed height limitations. Use the high rate of Classic if sicklepod seedlings are taller than two inches. A cultivation approximately 14 days after Classic application may be required to control sicklepod escapes.

<sup>1</sup>Classic is a Trademark of E.I. du Pont de Nemours and Company

\*Registered Trademark of American Cyanamid Company

ACCEPTED  
with COMMENTS  
in EPA Letter Dated

-1-

MAY 30 1990

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

241-307

replanted to soybeans. Rework the soil no deeper than the treated zone. DO NOT apply a second treatment of TRI-SCEPT.

## ROTATIONAL CROP RESTRICTIONS

The following rotational crops may be planted after applying TRI-SCEPT herbicide followed by Classic in soybeans:

1. Soybeans may be planted anytime.
2. Fifteen months after last herbicide application:

Barley	Grain sorghum	Rice
Edible beans	Oats	Tobacco
Field corn	Peanuts	Wheat
3. Sugarbeets may be planted 26 months after last herbicide application.
4. All other rotational crops, including cotton, may be planted 18 months after the last herbicide application unless specified otherwise in the PRECAUTIONS section of the Classic label.

Use of TRI-SCEPT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

ONLY rotational crops harvested at maturity may be used for feed or food.

DO NOT graze or feed treated soybean forage, hay or straw to livestock.

This label must be in the possession of the user at the time of herbicide application.

PR 34034 3/24/90



**CYANAMID**

American Cyanamid Company  
Agricultural Division  
Crop Protection Chemicals Department  
Wayne, NJ 07470 © 1990

EPA Reg. No. 241-307

**NOT REVIEWED**  
In Accordance with FR Notice 82-2.  
Based on Draft Labeling Dated

## FOR USE IN SOYBEANS

# TRI-SCEPT FOLLOWED BY CLASSIC<sup>1</sup> POSTEMERGENCE FOR CONTROL OF SICKLEPOD

FOR DISTRIBUTION AND USE ONLY IN ALABAMA, ARKANSAS, FLORIDA, GEORGIA,  
LOUISIANA, MISSISSIPPI, NORTH CAROLINA, SOUTH CAROLINA, TENNESSEE AND  
TEXAS

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE TRI-SCEPT AND CLASSIC LABELS BEFORE USING. SEE THE TRI-SCEPT AND CLASSIC LABELS FOR SPRAYING AND APPLICATION INFORMATION. SEE THE TRI-SCEPT LABEL FOR TRI-SCEPT USE RATES. ALWAYS FOLLOW THE MORE RESTRICTIVE LABEL.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product through any type of irrigation system.

A single postemergence application of Classic at a rate of 1/2 to 3/4 ounce per acre may follow registered TRI-SCEPT applications in soybeans. This treatment will control sicklepod in addition to the Weeds Controlled on the TRI-SCEPT label. Refer to the Classic label for additional weeds controlled by Classic.

Refer to the Classic label for weed height limitations. Use the high rate of Classic if sicklepod seedlings are taller than two inches. A cultivation approximately 14 days after Classic application may be required to control sicklepod escapes.

<sup>1</sup>Classic is a Trademark of E. I. du Pont de Nemours and Company  
\*Registered Trademark of American Cyanamid Company

**NOTE: If replanting is necessary in a field previously treated with TRI-SCEPT, the field may be replanted to soybeans. Rework the soil no deeper than the treated zone. DO NOT apply a second treatment of TRI-SCEPT.**

### **ROTATIONAL CROP RESTRICTIONS**

The following rotational crops may be planted after applying TRI-SCEPT herbicide followed by Classic in soybeans:

1. Soybeans may be planted anytime.
2. Fifteen months after last herbicide application:

Barley	Grain sorghum	Rice
Dry beans	Oats	Tobacco
Field corn	Peanuts	Wheat
3. Sugarbeets may be planted 26 months after last herbicide application.
4. All other rotational crops, including cotton, may be planted 18 months after the last herbicide application unless specified otherwise in the PRECAUTIONS section of the Classic label.

Use of TRI-SCEPT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

**ONLY** rotational crops harvested at maturity may be used for feed or food.

**DO NOT** graze or feed treated soybean forage, hay or straw to livestock.

This label must be in the possession of the user at the time of herbicide application.

