

EPA Reg. No. 241-307
WEED MANAGEMENT SYSTEMS
FOR USE IN SOUTHERN SOYBEANS

FOR DISTRIBUTION AND USE ONLY IN THE STATES OF ALABAMA, ARKANSAS,
 FLORIDA, GEORGIA, LOUISIANA, MISSISSIPPI, MISSOURI (ONLY IN BUTLER,
 DUNKLIN, MISSISSIPPI, NEW MADRID, PEMISCOT, SCOTT, AND STODDARD
 COUNTIES), OKLAHOMA, SOUTH CAROLINA, TENNESSEE, AND TEXAS

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE TRI-SCEPT AND TRIFLURALIN OR PROWL® LABELS BEFORE USING. SEE THE TRI-SCEPT LABEL FOR USE AREA, MIXING, SPRAYING AND APPLICATION INFORMATION, AND ROTATIONAL CROP RESTRICTIONS. ALWAYS FOLLOW THE MOST RESTRICTIVE LABEL.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. TRI-SCEPT may be applied preplant incorporated as stated on the TRI-SCEPT leaflet label. This label must be in the possession of the user at the time of herbicide application. DO NOT apply this product through any type of irrigation system.

SOIL APPLIED SYSTEMS

1. 2 1/3 Pints per Acre


TRI-SCEPT applied at 2 1/3 pints per acre provides the most consistent weed control over a wide range of environmental conditions to soybean producers utilizing low to moderate management intensity. The 2 1/3 pints per acre rate of TRI-SCEPT is the preferred application for heavy infestations and broad spectrum control. TRI-SCEPT may be applied preplant incorporated at 2 1/3 pints per acre from 30 days prior up to day of planting for control of the following weeds.

Grasses

Barnyardgrass	(<i>Echinochloa crus galli</i>)	Goosegrass	(<i>Elaeusine indica</i>)
Crabgrass	(<i>Digitaria spp.</i>)	Johnsongrass, Seedling	(<i>Sorghum halepense</i>)
Crowfootgrass	(<i>Dactyloctenium aegyptium</i>)	Panicum, Fall*	(<i>Panicum dichotomiflorum</i>)
Cupgrass, Woolly	(<i>Eriochloa villosa</i>)	Texas	(<i>Panicum texanum</i>)
Foxtail, Giant	(<i>Setaria faberii</i>)	Sandbur, Field	(<i>Cenchrus incertus</i>)
Green	(<i>Setaria viridis</i>)	Shattercane*	(<i>Sorghum bicolor</i>)
Robust	(<i>Setaria sp.</i>)	Signalgrass, Broadleaf	(<i>Brachiaria platyphylla</i>)
Yellow	(<i>Setaria glauca</i>)		

*See TRI-SCEPT plus trifluralin or PROWL herbicide Tank Mixture Section of the TRI-SCEPT leaflet label for use directions.

®Registered Trademark of American Cyanamid Company



Cocklebur, Common	(<i>Xanthium strumarium</i>)
Jimsonweed	(<i>Datura stramonium</i>)
Kochia	(<i>Kochia scoparia</i>)
Lambsquarters, Common	(<i>Chenopodium album</i>)
Mallow, Venice	(<i>Hibiscus trionum</i>)
Morningglory, Palm Leaf	(<i>Ipomoea wrightii</i>)
Pitted	(<i>Ipomoea lacunosa</i>)
Smallflower	(<i>Jacquemontia tamnifolia</i>)
Mustard Species	(<i>Brassica spp.</i>)
Nightshade	
Eastern Black	(<i>Solanum ptycanthum</i>)
Pigweed, Palmer	(<i>Amaranthus palmeri</i>)
Redroot	(<i>Amaranthus retroflexus</i>)
Smooth	(<i>Amaranthus hybridus</i>)
Spiny	(<i>Amaranthus spinosus</i>)
Waterhemp, Tall	(<i>Amaranthus tuberculatus</i>)

Pusley, Florida	(<i>Richardia scabra</i>)
Ragweed, Common	(<i>Ambrosia artemisiifolia</i>)
Giant	(<i>Ambrosia trifida</i>)
Redweed	(<i>Metochia corymbifolia</i>)
Sida, Prickly (Teaweed)	(<i>Sida spheosa</i>)
Smartweed, Ladysthumb	(<i>Polygonum persicaria</i>)
Pennsylvania	(<i>Polygonum pennsylvanicum</i>)
Sunflower, Common	(<i>Helianthus annuus</i>)
Velvetleaf	(<i>Abutilon theophrasti</i>)

TRI-SCEPT applied at 2 1/3 pints per acre will aid in the control of and reduce competition from:

Beggarweed, Florida	(<i>Desmodium tortuosum</i>)	Nutsedge, Yellow	(<i>Cyperus esculentus</i>)
Mexicanweed	(<i>Caperonia castanifolia</i>)	Sicklepod	(<i>Cassia obtusifolia</i>)
Morningglory, Entireleaf	(<i>Ipomoea hederacea</i> var. <i>intergruscula</i>)	Spurge, Spotted	(<i>Euphorbia maculata</i>)
Ivyleaf	(<i>Ipomoea hederacea</i>)	Texasweed	(<i>Caperonia palustris</i>)
Tall	(<i>Ipomoea purpurea</i>)		

2. 1 3/4 Pints per Acre plus trifluralin or PROWL

The 1 3/4 pints per acre option is intended for use by soybean producers utilizing intensive management practices including regular cultivation. TRI-SCEPT may be applied preplant incorporated at 1 3/4 pints per acre from 7 days prior up to day of planting.

Trifluralin (e.g. TRI-4™, Trellan®) or PROWL must be tank mixed with the 1 3/4 pint rate of TRI-SCEPT. Add trifluralin or PROWL at a rate of 3/8 pint per acre (based on 4 lbs. of active ingredient per gallon) to obtain a tank mix which is equivalent to 1/2 pint per acre of SCEPTER® herbicide plus 1 1/2 pints per acre of trifluralin or PROWL. Refer to the following table for additional tank-mix options.

TRI-SCEPT applied at a rate of:	+ Trifluralin or PROWL applied at a rate of:	= SCEPTER applied at a rate of:	+ Trifluralin or PROWL applied at a rate of:
2 1/3 pints/acre	---	2/3 pint/acre	1 1/2 pints/acre
2 1/3 pints/acre	1/2 pint/acre	2/3 pint/acre	2 pints/acre
1 3/4 pints/acre	3/8 pint/acre	1/2 pint/acre	1 1/2 pints/acre
1 3/4 pints/acre	7/8 pint/acre	1/2 pint/acre	2 pints/acre

DO NOT apply TRI SCEPT at rates below 1 3/4 pints per acre.

*Trellan is a trademark of Dow Chemical
 **Registered Trademark of American Cyanamid Company
 ***Trademark of American Cyanamid Company

1 3/4 Pints per Acre Plus trifluralin or PROWL

When applied as directed, TRI-SCEPT plus trifluralin or PROWL will control the following grasses and broad-leaf weeds.

GRASSES

Barnyardgrass	(<i>Echinochloa crus-galli</i>)	Goosegrass	(<i>Eleusine indica</i>)
Crabgrass	(<i>Digitaria spp.</i>)	Johnsongrass, seedling	(<i>Sorghum halepense</i>)
Crowfootgrass	(<i>Dactyloctenium aegyptium</i>)	Panicum, Fall	(<i>Panicum dichotomiflorum</i>)
Foxtail, Giant	(<i>Setaria faberi</i>)	Texas	(<i>Panicum texanum</i>)
Green	(<i>Setaria viridis</i>)	Sandbur, Field	(<i>Cenchrus incertus</i>)
Robust	(<i>Setaria sp.</i>)	Signalgrass, Broadleaf	(<i>Brachiaria platyphylla</i>)
Yellow	(<i>Setaria glauca</i>)		

BROADLEAF WEEDS

Carpelweed	(<i>Mollugo verticillata</i>)	Pigweed (continued)	
Cocklebur, Common	(<i>Xanthium strumarium</i>)	Waterhemp, tall	(<i>Amaranthus tuberculatus</i>)
Lambsquarters, Common	(<i>Chenopodium album</i>)	Purslane	(<i>Portulaca oleracea</i>)
Pigweed		Pusley, Florida	(<i>Richardia scabra</i>)
Palmer	(<i>Amaranthus palmeri</i>)	Sida, Prickly (Teaweed)	(<i>Sida spinosa</i>)
Redroot	(<i>Amaranthus retroflexus</i>)	Smartweed	
Smooth	(<i>Amaranthus hybridus</i>)	Ladythumb	(<i>Polygonum persicaria</i>)
Spiny	(<i>Amaranthus spinosus</i>)	Pennsylvania	(<i>Polygonum pennsylvanicum</i>)

Use of TRI-SCEPT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

Except where stated on approved supplemental labeling or the Sequential Treatments Section of the TRI-SCEPT leaflet label, **DO NOT** apply products containing chlorimuron ethyl (Classic, Canopy, Preview, Gemini, Lorox Plus²) or imazaquin (e.g. SCEPTER, SQUADRON[®], etc.) the same year as TRI-SCEPT. **DO NOT** apply products containing imazethapyr (e.g. PURSUIT[®], PURSUIT[®] PLUS, or PASSPORT[®]) the same year as TRI-SCEPT or injury to follow crops may occur.

¹Classic, Canopy, Preview, Gemini and Lorox Plus are trademarks of E. I. du Pont de Nemours and Company.
²Registered Trademarks of American Cyanamid Company
 PL 24025 12-8-90 (1)

