

FOR USE IN SOYBEANS
THIS LABEL EXPIRES APRIL 30, 1997

ROTATIONAL CROP RESTRICTIONS IN USE REGION 3

OBSERVE ALL PRECAUTIONARY STATEMENTS AND LIMITATIONS IN THE TRI-SCEPT LABEL BEFORE USING. SEE THE TRI-SCEPT LABEL FOR PRECAUTIONS, WORKER PROTECTION STANDARD REQUIREMENTS, APPLICATION INFORMATION AND USES WITH OTHER PRODUCTS.

DIRECTIONS FOR USE


It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

A single preplant incorporated application of TRI-SCEPT may be made at the 2.33 pint per acre rate.

ROTATIONAL CROP RESTRICTIONS

The following rotational crops may be planted in Use Region 3, as defined in the TRI-SCEPT label, which includes Nebraska (east of U.S. 81, north of U.S. 34 and also that area east of U.S. 283, south of U.S. 30, and west of U.S. 81), South Dakota (east of U.S. 81), Illinois (north of S.R. 116 west of Peoria; north of U.S. 24, east of Peoria), Wisconsin, Iowa (counties other than those listed in Use Region 2), and Minnesota (south of S.R. 210), following a soil application of TRI-SCEPT at 2.33 pints per acre.

Soybeans	No restrictions
Wheat	18 months*
Field Corn (non IMI-Corn™)	18 months**
IMI-Corn	9.5 months***
Barley & Oats	18 months
Edible Beans, Grain Sorghum, Peanuts, Tobacco	11 months
Sugar Beets & Red Table Beets	40 months
Other Crops	18 months****



*In Nebraska, east of U.S. 283, south of U.S. 30, and west of U.S. 81, wheat may be planted 4 months after a TRI-SCEPT application. In this geography, only field corn hybrids (IMI-Corn) which possess tolerance or resistance to TRI-SCEPT and other imidazolinone herbicides may be planted the spring of the year following an application of TRI-SCEPT (SEE NOTE ***).

**For USE REGION 3 of the TRI-SCEPT label, field corn (non IMI-Corn) may be planted as a rotational crop 18 months following the application of TRI-SCEPT unless extreme drought conditions develop (less than 15 inches of rainfall or irrigation is received from two weeks prior to the date of TRI-SCEPT application through November 15 of the same year). If the minimum rainfall requirement is not met, field corn (non IMI-Corn) may not be planted the spring of the year following the 18 month crop rotation period.

(see other side)

their chemical dealer, seed supplier, or American Cyanamid to obtain information regarding the availability of imidazolinone tolerant field corn hybrids (IMI-Corn) which are adapted to your area.

**** For USE REGION 3 as defined in the USE AREA section of the TRI-SCEPT label; canola, strawberries, cabbage, tomatoes, potatoes, carrots, celery, cole crops, garlic, onions, spinach, asparagus, cauliflower, and broccoli may be planted 26 months after a TRI-SCEPT application. Other crops may be planted 18 months after a TRI-SCEPT application.

Consult the TRI-SCEPT label for spraying and application information, weeds controlled, and use rates. When tank mixing, always follow the more restrictive label.

Use of TRI-SCEPT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risk associated with the use of TRI-SCEPT and, therefore, rotational crop injury is always possible.

®Registered Trademark of American Cyanamid Company

™Trademark of American Cyanamid Company

PE-XXXXX 12/15/95 SD



American Cyanamid Company

Agricultural Products Division

Crop Protection Chemicals Department

Wayne, NJ 07470

©1995