

**FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE, CRANBERRY, AND SMALL WHITE TYPE DRY BEANS, LIMA BEANS, AND SOUTHERN AND ENGLISH PEAS ONLY**

**REFER TO THE MAP ON PAGE 5 FOR GEOGRAPHICAL USE AREA**

**READ LABEL CAREFULLY - USE INSTRUCTIONS VARY FOR EACH GROUP**

**THIS LABEL EXPIRES DECEMBER 31, 1991**

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

**GENERAL INFORMATION**

PURSUIT application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

**DO NOT** apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application.

**USE PURSUIT ONLY** if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

\*Registered Trademark of American Cyanamid Company  
 †Dash is a trademark of BASF Corporation.

**DISCLAIMER**

**THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.**

**REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

**DO NOT** apply PURSUIT in any manner not specifically described in this label.


Sensitive crops such as sugarbeets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for post applications. **DO NOT** use crop oils, petroleum oils, liquid fertilizers or Dash<sup>†</sup>.

**BEST AVAILABLE COPY**

-1-



When PURSUIT is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

DO NOT tank-mix PURSUIT with postemergence grass herbicides. If postemergence grass herbicides are applied following PURSUIT, delay application of the postemergence grass herbicide at least 7 days following the application of PURSUIT.

## LIMA BEANS

DO NOT apply PURSUIT after June 30.

Allow at least 30 days between application and harvest.

**Preplant Incorporated Applications.** Apply PURSUIT at the broadcast rate of 2 ounces per acre to sand and loamy sand soils, or 3 ounces per acre to all other soils within 1 week before planting. Applied preplant incorporated, PURSUIT may be tank-mixed with a registered grass herbicide.

**Preemergence Applications.** Apply PURSUIT at the broadcast rate of 2 ounces per acre to sand and loamy sand soils, or 3 ounces per acre to all other soils immediately after, or up to 1 day after planting. PURSUIT may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

## NAVY, GREAT NORTHERN, RED KIDNEY, CRANBERRY, BLACK TURTLE, AND SMALL WHITE TYPE DRY BEANS

DO NOT apply PURSUIT to sand or loamy sand soils.

DO NOT apply PURSUIT after June 20.

Allow at least 60 days between application and harvest.

**Preplant Incorporated Applications.** Apply PURSUIT at the broadcast rate of 2 ounces per acre to sandy loam soils, or 3 ounces per acre to all other soils within 1 week before planting. Applied preplant incorporated, PURSUIT may be tank-mixed with a registered grass herbicide.

**Preemergence Applications.** Apply PURSUIT at the broadcast rate of 2 ounces per acre to sandy loam soils, or 3 ounces per acre to all other soils immediately after, or up to 1 day after

ing a preplant incorporated application of a registered grass herbicide.

## ENGLISH PEAS

DO NOT apply PURSUIT to sand or loamy sand soils.

Allow at least 30 days between application and harvest.

**Preplant Incorporated Applications.** Apply PURSUIT at the broadcast rate of 3 ounces per acre up to 1 week before planting. DO NOT apply PURSUIT to English peas that have been or will be treated with trifluralin.

**Preemergence Applications.** Apply PURSUIT at the broadcast rate of 3 ounces per acre immediately after, or up to 1 day after planting. DO NOT apply PURSUIT if English peas have been treated with trifluralin.

**Postemergence Applications.** Apply PURSUIT at 3 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture. **DO NOT APPLY PURSUIT POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY MAY RESULT.** DO NOT apply PURSUIT to English peas that have been treated with trifluralin.

## SOUTHERN PEAS

Allow at least 30 days between application and harvest.

**Preplant Incorporated Applications.** Apply PURSUIT at the broadcast rate of 4 ounces per acre within 1 week before planting. Applied preplant incorporated, PURSUIT may be tank-mixed with a registered grass herbicide.

**Preemergence Applications.** Apply PURSUIT at the broadcast rate of 4 ounces per acre immediately after, or up to 1 day after planting. PURSUIT may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

**Early Post Applications.** Apply PURSUIT at the broadcast rate of 4 ounces per acre when peas are at least 3 inches in height and weeds are 1 inch or less. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

**WEEDS CONTROLLED**

PURSUIT applied to sandy soils at the broadcast rate of 2 ounces per acre preplant incorporated or preemergence will control:

- Nightshade, black
- Nightshade, Eastern black
- Pigweed, redroot

PURSUIT applied at the broadcast rate of 3 ounces per acre preplant incorporated, preemergence or early postemergence will control:

- Mustard, wild
- Nightshade, black
- Nightshade, Eastern black
- Nightshade, hairy
- Pigweed, redroot

Postemergence applications of 3 ounces per acre must be to weeds less than 2 inches tall for best results.

PURSUIT applied preplant incorporated or preemergence at 4 ounces per acre will control:

**BROADLEAF WEEDS**

Carpetweed	Pigweed	Smartweed
Galinsoga	Palmer	ladysthumb
Jimsonweed*	redroot	Pennsylvania
Kochia	smooth	Spurge
Lambsquarters, common*	spiny	prostrate
Morningglory smallflower	Puncturevine	spotted
Mustard species	Purslane, common	Sunflower, common*
Nightshade black	Pusley, Florida	Velvetleaf*
Eastern Black hairy	Sida, prickly*	Waterhemp tall

**GRASSES**

Foxtail giant	Foxtail (Cont'd) robust white
giant green	yellow
green	Johnsongrass, seedling
robust purple	

\*PURSUIT must be applied preplant incorporated for best results.

Postemergence applications of 4 ounces per acre of PURSUIT applied as directed in this label will control:

**BROADLEAF WEEDS**

Weeds Controlled 1-2 leaves (1-2 inches)	Weeds Controlled 1-4 leaves (1-3 inches)	Weeds Controlled up to 8 leaves
Lambsquarters, common	Jimsonweed	Artichoke, Jerusalem (6-10")
Morningglory entireleaf	Kochia	Cocklebur (1-8")
ivyleaf	Marshelder	Pigweed (1-8")
pitted	Morningglory smallflower	Palmer
tali	Mustard species	redroot
Poinsettia, wild	Nightshade black	smooth
	Eastern black hairy	spiny
	Ragweed, common	Waterhemp, tall
	giant	
	Smartweed ladysthumb	
	Pennsylvania	
	Spurge prostrate	
	spotted	
	Sunflower, common	
	Velvetleaf	

**BEST AVAILABLE COPY**

475

## GRASSES

Weeds Controlled 1-3 leaves (1-3 inches)	Weeds Controlled 1-6 leaves (1-8 inches)
Barnyardgrass	Johnsongrass,
Crabgrass, large	seedling
smooth	Shattercane
Foxtail	
giant	
green	
giant green	
robust purple	
robust white	
yellow	
Red rice	

Postemergence applications of PURSUIT suppress and reduce competition from broadleaf signalgrass, yellow nutsedge, and rhizome Johnsongrass.

### ROTATIONAL CROP RESTRICTIONS

The following crops may be planted after applying PURSUIT as recommended in this label:

1. Two months after applying PURSUIT:  
  Snap beans (DE, MD, PA and VA only)
2. Four months after applying PURSUIT:  
  Barley  
  Edible beans (dry, snap) not listed on this label  
  Peas not listed on this label  
  Wheat
3. Nine and one-half months after applying PURSUIT:  
  Field corn  
  Peanuts  
  Tobacco

**BEST AVAILABLE COPY**

**DO NOT** plant sugarbeets, Irish potatoes, canola or rapeseed prior to 26 months after a PURSUIT application.

Sugarbeets are very sensitive to small amounts of PURSUIT in the soil. In addition to delaying the planting of sugarbeets 26 months, a field bioassay is required. The field bioassay will determine when it is safe to plant sugar beets. To conduct the field bioassay, plant a strip of sugarbeets through entire field and grow for 60 days. If no injury is observed, it is safe to plant sugarbeets the following year.

Soybeans and legume vegetable crops listed on this label can be planted anytime after a PURSUIT application.

**DO NOT** plant other rotational crops prior to 18 months after a PURSUIT application.


In the event of a crop loss, the bean or pea crop may be replanted.

**DO NOT** feed treated forage, hay, or straw to livestock.

**CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.**

595

PURSUIT can be applied to the edible legume vegetables listed only in the states or parts of states shaded in the following map:



Use area includes: South Dakota east of U.S. 81, Nebraska east of U.S. 81, Kansas east of U.S. 81 and Cloud, Ellsworth, Harvey, Jewell, Lincoln, McPherson, Mitchell, Ottawa, Republic, Saline, Sedgewick and Sumner counties, and all other states or portions of states shaded in the preceding map.



\*Registered Trademark of American Cyanamid Company  
PE-17087 2/4/91 DL