

PM-25

1910

FOR DISTRIBUTION AND USE ONLY WITHIN
ARIZONA, CALIFORNIA, NEVADA, AND UTAH

EPA Reg. No. 241-310

FOR USE IN RED KIDNEY BEANS ONLY

ACCEPTED

MAR 14 1991

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 241-310

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION
IN THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to red kidney beans. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. DO NOT apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application.

USE PURSUIT ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS. REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

*Registered Trademark of American Cyanamid Company
†Dash is a trademark of BASF Corporation

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system. This label must be in the possession of the user at time of herbicide application.

Sensitive crops such as sugarbeets, rapeseed, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT applications must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for post applications. DO NOT use crop oils, petroleum oils, liquid fertilizers or Dash[†].

DO NOT make more than one application of PURSUIT per year.

Allow at least 60 days between application and harvest.

When PURSUIT is used in combination with another herbicide, refer to respective labels for rates, methods of applications, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

DO NOT tank-mix PURSUIT with postemergence grass herbicides. If postemergence grass herbicides are applied following PURSUIT, delay application of the postemergence grass herbicide at least 7 day following the application of PURSUIT.

BEST AVAILABLE COPY

broadcast rate of 4 ounces per acre within 1 week before planting. Applied preplant incorporated, PURSUIT may be tank-mixed with a registered grass herbicide.

Preemergence Applications. Apply PURSUIT at the broadcast rate of 4 ounces per acre immediately after, or up to 2 days after planting. PURSUIT may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

Postemergence Applications. Apply PURSUIT at the broadcast rate of 4 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture. **DO NOT** apply PURSUIT postemergence before crop has at least one true leaf or crop injury may result. **DO NOT** apply PURSUIT to red kidney beans that have been treated with trifluralin.

PURSUIT applied preplant incorporated or preemergence at 4 ounces per acre will control:

BROADLEAF WEEDS

Carpetweed	Pigweed	Smartweed
Galinsoga	Palmer	ladysthumb
Jimsonweed*	redroot	Pennsylvania
Kochia	smooth	Spurge
Lambsquarters, common*	spiny	prostrate
Morningglory smallflower	Puncturevine	spotted
Mustard species	Purslane, common	Sunflower common*
Nightshade black	Pusley, Florida	Velvetleaf*
Eastern black hairy	Sida prickly*	Waterhemp, tall

GRASSES

Foxtail*
giant
giant green
green
robust purple
robust white
yellow
Johnsongrass, seedling*

*Must be preplant incorporated for best results.

BROADLEAF WEEDS

Weeds Controlled 1-2 leaves (1-2 inches)

Lambsquarters, common
Morningglory
entireleaf
ivy leaf
pitted
tall

Weeds Controlled 1-4 leaves (1-3 inches)

Jimsonweed	Ragweed,
Kochia	common
Marshelder	giant
Morningglory smallflower	Smartweed
Mustard species	ladysthumb
Nightshade black	Pennsylvania
Eastern black hairy	Spurge
	prostrate
	spotted
	Sunflower, common
	Velvetleaf

Weeds Controlled up to 8 leaves

Artichoke	Waterhemp,
Jerusalem (6-10")	tall
Cocklebur (1-8")	
Pigweed (1-8")	
Palmer	
redroot	
smooth	
spiny	

GRASSES

Weeds Controlled 1-3 leaves (1-3 inches)

Barnyardgrass
Crabgrass,
large
smooth
Foxtail
giant
green
robust purple
robust white
yellow

BEST AVAILABLE COPY

Weeds Controlled 1-6 leaves (1-8 inches)

Johnsongrass,
seedling
Shattercane

Postemergence applications of PURSUIT suppress and reduce competition from broadleaf signalgrass, yellow nutsedge, and rhizome Johnsongrass.

ROTATIONAL CROP RESTRICTIONS

The following crops may be planted after applying PURSUIT as recommended in this label:

1. Four months after applying PURSUIT:
 - Barley
 - Edible beans (dry, lima, snap) not listed on this label
 - Peas
 - Soybeans
 - Wheat
2. Nine and one-half months after applying PURSUIT:
 - Field corn
 - Peanuts
 - Tobacco

DO NOT plant sugarbeets, Irish potatoes, rapeseed or canola prior to 26 months after a PURSUIT application.

Sugarbeets are very sensitive to small amounts of PURSUIT in the soil. In addition to delaying the planting of sugarbeets 26 months, a field bioassay is required. The field bioassay will determine when it is safe to plant sugarbeets. To conduct the field bioassay, plant a strip of sugarbeets through entire field and grow for 60 days. If no injury is observed, it is safe to plant sugarbeets the following year.

DO NOT plant other rotational crops prior to 18 months after a PURSUIT application.

In the event of a crop loss, red kidney beans may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER RESTRICTIONS.

PE _____ 291 DL

 **CYANAMID**
American Cyanamid Company
Agricultural Division
Crop Protection Chemicals Department
Wayne, NJ 07470 1991

BEST AVAILABLE COPY

ACCEPTED
MAR 14 1991
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 241-310

4 2 10

EPA Reg. No. 241-310

FOR USE IN SUCCULENT PEAS, DRY EDIBLE PEAS, LENTILS, AND CHICKPEAS

PURSUIT can be used on dry edible peas (including chickpeas) and lentils in the Palouse or Camas Prairies of Washington and Idaho, and in Morrow, Umatilla, Grant and Union counties of Oregon only. PURSUIT can be used on non-irrigated succulent peas in the Palouse or Camas Prairies only.

THIS LABEL EXPIRES DECEMBER 31, 1991

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to peas and lentils. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application.

USE PURSUIT ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply PURSUIT in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT applications must be drained and thoroughly cleaned with water before being used to apply other products.

DO NOT make more than one application of PURSUIT per year.

Allow at least 30 days between application and harvest of succulent peas.

Allow at least 60 days between application and harvest of dry edible peas, chickpeas, and lentils.

When PURSUIT is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

*Registered Trademark of American Cyanamid Company

BEST AVAILABLE COPY

Preemergence Applications. Apply PURSUIT at the broadcast rate of 3 ounces per acre after planting, but prior to crop emergence.

PURSUIT may be tank-mixed with Sencor DF¹ or Lexone DF² to assist in the control of lambsquarters or mayweed-chamomile (dogfennel). Refer to the Sencor or Lexone label for proper application rates and restrictions.

WEEDS CONTROLLED

PURSUIT applied at the broadcast rate of 3 ounces per acre preplant incorporated or preemergence will control:

Mustard, wild
Nightshade, black
Nightshade, Eastern black
Nightshade, hairy
Pigweed, redroot

ROTATIONAL CROP RESTRICTIONS

The following crops may be planted after applying PURSUIT as recommended in this label:

1. Four months after applying PURSUIT:
 - Barley
 - Edible beans (dry, lima, snap)
 - Soybeans
 - Wheat
2. Nine and one-half months after applying PURSUIT:
 - Field corn
 - Peanuts
 - Tobacco

DO NOT plant sugarbeets, Irish potatoes, canola or rapeseed prior to 26 months after a PURSUIT application.

Sugarbeets are very sensitive to small amounts of PURSUIT in the soil. In addition to delaying the planting of sugarbeets 26 months, a field bioassay is required. The field bioassay will determine when it is safe to plant sugarbeets. To conduct the field bioassay, plant a strip of sugarbeets through entire field and grow for 60 days. If no injury is observed, it is safe to plant sugarbeets the following year.

Legume vegetable crops listed on this label can be planted anytime after a PURSUIT application.

DO NOT plant other rotational crops prior to 18 months after a PURSUIT application.

In the event of a crop loss, peas, lentils or chickpeas may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.

¹Sencor is a trademark of the Parent Company of Farbenfabriken Bayer GmbH, Leverkusen
²Lexone is a trademark of E. I. duPont de Nemours and Company, Inc.

PE-17086 2/691 DL

 **CYANAMID**
American Cyanamid Company
Agricultural Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1991

BEST AVAILABLE COPY

EPA Reg. No. 241-310

6910

**FOR USE IN RED KIDNEY TYPE DRY BEANS, LIMA BEANS,
AND ENGLISH PEAS ONLY**

FOR DISTRIBUTION AND USE ONLY IN ILLINOIS, IOWA, AND WISCONSIN

READ LABEL CAREFULLY - USE INSTRUCTIONS VARY FOR EACH GROUP

THIS LABEL EXPIRES DECEMBER 31, 1991

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION
IN THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

GENERAL INFORMATION

PURSUIT application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT only to the legume vegetables specifically listed in this label. DO NOT apply PURSUIT in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. DO NOT apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application.

USE PURSUIT ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

*Registered Trademark of American Cyanamid Company
'Dash' is a trademark of BASF Corporation

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply PURSUIT in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for post applications. DO NOT use crop oils, petroleum oils, liquid fertilizers or Dash¹.

BEST AVAILABLE COPY

ACCEPTED
MAR 14 1991
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 241-310

in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

DO NOT tank-mix PURSUIT with postemergence grass herbicides. If postemergence grass herbicides are applied following PURSUIT, delay application of the postemergence grass herbicide at least 7 days following the application of PURSUIT.

RED KIDNEY TYPE DRY BEANS AND LIMA BEANS

DO NOT apply PURSUIT after June 30.

Allow at least 30 days between application and harvest of lima beans.

Allow at least 60 days between application and harvest of red kidney beans.

Preplant Incorporated Applications. Apply PURSUIT at the broadcast rate of 3 ounces per acre within 1 week before planting. Applied preplant incorporated, PURSUIT may be tank-mixed with a registered grass herbicide.

Preemergence Applications. Apply PURSUIT at the broadcast rate of 3 ounces per acre immediately after, or up to 1 day after planting. PURSUIT may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

ENGLISH PEAS

DO NOT apply PURSUIT to sand or loamy sand soils.

Allow at least 30 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT at the broadcast rate of 3 ounces per acre up to 1 week before planting. DO NOT apply PURSUIT to English peas that have been or will be treated with trifluralin.

Preemergence Applications. Apply PURSUIT at the broadcast rate of 3 ounces per acre immediately after, or up to 1 day after planting. DO NOT apply PURSUIT if English peas have been treated with trifluralin.

Postemergence Applications. Apply PURSUIT at 3 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

WEEDS CONTROLLED

PURSUIT applied at the broadcast rate of 3 ounces per acre preplant incorporated, preemergence, or early postemergence will control:

Mustard, wild
Nightshade, black
Nightshade, Eastern black
Nightshade, hairy
Pigweed, redroot

Postemergence applications of 3 ounces per acre must be to weeds less than 2 inches tall for best results.

ROTATIONAL CROP RESTRICTIONS

The following crops may be planted after applying PURSUIT as recommended in this label:

- Four months after applying PURSUIT:
 - Barley
 - Edible beans (dry, snap) not listed on this label
 - Peas not listed on this label
 - Wheat
- Nine and one-half months after applying PURSUIT:
 - Field corn
 - Peanuts
 - Tobacco

DO NOT plant sugarbeets, Irish potatoes, canola or rapeseed prior to 26 months after a PURSUIT application.

Sugarbeets are very sensitive to small amounts of PURSUIT in the soil. In addition to delaying the planting of sugarbeets 26 months, a field bioassay is required. The field bioassay will determine when it is safe to plant sugarbeets. To conduct the field bioassay, plant a strip of sugarbeets through entire field and grow for 60 days. If no injury is observed, it is safe to plant sugarbeets the following year.

Soybeans and legume vegetable crops listed on this label can be planted anytime after a PURSUIT application.

DO NOT plant other rotational crops prior to 18 months after a PURSUIT application.

In the event of a crop loss, red kidney beans, lima beans, or English peas may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.

ACCEPTED
MAR 14 1991
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 241-310

8910

EPA Reg. No. 241-310

**FOR PREEMERGENCE USE IN SNAP BEANS GROWN
IN NEW JERSEY ONLY**

THIS LABEL EXPIRES DECEMBER 31, 1991

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION
IN THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to snap beans. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application.

USE PURSUIT ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply PURSUIT in any manner not specifically described in this label.

Sensitive crops such as sugarbeets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT applications must be drained and thoroughly cleaned with water before being used to apply other products.

DO NOT apply PURSUIT after June 20.

Allow at least 30 days between application and harvest.

DO NOT make more than one application of PURSUIT per year.

When PURSUIT is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

*Registered Trademark of American Cyanamid Company

BEST AVAILABLE COPY

Preemergence Applications. For suppression of pigweed, nightshade and common purslane, apply PURSUIT at the broadcast rate of 1.5 ounces per acre immediately after, or up to 1 day after planting. PURSUIT may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

ROTATIONAL CROP RESTRICTIONS

The following crops may be planted after applying PURSUIT as recommended in this label:

1. Four months after applying PURSUIT:

Barley
Wheat

2. Nine and one-half months after applying PURSUIT:

Field corn
Peanuts
Tobacco

DO NOT plant sugarbeets, Irish potatoes, canola or rapeseed prior to 26 months after a PURSUIT application.

Sugarbeets are very sensitive to small amounts of PURSUIT in the soil. In addition to delaying the planting of sugarbeets 26 months, a field bioassay is required. The field bioassay will determine when it is safe to plant sugarbeets. To conduct the field bioassay, plant a strip of sugarbeets through entire field and grow for 60 days. If no injury is observed, it is safe to plant sugarbeets the following year.

Soybeans and legume vegetable crops can be planted anytime after a PURSUIT application.

DO NOT plant other rotational crops prior to 18 months after a PURSUIT application.

In the event of a crop loss, snap beans may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.



CYANAMID
American Cyanamid Company
Agricultural Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1991

PE-17085 2/6/91 DL

BEST AVAILABLE COPY

ACCEPTED

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 241-310

Pursuit® herbicide

ACTIVE INGREDIENT:
 Ammonium salt of imazethapyr
 (±)-2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-5-ethyl-3-pyridinecarboxylic acid* 22.87%

INERT INGREDIENTS 77.13%

TOTAL 100.00%

*Equivalent to 21.6% (±)-2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-5-ethyl-3-pyridinecarboxylic acid

(1 gallon contains 2.0 pounds of active ingredient as the free acid)

EPA Reg. No. 241-310 EPA Est. No. 241-PR-2

KEEP OUT OF REACH OF CHILDREN CAUTION!

PRECAUCION! PRECAUCION AL USUARIO: Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

STATEMENT OF PRACTICAL TREATMENT


IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF IN EYES: Flush with plenty of water.

In case of an emergency endangering life or property involving this product, call collect, day or night, area code 201-835-3900

Net Contents: 15 gallons 1/51

© Registered Trademark of American Cyanamid Company



FOR USE IN SOYBEANS

CYANAMID
 American Cyanamid Company
 Agricultural Division
 Corporate Headquarters, Department
 Wayne, NJ 07470

BEST AVAILABLE COPY

PRECAUCION!
 CAUTION!
 Harmful if inhaled. Avoid breathing dust, skin, eyes, soap and water. Wash thoroughly after use.
 ENVIRONMENTAL PRECAUTIONS:
 DO NOT apply to swamps, wetlands, or other sensitive areas.
 DO NOT use equipment that has been used for herbicides.
 It is a violation of the Federal Insecticide, Fungicide, and Rodenticide Act to use this product in a manner inconsistent with the label.
 Observe all directions and restrictions on the label and on the container.
 and on the container.
 with PURSUIT herbicide.
 tent with the container.
 Keep containers tightly closed to prevent evaporation and contamination.
 DO NOT use for irrigation purposes.
 See complete label for details.
 STORAGE:
 PROHIBIT smoking, drinking, or eating in areas where this product is used.
 KEEP FROM children and animals.
 DO NOT use for irrigation purposes.
 Wastes resulting from the use of this product should be disposed of in accordance with local, state, and federal regulations.
 CONTAINERS:
 Return empty containers to the manufacturer for recycling.
 See Disposal Instructions on the label.