

Reg # 241-310

PM-25


174

EPA Reg. No. 241-310

FOR USE IN SOYBEANS

TANK-MIXTURE OF PURSUIT HERBICIDE AND COBRA¹ HERBICIDE
FOR ENHANCED CONTROL OF COMMON RAGWEED IN SOYBEANS

FOR DISTRIBUTION AND USE ONLY IN THE STATES OF
INDIANA, KENTUCKY, OHIO AND MICHIGAN



OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT LEAFLET LABEL BEFORE USING. SEE THE PURSUIT LEAFLET LABEL FOR USE AREA, MIXING AND APPLICATION INFORMATION, WEEDS CONTROLLED AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system.

GENERAL INFORMATION

Cobra provides fast burn-down of susceptible broadleaf weeds. Temporary bronzing of the soybean leaves soon after application may occur. Soybeans soon outgrow early symptoms; yield will not be effected by adding Cobra to PURSUIT.

APPLICATION RATES

Apply PURSUIT at a broadcast rate of 4 ounces per acre (1/4 pint) and Cobra at 4 ounces per acre.

WEEDS CONTROLLED

The addition of Cobra to PURSUIT at the above recommended rate will enhance the control of common ragweed. Apply to common ragweeds with up to 4 leaves.

¹Trademark of Valent U.S.A. Corporation

²Trademark of BASF Corporation

[®]Registered Trademark of American Cyanamid Company

[™]Trademark of Agsco, Inc.

BEST AVAILABLE COPY

SURFACTANTS: Use a non-ionic surfactant containing at least 80% active ingredient. Apply the surfactant at the rate of 1 qt./100 gal.

CROP OIL CONCENTRATE: Instead of a surfactant, a petroleum or vegetable seed oil (such as SUN-IT II[™]) based crop oil concentrate may be used. SUN-IT II is recommended when weeds are under moisture or temperature stress. Use SUN-IT II at the rate of 1.5 to 2 pts./acre (use the higher rate when weeds are at the maximum label size). Other crop oil concentrates should be used at 1.5 pts./acre.

II. LIQUID FERTILIZER SOLUTION

Recommended nitrogen based fertilizers include liquid fertilizers (such as 28%N, 32%N or 10-34-0) at the rate of 1 - 2 qts./acre. Use the higher rate when weeds are under moisture or temperature stress. Instead of a liquid fertilizer, spray grade ammonium sulfate may be used at the rate of 4 lbs./acre.

This label must be in the possession of the user at the time of herbicide application.

PE-17095 4/91 DL

American Cyanamid Company
Agricultural Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1991

BEST AVAILABLE COPY

3 8 4

ACCEPTED
MAY 21 1991
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 241-310

FOR DISTRIBUTION AND USE ONLY WITHIN CALIFORNIA

EPA Reg. No. 241-310

FOR USE IN RED KIDNEY BEANS ONLY

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT LEAFLET LABEL BEFORE USING.

GENERAL INFORMATION

The growth, maturity, quality and/or yield of red kidney beans treated with PURSUIT may be adversely affected if improper agronomic practices are utilized or if cold and/or wet conditions occur at or within 3 weeks after planting. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

SINCE GROWING CONDITIONS, ENVIRONMENTAL CONDITIONS AND GROWER AGRONOMIC PRACTICES ARE BEYOND THE CONTROL OF AMERICAN CYANAMID COMPANY, ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO RED KIDNEY BEANS SHALL BE ASSUMED BY THE USER. REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

ROTATIONAL CROP RESTRICTIONS

The following crops may be planted after applying PURSUIT as recommended in this label:

1. Anytime
 - Lima Beans
 - Southern (Blackeye) Peas
 - Pink & Small White Dry Beans
2. Four months after last PURSUIT application:
 - Barley
 - Wheat
3. Nine and one-half months after last PURSUIT application:
 - Field corn
 - Peanuts
 - Tobacco

DO NOT plant sugarbeets, rapeseed or canola prior to 26 months after a PURSUIT application.

DO NOT plant other rotational crops prior to 18 months after a PURSUIT application.

DO NOT feed treated red kidney beans, hay, or straw to livestock.

CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER RESTRICTIONS.

*Registered Trademark of American Cyanamid Company

BEST AVAILABLE COPY

any type of irrigation system.

Sensitive crops such as sugarbeets, rapeseed, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT applications must be drained, cleaned with detergent and water, and thoroughly rinsed before being used to apply other products.

DO NOT apply by aerial application.

DO NOT make more than one application of PURSUIT per year.

Allow at least 60 days between application and harvest.

When PURSUIT is used in combination with another herbicide, refer to respective labels for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

DO NOT apply sulfonyl urea herbicides the same year as PURSUIT or injury to follow crops may occur.

DO NOT tank-mix PURSUIT with postemergence grass herbicides. If postemergence grass herbicides are applied following PURSUIT, delay application of the postemergence grass herbicide at least 3 days following the application of PURSUIT.

DO NOT combine PURSUIT with a product whose label prohibits such a mixture.

APPLICATION RATE

Postemergence Applications. Apply PURSUIT at the broadcast rate of 4 ounces per acre when weeds are actively growing and red kidney beans have at least 1 fully expanded trifoliate leaf. **DO NOT** apply PURSUIT postemergence when the crop and weeds have been subjected to stress conditions such as temperature and moisture extremes.

FOR POSTEMERGENCE APPLICATIONS, A NONIONIC SURFACTANT MUST BE ADDED TO SPRAY SOLUTIONS CONTAINING PURSUIT. THE NONIONIC SURFACTANT SHOULD CONTAIN AT LEAST 80% ACTIVE INGREDIENT AND SHOULD BE USED AT A RATE OF 2 PINTS PER 100 GALLONS OF SPRAY MIXTURE.

PE-17010 3/9/91 DL



CYANAMID

American Cyanamid Company
Agricultural Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1991

Weeds Controlled 1-2 leaves (1-2 inches)

Lambsquarters, common
Morningglory
entireleaf
ivyleaf
pitted
tall

Weeds Controlled 1-4 leaves (1-3 inches)

Jimsonweed	Ragweed,
Kochia	common
Morningglory	giant
smallflower	Smartweed
Mustard, wild	ladysthumb
Nightshade	Pennsylvania
black	Spurge
Eastern black	prostrate
hairy	spotted
	Sunflower, common
	Velvetleaf

Weeds Controlled up to 8 leaves

Cocklebur (1-8")
Pigweed (1-8")
 redroot
 tall waterhemp

GRASSES

Weeds Controlled 1-3 leaves (1-3 inches)

Barnyardgrass	Foxtail
Crabgrass,	giant
large	green
smooth	robust purple
	yellow

Weeds Controlled 1-6 leaves (1-8 inches)

Johnsongrass,
 seedling
Shattercane

Postemergence applications of PURSUIT suppress and reduce competition from broadleaf signalgrass, yellow nutsedge, and rhizome Johnsongrass.

This label must be in the possession of the user at time of herbicide application.

BEST AVAILABLE COPY