


PURSUIT® HERBICIDE SUPPLEMENTAL LABELING

EPA Reg. No. 241-310

**FOR USE IN CHEMICAL FALLOW WEED CONTROL
IN WHEAT PRODUCTION**



OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT LEAFLET LABEL BEFORE USING. SEE THE PURSUIT LEAFLET LABEL FOR USE AREA, MIXING AND APPLICATION INFORMATION, WEEDS CONTROLLED AND ROTATIONAL CROP RESTRICTIONS.

GENERAL INFORMATION

PURSUIT is effective in controlling broadleaf and certain grasses common to wheat fields. It is effective when used in combination with other herbicides and/or tillage to maintain weed free land in the fallow period of continuous wheat production. The use of PURSUIT minimizes the need for tillage. Generally tillage is recommended to prepare a seedbed prior to planting wheat.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system.

APPLICATION RATES

Apply PURSUIT at the broadcast rate of 3 to 4 ounces/acre. Use 4 ounces/acre when weeds are at the maximum recommended size or under drought stress.

APPLICATIONS TIMING/TANK-MIXES

PURSUIT should be tank-mixed with:

- 1) glyphosate containing products such as Roundup¹, Fallow Master¹, Landmaster¹ BW or Landmaster¹ II or
- 2) atrazine

for enhanced control of broadleaf weeds and grasses.

SUMMER TO FALL APPLICATION:

Apply PURSUIT to the wheat stubble following wheat harvest. If no weeds are present at application, apply PURSUIT alone. If vegetation is present at application, include Roundup in the spray mixture for control of large weeds or weeds not controlled by PURSUIT.

A tank-mix of PURSUIT plus atrazine at 0.5 lb ai/A will control downy brome, volunteer wheat and many other weeds.

To control large weeds that have been cut by the wheat combine, or downy brome or volunteer wheat, apply PURSUIT with both Roundup and atrazine.

[®]Registered Trademark of American Cyanamid Company

¹Trademarks of Monsanto Agricultural Products Company

BEST AVAILABLE COPY

SPRING APPLICATION:

Apply PURSUIT in the spring before the weeds exceed 2 inches in size. When downy brome, volunteer wheat are present, or weeds are larger than 2 inches, a tank-mixture of PURSUIT and Roundup should be used. (Other registered glyphosate containing products may be substituted for Roundup.)

ADDITIVES

APPLICATIONS OF PURSUIT REQUIRE THE ADDITION OF A SURFACTANT OR A CROP OIL CONCENTRATE AND A LIQUID FERTILIZER SOLUTION.

SURFACTANTS: Use a non-ionic surfactant containing at least 80% active ingredient. Apply the surfactant at the rate of 1 qt./100 gal.

OR

CROP OIL CONCENTRATE: Instead of a surfactant, a petroleum or vegetable seed oil (such as SUN-IT II[®] adjuvant) based crop oil concentrate may be used. Use SUN-IT II at the rate of 1.5 to 2 pints/acre (use the high rate when weeds are at the maximum label size). Other crop oil concentrates should be used at 1.5 pints/acre.

AND

The addition of a fertilizer solution (10-34-0, 28 or 32% nitrogen) will enhance the weed control from PURSUIT. Add the fertilizer solution at the rate of 1-2 quarts/acre. Use the higher rate when weeds are under moisture or temperature stress. Instead of a liquid fertilizer, ammonium sulfate may be used at the rate of 4 lbs./acre

When PURSUIT is used in combination with other herbicides, refer to the respective label for rates, method of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

SPRAYING INSTRUCTIONS

Use a minimum of 20 gallons of water per acre when applying PURSUIT for wheat fallow weed control, unless tank-mixed with glyphosate containing products. When mixed with glyphosate use 10 to 20 gallons of water per acre.

[®]Trademark of Agsco, Inc.

WEEDS CONTROLLED

PURSUIT applied at 4 ounces/acre will control the following weeds up to 2 inches in size:

Cutleaf nightshade
Field pennycress
Flixweed
Knotweed,
 erect
 prostrate
Kochia
Mustard,
 tansy
 tumble
 wild

Pigweed,
 prostrate
 smooth
 redroot
Puncturevine
Russian thistle
Shepardspurse
Skeletonleaf bursage
Wild buckwheat

Refer to the PURSUIT leaflet label for ROTATIONAL CROP RESTRICTIONS

Consult the PURSUIT leaflet for additional precautions.

DO NOT plant wheat into treated fields within 4 months of a PURSUIT application.

This label must be in the possession of the user at the time of herbicide application.

PE-17130

11/91

American Cyanamid Company
Agricultural Division
Crop Protection Chemicals Department
Wayne, NJ 07470 •1991