

**FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE,
CRANBERRY, AND SMALL WHITE TYPE DRY BEANS, LIMA BEANS, AND
SOUTHERN AND ENGLISH PEAS ONLY**

REFER TO THE MAP ON PAGE 6 FOR GEOGRAPHICAL USE AREA

READ LABEL CAREFULLY - USE INSTRUCTIONS VARY FOR EACH GROUP

THIS LABEL EXPIRES DECEMBER 31, 1992

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

GENERAL INFORMATION

PURSUIT application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

*Dash is a trademark of RASP Corporation
*Registered Trademark of American Cyanamid Company

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE


It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply PURSUIT in any manner not specifically described in this label.

Sensitive crops such as sugarbeets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for postemergence applications. **DO NOT** use crop oils, petroleum oils, liquid fertilizers or Dash¹.



non treatments. When PURSUIT is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

LIMA BEANS

DO NOT apply PURSUIT after June 30.

Allow at least 30 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT at the broadcast rate of 2 ounces per acre to sand and loamy sand soils, or 3 ounces per acre to all other soils within 1 week before planting. Applied preplant incorporated, PURSUIT may be tank-mixed with a registered grass herbicide.

Preemergence Applications. Apply PURSUIT at the broadcast rate of 2 ounces per acre to sand and loamy sand soils, or 3 ounces per acre to all other soils immediately after, or up to 1 day after planting. PURSUIT may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

NAVY, GREAT NORTHERN, RED KIDNEY, CRANBERRY, BLACK TURTLE, AND SMALL WHITE TYPE DRY BEANS

DO NOT apply to Domino variety black turtle beans.

DO NOT apply PURSUIT to sand or loamy sand soils.

DO NOT apply PURSUIT after June 20.

Allow at least 60 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT within 1 week before planting. Applied preplant incorporated, PURSUIT may be tank-mixed with a registered grass herbicide. Refer to the following table for use rate.

Soil Type	Use Rate (oz/acre)
Sands, loamy sands	DO NOT USE
Sandy loams and all other soils with \leq 2% O.M.	2
All other soils with $>$ 2% O.M.	3

plant incorporated application of a registered grass herbicide.

ENGLISH PEAS

DO NOT apply PURSUIT to sand or loamy sand soils.

Allow at least 30 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT at the broadcast rate of 3 ounces per acre up to 1 week before planting. **DO NOT** apply PURSUIT to English peas that have been or will be treated with trifluralin.

Preemergence Applications. Apply PURSUIT at the broadcast rate of 3 ounces per acre immediately after, or up to 1 day after planting. **DO NOT** apply PURSUIT if English peas have been treated with trifluralin.

Postemergence Applications. Apply PURSUIT at 3 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture. **DO NOT APPLY PURSUIT POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY MAY RESULT.** The use of trifluralin prior to PURSUIT application may increase the likelihood and severity of crop injury.

SOUTHERN PEAS

Allow at least 30 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT at the broadcast rate of 4 ounces per acre within 1 week before planting. Applied preplant incorporated, PURSUIT may be tank-mixed with a registered grass herbicide.

Preemergence Applications. Apply PURSUIT at the broadcast rate of 4 ounces per acre immediately after, or up to 1 day after planting. PURSUIT may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

Early Postemergence Applications. Apply PURSUIT at the broadcast rate of 4 ounces per acre when peas are at least 3 inches in height. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

WEEDS CONTROLLED

PURSUIT applied to low organic matter soils at the broadcast rate of 2 ounces per acre preplant incorporated or preemergence will control light to moderate infestations of:

- Nightshade, black
- Nightshade, Eastern black
- Pigweed, redroot

PURSUIT applied at the broadcast rate of 3 ounces per acre preplant incorporated, preemergence or early postemergence will control:

- Mustard, wild
- Nightshade, black
- Nightshade, Eastern black
- Nightshade, hairy
- Pigweed, redroot

Postemergence applications of 3 ounces per acre must be to weeds less than 2 inches tall for best results.

When applied as directed at the broadcast rate of 4 ounces per acres, PURSUIT will control or reduce competition from the weeds listed below:

NOTE: C = Control, R = Reduced Competition

(The number under Maximum Leaf Stage indicates the MAXIMUM number of leaves at which weeds should be sprayed postemergence).

BROADLEAF WEEDS

WEEDS CONTROLLED	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Anoda, spurred	C	2	1 - 2
Artichoke, Jerusalem		8	6 - 10
Buffalobur	C ^{**}		
Bristly starbur		2	1 - 2
Carpetweed	C		
Cocklebur, common	C ^{**}	8	1 - 8
Galinsoga	C		
Jimsonweed	C [*]	4	1 - 3
Kochia	C	4	1 - 3
Lambsquarters	C [*]	2	1 - 2
Mallow, Venice	R		
Morningglory,			
entireleaf	R	2	1 - 2
ivyleaf	R	2	1 - 2
pitted	R	2	1 - 2
smallflower	C	4	1 - 3
tall	R	2	1 - 2
Mustard sp.	C	4	1 - 3
Nightshade,			
black	C	4	1 - 3
Eastern black	C	4	1 - 3
hairy	C	4	1 - 3
Pigweed,			
palmer	C	8	1 - 8
redroot	C	8	1 - 8
smooth	C	8	1 - 8
spiny	C	8	1 - 8
Poinsettia, wild	C		
Puncturevine	C		
Purslane, common	C		
Pusley, Florida	C		
Sida, prickly	C [*]		

BROADLEAF WEEDS (Continued)

WEEDS CONTROLLED	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Ragweed,			
common	R	4	1 - 3
giant	R	4	1 - 3
Sage, barnyard		R	1 - 3
Smartweed,			
ladysthumb	C	4	1 - 3
Pennsylvania	C	4	1 - 3
Spurge,			
prostrate	C	4	1 - 3
spotted	C	4	1 - 3
Sunflower, common	C*	4	1 - 3
Velvetleaf	C*	4	1 - 3
Thistle, Canada		R	1 - 3
Waterhemp, tall	C	4	1 - 3

*When soil applied, common lambsquarters, jimsonweed, prickly sida, velvetleaf and common sunflower are more consistently controlled by preplant incorporated treatments.

**Use soil applications for light to moderate infestations only. Must be preplant incorporated for best results.

GRASS WEEDS*

WEEDS CONTROLLED	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Barnyardgrass	R	3	1 - 3
Crabgrass,			
large	R	3	1 - 3
smooth	R	3	1 - 3
Cupgrass, woolly		3**	1 - 3
Foxtail,			
giant	C	6	1 - 6
green	C	3	1 - 3
giant green	C	3	1 - 3
robust purple	C	3	1 - 3
robust white	C	3	1 - 3
yellow	C	3	1 - 3
Goosegrass	R		
Johnsongrass,			
seedling	C	6	1 - 8
rhizome		R	1 - 8
Panicum,			
fall	R		
Texas	R		

*When soil applied to grasses, more consistent control can be obtained from preplant incorporated treatments.

**PURSUIT controls emerged woolly cupgrass only.

GRASS WEEDS (Continued)

WEEDS CONTROLLED	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Red rice		3	1 - 3
Shattercane	R	6	1 - 8
Signalgrass, broadleaf	R	4	1 - 8
SEDGES			
Nutsedge,			
purple	R	R	1 - 3
yellow	R	R	1 - 3

DO NOT count cotyledon leaves when determining weed stage of growth.

ROTATIONAL CROP RESTRICTIONS

The following crops may be planted after applying PURSUIT as recommended in this label:

1. Anytime
 - Soybeans
 - Peanuts
 - Edible bean and pea types listed on this label (Refer to heading on page 1)
2. Two months after PURSUIT application:
 - Snap beans (DE, MD, PA and VA only)
3. Four months after PURSUIT application:
 - Barley
 - Edible bean and pea types not listed on this label (Refer to heading on page 1)
 - Rye
 - Wheat
4. Nine and one-half months after PURSUIT application:
 - Field corn
 - Field corn grown for seed*
 - Tobacco
5. Twenty-six months after PURSUIT application:

Canola (Rapeseed)	Onions
Carrots	Potatoes
Celery	Spinach
Cole crops	Sugarbeets
Garlic	Tomatoes

DO NOT plant other rotational crops prior to 18 months after a PURSUIT application.

In the event of a crop loss, the bean or pea crop may be replanted.


DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT shall be assumed by the user.


CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.

USE AREA MAP

PURSUIT can be applied to the edible legume vegetables listed only in the states or parts of states shaded in the following map:



Use area includes: South Dakota east of U.S. 81, Nebraska east of U.S. 81, Kansas east of U.S. 81, and all other states or portions of states shaded in the preceding map.



7 OF 15

EPA Reg. No. 241-310

**FOR PREEMERGENCE USE IN SNAP BEANS GROWN
IN NEW JERSEY ONLY**

THIS LABEL EXPIRES DECEMBER 31, 1992

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to snap beans. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application.

USE PURSUIT ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

*Registered Trademark of American Cyanamid Company

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply PURSUIT in any manner not specifically described in this label.

Sensitive crops such as sugarbeets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.


Spray equipment used for PURSUIT applications must be drained and thoroughly cleaned with water before being used to apply other products.

DO NOT apply PURSUIT after June 20.

Allow at least 30 days between application and harvest.

DO NOT make more than one application of PURSUIT per year.

When PURSUIT is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.



Preemergence Applications. For suppression of pigweed, nightshade and common purslane, apply PURSUIT at the broadcast rate of 1.5 ounces per acre immediately after, or up to 1 day after planting. PURSUIT may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

ROTATIONAL CROP RESTRICTIONS

The following crops may be planted after applying PURSUIT as recommended in this label:

1. Anytime
 - Soybeans
 - Peanuts
 - Edible beans and peas
2. Four months after PURSUIT application:
 - Barley
 - Rye
 - Wheat
3. Nine and one-half months after PURSUIT application:
 - Field corn
 - Field corn grown for seed*
 - Tobacco
4. Twenty-six months after PURSUIT application:
 - Canola (Rapeseed)
 - Carrots
 - Celery
 - Cole crops
 - Garlic
 - Onions
 - Potatoes
 - Spinach
 - Sugarbeets
 - Tomatoes (direct-seeded)

DO NOT plant other rotational crops prior to 18 months after a PURSUIT application.

In the event of a crop loss, snap beans may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT shall be assumed by the user.

CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.


9 of 15

FOR DISTRIBUTION AND USE ONLY WITHIN CALIFORNIA

EPA Reg. No. 241-310

FOR USE IN RED KIDNEY BEANS ONLY

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT LEAFLET LABEL BEFORE USING



GENERAL INFORMATION

Entry into treated fields is not permitted without protective clothing until sprays have dried.

The growth, maturity, quality and/or yield of red kidney beans treated with PURSUIT may be adversely affected if improper agronomic practices are utilized or if cold and/or wet conditions occur at or within 3 weeks after planting. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely. **DO NOT** apply PURSUIT postemergence after crop has begun to flower or crop injury may result.

The mode of weed killing activity involves uptake of PURSUIT by weed roots and/or foliage and rapid translocation to the growing points. Adequate soil moisture is important for optimum PURSUIT activity. When adequate soil moisture is present, PURSUIT will provide residual control of susceptible germinating weeds; activity on established weeds will depend on the weed species and the location of its root system in the soil.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

The label instructions for the use of this product reflect the opinion of experts based on research and field use. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the use of, or application of the product contrary to label instructions, all of which are beyond the control of American Cyanamid Company. All such risks shall be assumed by the user.

American Cyanamid Company shall not be responsible for losses or damages resulting from use of this product in any manner not set forth on this label. User assumes all risks associated with the use of this product in any manner not specifically set forth on this label.

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use, subject to the risks referred to above.

American Cyanamid Company's liability will be limited to a refund of the purchase price of PURSUIT. In no case shall Cyanamid or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product.

American Cyanamid Company makes no other express or implied warranty, including other express or implied warranty of FITNESS or of MERCHANTABILITY. USER assumes the risk of any use contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable by American Cyanamid Company.

USES WITH OTHER PRODUCTS (TANK-MIXES)

This product is not to be used in combination with any other product except as specifically recommended in writing by American Cyanamid Company and American Cyanamid Company shall have no liability for any loss, damage or injury arising out of its use in any such combination not so specifically recommended. If used in combination recommended by American Cyanamid Company, the liability of American Cyanamid Company shall in no manner extend to any damage, loss or injury not directly caused by the inclusion of the American Cyanamid Company product in such combination use, and in any event shall be limited to return of the account of the purchase price of the American Cyanamid Company product.

PURSUIT as recommended in this label:

1. Anytime
 - Lima Beans
 - Southern (Blackeye) Peas
 - Soybeans
 - Pink & Small White Dry Beans
2. Four months after last PURSUIT application:
 - Barley
 - Wheat
3. Nine and one-half months after last PURSUIT application:
 - Field corn
 - Peanuts
 - Tobacco
4. Twenty-six months after last PURSUIT application:

Canola (rapeseed)	Onions
Carrots	Potatoes
Celery	Spinach
Cole crops	Sugarbeets
Garlic	Tomatoes

Sugarbeets are very sensitive to small amounts of PURSUIT in the soil. In addition to delaying the planting of sugarbeets 26 months, a field bioassay is required. The field bioassay will determine when it is safe to plant sugarbeets. To conduct the field bioassay, plant a strip of sugarbeets through the entire field and grow for 60 days. If no injury is observed, it is safe to plant sugarbeets the following year.

DO NOT plant other rotational crops prior to 18 months after PURSUIT application.

DO NOT feed treated red kidney beans, hay, or straw to livestock.

Only rotational crops harvested at maturity may be used for feed or food.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system.

DO NOT apply by aerial application.

DO NOT make more than one application of PURSUIT per year.

Allow at least 60 days between application and harvest.

When PURSUIT is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label do-ages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

postemergence grass herbicides are applied following PURSUIT, delay application of the postemergence grass herbicide at least 7 days following the application of PURSUIT.

DO NOT combine PURSUIT with a product whose label prohibits such a mixture.

MIXING INSTRUCTIONS

Fill the spray tank one-half to three-quarters full with clean water. Use a calibrated measuring device to measure the required amount of PURSUIT. Add PURSUIT to the spray tank while agitating. Fill the remainder of the tank with water.

For postemergence applications, add a non-ionic surfactant as the last ingredient in the tank. Maintain agitation while spraying to ensure a uniform spray mixture. An antifoaming agent may be added to the tank if needed.

When tank mixing PURSUIT with recommended herbicides, add the other herbicide to the spray tank first and make sure it is thoroughly mixed before adding PURSUIT.

SPRAYING INSTRUCTIONS

DO NOT apply when wind velocity is greater than 10 mph, or when spray may be carried to sensitive crops. Sensitive crops include leafy vegetables, sugarbeets, and cotton.

GROUND APPLICATIONS:

Uniformly apply with properly calibrated ground equipment in 10 or more gallons of water per acre. A spray pressure of 20 to 40 psi is recommended.

Adjust the boom height to ensure proper coverage of weed foliage (according to the manufacturer's recommendation). The nozzle spacing on the boom should be 20 inches. Use only standard flat-fan nozzle tips. Avoid overlaps when spraying.

To avoid injury to sensitive crops, spray equipment used for PURSUIT applications must be drained and thoroughly cleaned with water before being used to apply other products.

APPLICATION RATE

Postemergence Applications. Apply PURSUIT at the broadcast rate of 4 ounces per acre when weeds are actively growing and red kidney beans have at least 1 fully expanded trifoliate leaf. DO NOT apply PURSUIT postemergence when the crop and weeds have been subjected to stress conditions such as temperature and moisture extremes.

FOR POSTEMERGENCE APPLICATIONS, A NONIONIC SURFACTANT MUST BE ADDED TO SPRAY SOLUTIONS CONTAINING PURSUIT. THE NONIONIC SURFACTANT SHOULD CONTAIN AT LEAST 80% ACTIVE INGREDIENT AND SHOULD BE USED AT A RATE OF 2 PINTS PER 100 GALLONS OF SPRAY MIXTURE.

For maximum weed control, cultivate 7 - 10 days following a postemergence PURSUIT application. This timely cultivation will enhance residual weed control, especially under dry conditions.

If air temperatures reach or stay below 40° for 10 or more hours, delay a PURSUIT application for 48 hours from the time temperatures increase above 40°. Unusually cool temperatures (40° or less) reduces photosynthesis and transpiration. This reduces the uptake and translocation (and effectiveness) of PURSUIT in weeds.

PURSUIT should be applied a minimum of one hour before rainfall or overhead irrigation.


WEEDS CONTROLLED

Postemergence applications of 4 ounces per acre of PURSUIT applied as directed in this label will control:

BROADLEAF WEEDS		BROADLEAF WEEDS (Continued)	
Weeds Controlled 1-2 leaves (1-2 inches)		Weeds Controlled up to 8 leaves	
Lambsquarters, common	Morningglory species entireleaf ivy leaf pitted tall	Cocklebur (1-8")	Pigweed (1-8") redroot tall waterhemp
Weeds Controlled 1-4 leaves (1-3 inches)		GRASSES	
Weeds Controlled 1-4 leaves (1-3 inches)		Weeds Controlled 1-3 leaves (1-3 inches)	
Jimsonweed	Ragweed, common giant	Barnyardgrass	Foxtail, giant green robust purple yellow
Kochia	Smartweed, ladysthumb Pennsylvania	Crabgrass, large smooth	
Morningglory, smallflower	Spurge, prostrate spotted	Weeds Controlled (1-6 leaves (1-8 inches))	
Mustard, wild	Sunflower, common	Johnsongrass, seedling	Shattercane
Nightshade, black Eastern black hairy	Velvetleaf		

Postemergence applications of PURSUIT suppress and reduce competition from broadleaf signalgrass, yellow nutsedge, and rhizome Johnsongrass.

This label must be in the possession of the user at time of herbicide application.



12 of 15

EPA Reg. No. 241-310

**FOR USE IN RED KIDNEY TYPE DRY BEANS, WHITE LUPINS,
LIMA BEANS, AND ENGLISH PEAS ONLY**

**FOR DISTRIBUTION AND USE ONLY IN ILLINOIS, IOWA, AND WISCONSIN
READ LABEL CAREFULLY - USE INSTRUCTIONS VARY FOR EACH GROUP**

THIS LABEL EXPIRES DECEMBER 31, 1992

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

GENERAL INFORMATION

PURSUIT application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application. **DO NOT** apply PURSUIT postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply PURSUIT in any manner not specifically described in this label.

Sensitive crops such as sugarbeets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.


Spray equipment used for PURSUIT application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for post applications.

DO NOT use crop oils, petroleum oils, liquid fertilizers or Dash¹.

DO NOT make more than one application of PURSUIT per year.

¹Dash is a trademark of BASF Corporation
*Registered Trademark of American Cyanamid Company



label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

RED KIDNEY TYPE DRY BEANS, WHITE LUPINS AND LIMA BEANS

DO NOT apply PURSUIT after June 30.

DO NOT apply PURSUIT to white lupins grown on sand or loamy sand soils.

Allow at least 30 days between application and harvest of lima beans.

Allow at least 60 days between application and harvest of red kidney beans or white lupins.

Preplant Incorporated Applications. Apply PURSUIT at the broadcast rate of 3 ounces per acre within 1 week before planting. Applied preplant incorporated, PURSUIT may be tank-mixed with a registered grass herbicide.

Preemergence Applications. Apply PURSUIT at the broadcast rate of 3 ounces per acre immediately after, or up to 1 day after planting. PURSUIT may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

ENGLISH PEAS

Allow at least 30 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT at the broadcast rate of 3 ounces per acre up to 1 week before planting. DO NOT apply PURSUIT to English peas that have been or will be treated with trifluralin.

Preemergence Applications. Apply PURSUIT at the broadcast rate of 3 ounces per acre immediately after, or up to 1 day after planting. DO NOT apply PURSUIT if English peas have been treated with trifluralin.

Postemergence Applications. Apply PURSUIT at 3 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

DO NOT APPLY PURSUIT POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY MAY RESULT. If English peas have been treated with trifluralin, PURSUIT may be applied ONLY at 2 ounces per acre for suppression of black nightshade. The use of trifluralin prior to PURSUIT application may increase the likelihood and severity of crop injury.

Mustard, wild
Nightshade, black
Nightshade, Eastern black

Nightshade, hairy
Pigweed, redroot

Postemergence applications of 3 ounces per acre must be to weeds less than 2 inches tall for best results.

ROTATIONAL CROP RESTRICTIONS

The following crops may be planted after applying PURSUIT as recommended in this label:

1. Anytime
 - Soybeans
 - Peanuts
 - Edible bean and pea types listed on this label (Refer to heading on page 1).
2. Four months after PURSUIT application:
 - Barley
 - Edible bean and pea types not listed on this label (Refer to heading on page 1).
 - Rye
 - Wheat
3. Nine and one-half months after PURSUIT application:
 - Field corn
 - Field corn grown for seed*
 - Tobacco
4. Twenty-six months after PURSUIT application:

Canola (Rapeseed)	Onions
Carrots	Potatoes
Celery	Spinach
Cole Crops	Sugarbeets
Garlic	Tomatoes

DO NOT plant other rotational crops prior to 18 months after a PURSUIT application.

In the event of a crop loss, red kidney beans, white lupins, lima beans, or English peas may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT shall be assumed by the user.

CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.

ACCEPTED

NAIS 1992

Under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 241-310

EPA Reg. No. 241-310

14 OF 15

FOR USE IN SUCCULENT PEAS, DRY EDIBLE PEAS, LENTILS, AND CHICKPEAS

PURSUIT can be used on succulent peas, dry edible peas (including chickpeas) and lentils in Washington (Adams, Asotin, Columbia, Garfield, Spokane, Walla Walla and Whitman counties), Idaho (Benewah, Clearwater, Idaho, Kootenai, Latah, Lewis, and Nez Perce counties), Oregon (Union, Umatilla and Wallowa counties) and Montana only.

THIS LABEL EXPIRES DECEMBER 31, 1992

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to peas and lentils. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application.

USE PURSUIT ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans. Plant peas or lentils at least 1/2 inch deep to reduce risk of crop injury.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply PURSUIT in any manner not specifically described in this label.

Sensitive crops such as sugarbeets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT applications must be drained and thoroughly cleaned with water before being used to apply other products.

DO NOT make more than one application of PURSUIT per year.

Allow at least 30 days between application and harvest of succulent peas.

Allow at least 60 days between application and harvest of dry edible peas, chickpeas, and lentils.

When PURSUIT is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

*Registered Trademark of American Cyanamid Company

DO NOT incorporate deeper than 3 inches. Refer to the PURSUIT leaflet label for incorporation instructions.

Preemergence Applications. Apply PURSUIT at the broadcast rate of 3 ounces per acre after planting, but prior to crop emergence.

PURSUIT may be tank-mixed with Sencor DF¹ or Lexone DF² to assist in the control of lambsquarters or mayweed-chamomile (dogfennel). Refer to the Sencor or Lexone label for proper application rates and restrictions.

WEEDS CONTROLLED

PURSUIT applied at the broadcast rate of 3 ounces per acre preplant incorporated or preemergence will control:

Buckwheat, wild	Nightshade, black	Pigweed, redroot
Kochia	Nightshade, Eastern black	Shepherdspurse
Mustard, wild	Nightshade, hairy	Thistle, Russian

ROTATIONAL CROP RESTRICTIONS

The following crops may be planted after applying PURSUIT as recommended in this label:

- Anytime
 - Chickpeas
 - Lentils
 - Peas
 - Peanuts
 - Soybeans
- Four months after PURSUIT application:
 - Barley
 - Edible beans (dry, lima, snap)
 - Rye
 - Wheat
- Nine and one-half months after PURSUIT application:
 - Field corn
 - Field corn grown for seed*
 - Tobacco
- Twenty-six months after PURSUIT application:
 - Canola (Rapeseed)
 - Carrots
 - Celery
 - Cole crops
 - Garlic
 - Onions
 - Potatoes
 - Spinach
 - Sugarbeets
 - Tomatoes

DO NOT plant other rotational crops prior to 18 months after a PURSUIT application.

In the event of a crop loss, peas, lentils or chickpeas may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

*Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into field treated previously with PURSUIT shall be assumed by the user.

CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.

Sencor is a trademark of the Parke-Davis Company of Fort Worth, Texas.
Lexone is a trademark of E. I. du Pont de Nemours and Company, Inc.

PE-17086 12/18/91 DL

