

PM 25 241-310

1047

PURSUIT® SUPPLEMENTAL LABELING


EPA Reg. No. 241-310

FOR USE IN SOYBEANS

**POSTEMERGENCE APPLICATIONS OF PURSUIT®
FOLLOWING SOIL APPLICATIONS OF
SCEPTER®, SQUADRON® OR TRI-SCEPT®**

FOR USE ONLY IN THE STATE OF KENTUCKY

THIS LABEL EXPIRES DECEMBER 31, 1992



OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT LEAFLET LABEL BEFORE USING. SEE THE PURSUIT LEAFLET LABEL FOR MIXING AND APPLICATION INFORMATION, WEEDS CONTROLLED AND ROTATIONAL C. P RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system.

GENERAL INFORMATION

PURSUIT herbicide may be applied postemergence to control weeds such as common cocklebur, giant ragweed, morningglories, foxtails, crabgrass or seedling Johnsongrass that escape control from soil applications of SCEPTER, SQUADRON or TRI-SCEPT.

APPLICATION RATES

Apply SCEPTER at 1/2 pts./A or SQUADRON at 2.25 pts./A preplant incorporated or preemergence, or apply TRI-SCEPT at 1.75 pts./A preemergence.

Followed by:

PURSUIT at 4 ozs./A postemergence.

®Registered Trademark of American Cyanamid Company

2011

WEEDS CONTROLLED

Postemergence applications of PURSUIT will control the following weeds:

<u>Weeds controlled</u>	<u>Maximum Leaf Stage</u>	<u>(inches)</u>
Cocklebur, common	8	1-8
Morningglories	2	1-2
Ragweed, giant	4	1-3
Crabgrass	3	1-3
Foxtails	3	1-3
Johnsongrass, seedling	6	1-8

Refer to the PURSUIT leaflet label for additional weeds controlled.

POSTEMERGENCE APPLICATIONS OF PURSUIT REQUIRE THE ADDITION OF AN ADJUVANT AND A LIQUID FERTILIZER SOLUTION.

I. ADJUVANTS

SURFACTANTS: Use a non-ionic surfactant containing at least 80% active ingredient. Apply the surfactant at the rate of 1 qt./100 gal.

OR

CROP OIL CONCENTRATE: Instead of a surfactant, a petroleum or vegetable seed oil (such as SUN-IT II[™]) based crop oil concentrate may be used. SUN-IT II is recommended when weeds are under moisture or temperature stress. Use SUN-IT II at the rate of 1.5 to 2 pts./acre (use the higher rate when weeds are at the maximum label size). Other crop oil concentrates should be used at 1.5 pts./acre.

AND

II. LIQUID FERTILIZER SOLUTION

Recommended nitrogen based fertilizers include liquid fertilizers (such as 28%N, 32%N or 10-34-0) at the rate of 1 - 2 qts./acre. Use the higher rate when weeds are under moisture or temperature stress. Instead of a liquid fertilizer, spray grade ammonium sulfate may be used at the rate of 4 lbs./acre.

BEST AVAILABLE COPY

2411

ROTATIONAL CROP GUIDELINES

When using PURSUIT applied after applications of SCEPTER, SQUADRON or TRI-SCEPT, rotational crops may be planted at the following intervals:

1. Anytime:
Soybeans
2. Nine and one-half months after PURSUIT application:
Wheat
3. Eighteen months after PURSUIT application.
All crops not listed elsewhere in the ROTATIONAL CROP GUIDELINES section
4. Twenty six months after PURSUIT application.
Canola (Rapeseed) Onions
Carrots Potatoes
Celery Spinach
Cole crops Sugarbeets
Garlic Tomatoes

Refer to the ROTATIONAL CROP RESTRICTIONS on the SCEPTER, SQUADRON or TRI-SCEPT label. Follow the more restrictive label (SCEPTER, SQUADRON, or TRI-SCEPT and PURSUIT).

This label must be in the possession of the user at the time of herbicide application.

PE-17140

1/92


American Cyanamid Company
Agricultural Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1992

PURSUIT® SUPPLEMENTAL LABELING

EPA Reg. No. 241-310

FOR USE IN SOYBEANS

PURSUIT AND GRASS HERBICIDES FOR CONTROL OF
VOLUNTEER CORN AND GRASSES



OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT LEAFLET LABEL BEFORE USING. SEE THE PURSUIT LEAFLET LABEL FOR USE AREA, MIXING AND APPLICATION INFORMATION, WEEDS CONTROLLED AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system.

A selective postemergence grass herbicide such as Select¹, Poast Plus², Fusilade 2000³, Assure⁴ or Option⁵ may be mixed with PURSUIT to control volunteer corn or grasses not controlled by PURSUIT. In some cases the activity of the grass herbicide may be reduced when mixed with PURSUIT. The reduction in activity may be overcome by delaying the application of the postemergence grass herbicide 7 days following the application of PURSUIT. Refer to the respective grass herbicide label for recommended application rate, weed size and restrictions.

When PURSUIT is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

This label must be in the possession of the user at the time of herbicide application.

¹Select is a trademark of Valent Corporation

²Poast Plus is a trademark of BASF Corporation

³Fusilade 2000 is a trademark of ICI Americas, Inc.

⁴Assure is a trademark of E. I. duPont de Nemours & Co., Inc.

⁵Option is a trademark of Hoechst-Roussel Agri-Vet Company.

•Registered Trademark of American Cyanamid Company

PE-17148 2/92

BEST AVAILABLE COPY

American Cyanamid Company
Agricultural Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1992


7110 241-310 5071

PURSUIT® HERBICIDE SUPPLEMENTAL LABELING

EPA Reg. No. 241-310

**FOR USE IN SOYBEANS
IN NORTH DAKOTA AND NORTHERN MINNESOTA
(SEE USE AREA BELOW)**

THIS LABEL EXPIRES DECEMBER 31, 1992



OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT LEAFLET LABEL BEFORE USING. SEE THE PURSUIT LEAFLET LABEL FOR ADDITIONAL USE AREAS, MIXING AND APPLICATION INFORMATION, WEEDS CONTROLLED AND ROTATIONAL CROP RESTRICTIONS. THE ROTATIONAL CROP RESTRICTIONS IN THIS SUPPLEMENTAL LABEL SUPERSEDE THE RESTRICTIONS LISTED IN THE LEAFLET LABEL.

GENERAL INFORMATION

PURSUIT Herbicide is a broad spectrum herbicide that provides season long control of many weeds. Under some conditions (such as heavy texture soil, high organic matter, low pH or low rainfall) PURSUIT may cause injury to subsequent planted crops. Sugar beets are particularly sensitive to PURSUIT residues in soil. PURSUIT is not recommended in crop rotations that include sugar beets. See the ROTATIONAL CROP section of this supplemental label for rotations to other sensitive crops.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system.

USE AREA

PURSUIT may be used in North Dakota and in Minnesota north of Highway #210.

BEST AVAILABLE COPY

*Registered Trademark of American Cyanamid Company

6077

APPLICATION RATES

Apply PURSUIT at 3 ounces per acre postemergence only.

WEEDS CONTROLLED

<u>Weeds controlled</u>	<u>Maximum Leaf Stage</u>	<u>(inches)</u>
Nightshade, black	4	1-3
Eastern black hairy	4	1-3
Kochia	4	1-3
Pigweed, redroot	4	1-4
Mustard species	4	1-3

ADJUVANTS

POSTEMERGENCE APPLICATIONS OF PURSUIT REQUIRE THE ADDITION OF AN ADJUVANT AND A LIQUID FERTILIZER SOLUTION.

I. ADJUVANTS

SURFACTANTS: Use a non-ionic surfactant containing at least 80% active ingredient. Apply the surfactant at the rate of 1 qt./100 gal.

OR

CROP OIL CONCENTRATE: Instead of a surfactant, a petroleum or vegetable seed based oil concentrate (such as SUN-IT II[™]) may be used. SUN-IT II is recommended when weeds are under moisture or temperature stress. Use SUN-IT II at the rate of 1.5 to 2 pts./acre (use the higher rate when weeds are at the maximum label size). Other crop oil concentrates should be used at 1.5 pts./A.

AND

II. LIQUID FERTILIZER SOLUTION

Recommended nitrogen based fertilizers include liquid fertilizers (such, as 28%N, 32%N or 10-34-0) at the rate of 1-2 qts./acre. Use the higher rate when weeds are under moisture or temperature stress. Instead of a liquid fertilizer, spray grade ammonium sulfate may be used at the rate of 4 lbs./acre.

BEST AVAILABLE COPY

7/9/11

ROTATIONAL CROPS

The following crops may be planted after applying PURSUIT at the recommended rate in soybeans.

1. Anytime
Soybeans
2. Nine and one-half months after PURSUIT application:
Field Corn
Edible beans and peas
wheat
3. Eighteen months after PURSUIT application:
Barley
Oats
Rye
4. Twenty six months after PURSUIT application:
Canola (rapeseed)
Carrots
Celery
Cole crops
Garlic
Flax
Onions
Potatoes
Spinach
Sunflower
Tomatoes
5. Forty months after PURSUIT application.
Sugar beets

DO NOT plant other rotational crops prior to 18 months after a PURSUIT application.

This label must be in the possession of the user at the time of herbicide application.

PE-17132 1/92

American Cyanamid Company
Agricultural Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©19921