

**FOR USE IN FIELD CORN**  
**(APPLY ONLY ON PURSUIT RESISTANT/TOLERANT**  
**FIELD CORN HYBRIDS)**

**USE DIRECTIONS FOR THE USE OF PURSUIT FOR WEED CONTROL**  
**IN PURSUIT RESISTANT/TOLERANT FIELD CORN HYBRIDS**

**ACCEPTED**  
 DEC 15 1992  
 Under the Federal Insecticide,  
 Fungicide, and Rodenticide Act  
 as amended, for the pesticide  
 registered under  
 EPA Reg. No. 241-310

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT LEAFLET LABEL BEFORE USING. SEE THE PURSUIT LEAFLET LABEL FOR PRECAUTIONS, APPLICATION INFORMATION, USES WITH OTHER PRODUCTS AND ROTATIONAL CROP RESTRICTIONS.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**DO NOT** apply this product through any type of irrigation system.

This label expires December 1993.

**GENERAL INFORMATION**

Apply PURSUIT only on selected field corn hybrids warranted by the seed company to possess resistance/tolerance to direct application of PURSUIT (for example: Pioneer IR varieties). **DO NOT** apply PURSUIT to corn hybrids which lack genetic resistance/tolerance to PURSUIT herbicide. Contact your chemical dealer, seed supplier or American Cyanamid to obtain information regarding PURSUIT resistant/tolerant field corn hybrids.

PURSUIT can be applied as an early preplant, preplant incorporated, preemergence, postemergence or directed-postemergence treatment on field corn hybrids which are classified as PURSUIT resistant/tolerant.

Crops growing under stressful environmental conditions can exhibit various injury symptoms which may be more pronounced if herbicides are used. Corn plants treated with PURSUIT may exhibit yellowing on new growth. Such effects occur infrequently and are temporary. Normal growth and appearance should resume within 1 to 2 weeks.

**Replanting:** If replanting is necessary in a field previously treated with PURSUIT, the field may be replanted to PURSUIT resistant/tolerant field corn, soybeans or peanuts. Rework the soil no deeper than the treated zone. **DO NOT** apply a second treatment of PURSUIT.

<sup>1</sup>Bronco, Roundup, and Lasso are Trademarks of Monsanto Agricultural Products Company.

<sup>2</sup>Gramoxone Extra, Sutan+, and Eradicane are Trademarks of ICI Americas, Inc.

<sup>3</sup>Dual is a Trademark of Ciba Geigy Corporation.

<sup>4</sup>Buctril is a Trademark of Rhone-Poulenc Ag Company.

<sup>5</sup>Banvel and Marksman are Trademarks of Sandoz Crop Protection Corporation.

<sup>6</sup>Registered Trademark of American Cyanamid Company.

**MIXING INSTRUCTIONS**

When tank-mixing PURSUIT with recommended herbicides, add the other herbicide to the spray tank first and make sure it is thoroughly mixed before adding PURSUIT. When tank-mixing atrazine, Bronco<sup>1</sup>, Roundup<sup>1</sup>, Lasso<sup>1</sup>, Gramoxone Extra<sup>2</sup>, Eradicane<sup>2</sup>, Sutan+<sup>2</sup>, Dual<sup>3</sup>, Buctril<sup>4</sup>, Banvel<sup>5</sup> or Marksman<sup>5</sup> with PURSUIT and PROWL<sup>6</sup> herbicide, add the herbicides in the following order: (1) atrazine, (2) Buctril, Banvel, PROWL, Marksman, Eradicane, Sutan+, Lasso or Dual, (3) Bronco, Roundup, Gramoxone Extra, (4) PURSUIT, and (5) nonionic surfactant.

**USE RATE**

**(4 OUNCES PER ACRE)**

Apply PURSUIT at a broadcast rate of 4 ounces per acre (1/4 pint) for all methods of application: early preplant, preplant incorporated, preemergence (including minimum and no-till), postemergence and directed-postemergence. At this broadcast rate, one gallon of PURSUIT will treat 32 acres of PURSUIT resistant/tolerant field corn.

**NOTE:** Only one application of PURSUIT may be made during the growing season.

**APPLICATION INSTRUCTIONS**  
**POSTEMERGENCE**

Apply PURSUIT herbicide as a postemergence treatment to PURSUIT resistant/tolerant field corn when crop and weeds are actively growing. Apply PURSUIT before weeds exceed a height of 3 inches, unless otherwise indicated (see PURSUIT leaflet label) and before corn exceeds the eight-leaf stage. More restrictive crop growth stage limitations of tank-mix partners must be followed.

## I. ADJUVANTS

**SURFACTANTS:** Use a non-ionic surfactant containing at least 80% active ingredient. Apply the surfactant at the rate of 1 qt./100 gal. of spray solution

OR

**CROP OIL CONCENTRATE:** Instead of a surfactant, a petroleum or vegetable seed oil based crop oil concentrate (such as SUN-IT II<sup>®</sup>) may be used. SUN-IT II is recommended when weeds are under moisture or temperature stress. Use SUN-IT II/other crop oil concentrates at the rate of 1.5 to 2 pts./A (use the higher rate when weeds are at the maximum label size).

AND

## II. LIQUID FERTILIZER SOLUTION

Recommended nitrogen based fertilizers include liquid fertilizers (such as 28%N, 32%N or 10-34-0) at the rate of 1-2 qts./A. Use the higher rate when weeds are under moisture or temperature stress. Instead of a liquid fertilizer, spray grade ammonium sulfate may be used at the rate of 2.5 lbs./A.

## SOIL APPLICATIONS

PURSUIT can be applied as an early preplant, preplant incorporated, or preemergence treatment in PURSUIT resistant/tolerant field corn.

## HERBICIDE COMBINATIONS POSTEMERGENCE

Products that can be tank-mixed postemergence with PURSUIT include PROWL, atrazine, Buctril, Banvel and Marksman. Postemergence applications of PURSUIT with PROWL or Banvel can be made to corn up to the five-leaf stage. Atrazine can be tank-mixed with PURSUIT for application to corn up to 12" in height. Buctril can be tank-mixed with PURSUIT for application to corn between the three-leaf and eight-leaf stages. Some corn leaf burn may result with Buctril or atrazine postemergence combinations with PURSUIT. Applications of Banvel herbicide to corn during periods of rapid growth may result in temporary leaning. **DO NOT** use crop oil concentrates as adjuvants in PURSUIT combinations with Banvel. In some cases the grass activity of PURSUIT will be reduced when used in combination with atrazine, Buctril or Banvel.

<sup>\*</sup>PURSUIT applications may be made with a non-ionic surfactant only (liquid fertilizer not required) in the states of Arkansas, Southern Missouri, Tennessee, South Carolina, Louisiana, Mississippi, Alabama, and Georgia.

<sup>†</sup>SUN-IT II is a Trademark of Agcon, Inc.

<sup>‡</sup>Accent is a Trademark of E. I. DuPont de Nemours and Company, Inc.

<sup>§</sup>Beacon is a Trademark of Ciba Geigy Corporation.

to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions.

## HERBICIDE COMBINATIONS SOIL APPLICATIONS

When heavy grass pressure is anticipated, PURSUIT should be used in combination with one of the following registered soil-applied grass herbicides; PROWL (preemergence or early postemergence only, **DO NOT** incorporate PROWL), Lasso or Dual (preemergence or preplant incorporated), Eradicane or Sutan+ (preplant incorporated only).

The application of a soil applied grass herbicide underlay (such as Lasso, Dual, Sutan+, or Eradicane) or PROWL applied preemergence or early postemergence (**DO NOT** incorporate PROWL treatments) will control grass weeds not on the PURSUIT label and enhance the control of certain broadleaf weeds such as common lambsquarters.

Refer to the PROWL (or other grass herbicide) label for specific use recommendations, rates and precautions.

**DO NOT** tank-mix PURSUIT with Accent<sup>®</sup> or Beacon<sup>®</sup>. **DO NOT** apply separate applications of these herbicides before or after PURSUIT treatments.

## Weeds Controlled by Preplant and Preemergence Combination Soil Treatments.

When applied as directed, PURSUIT combination soil treatments with PROWL (Preemergence only, **DO NOT** incorporate PROWL), Lasso, Dual, Eradicane, or Sutan+ will control the weeds listed in the following table, in addition to those controlled by PURSUIT alone.

GRASSES	PROWL <sup>a</sup>	Eradicane <sup>b</sup>	Sutan+ <sup>c</sup>	Lasso <sup>d</sup>	Dual <sup>e</sup>
Barnyardgrass	X	X	X	X	X
Crabgrass, large	X	X	X	X	X
Crabgrass, smooth	X	X	X	X	X
Crowfootgrass	X	X	X		
Goosegrass	X	X	X	X	X
Ichgrass		X	X		
Panicum, fall	X	X	X	X	X
Panicum, Texas	X	X	X		
Rhizome johnsongrass		X	X		
Sandbur, field	X	X	X		
Shattercane	X	X	X		
Signalgrass broadleaf	X	X	X	X	X
Wild proso millet	X	X	X		
Witchgrass	X	X	X	X	X
Woolly Cupgrass	X	X			

<sup>a</sup>Preemergence treatments only. **DO NOT** apply PROWL preplant incorporated. PROWL does not control emerged grasses.

<sup>b</sup>Preplant incorporated treatments only.

<sup>c</sup>Preplant incorporated or preemergence treatments.

**BROADLEAF WEEDS**

**GRASS WEEDS**

Weeds Controlled	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (Inches)
Anoda, spurred	C	2	1-2
Artichoke, Jerusalem		8	6-10
Buffalobur	C		
Bristly starbur		2	1-2
Carpetweed	C		
Cocklebur, common	C**	8	1-8
Galinsoga	C		
Jimsonweed	C	4	1-3
Kochia	C	4	1-3
Lambsquarters, common	C	2	1-2
Mallow, Venice	R		
Marshelder	C	4	1-3
Morningglory,			
entireleaf	R	2	1-2
ivyleaf	R	2	1-2
pitted	R	2	1-2
smallflower	C	4	1-3
tall	R	2	1-2
Mustard sp.	C	4	1-3
Nightshade,			
black	C	4	1-3
Eastern black	C	4	1-3
airy	C	4	1-3
Pigweed,			
palmer	C	8	1-8
redroot	C	8	1-8
smooth	C	8	1-8
spiny	C	8	1-8
Poinsettia, wild	C		
Puncturevine	C		
Purslane, common	C		
Pusley, Florida	C		
Sida, prickly	C		
Ragweed,			
common	R	4	1-3
giant	R	4	1-3
Sage, banyard	R		1-3
Smartweed,			
ladysthumb	C	4	1-3
Pennsylvania	C	4	1-3
Spurge,			
prostrate	C	4	1-3
spotted	C	4	1-3
Sunflower, common	C	4	1-3
Velvetleaf	C	4	1-3
Thistle, Canada	R		1-3
Waterhemp, tall	C	4	1-3

Weeds Controlled	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (Inches)
Barnyardgrass	R	3	1-3
Crabgrass,			
large	R	3	1-3
smooth	R	3	1-3
Cupgrass, woolly		3**	1-3
Foxtail,			
giant	C	6	1-6
green	C	3	1-3
giant green	C	3	1-3
robust purple	C	3	1-3
robust white	C	3	1-3
yellow	C	3	1-3
Goosegrass	R		
Johnsongrass,			
seedling	C	6	1-8
rhizome		R	1-8
Millet, wild proso	R	R	1-3
Panicum,			
tall	R		
Texas	R		
Red rice		3	1-3
Sandbar, field	R	R	≤1
Shattercane	R	6	1-8
Signalgrass, broadleaf	R	4	1-8
Sorghum alnum	R	6	1-3

**SEDGES**

Nutsedge,			
purple	R	R	1-3
yellow	R	R	1-3

\*When soil applied to grasses, more consistent control can be obtained from preplant incorporated treatments.

\*\*PURSUIT control - emerged woolly cupgrass only.

When applied as directed at the broadcast rate of 4 ounces per acre, PURSUIT will control or reduce competition from the weeds listed above.

NOTE: C = Control, R = Reduced Competition

(The number under Maximum Leaf Stage indicates the MAXIMUM number of leaves at which weeds should be sprayed postemergence).

PURSUIT is active against many broadleaf and grass species. However, when heavy grass pressure is anticipated, PURSUIT should be used in combination with a soil-applied grass herbicide. (See HERBICIDE COMBINATIONS section).

\*When soil applied, common lambsquarters, jimsonweed, prickly sida, velvetleaf and common sunflower are more consistently controlled by preplant incorporated treatments.

\*\*Must be preplant incorporated for best results. Cultivation and/or a postemergence herbicide may be required for season long results.

DO NOT count cotyledon leaves when determining weed stage of growth.

40753

## ROTATIONAL CROPS

Refer to the PURSUIT leaflet label for further rotational crop restrictions.

The following rotational crops may be planted after applying PURSUIT at the recommended rate in corn:

1. Anytime
 

Lima beans	Peanuts
Southern peas	Resistant/tolerant field corn
Soybeans	
2. Four months after PURSUIT application:
  - Rye
  - Wheat
  - Edible beans and peas (other than lima beans and Southern peas)
3. Nine and one-half months after PURSUIT application:
 

Barley	Field corn grown for seed*
Field corn	Turkey
4. Eighteen months after PURSUIT application:
 

Alfalfa	Lettuce	Popcorn	Sorghum
Cotton	Oats	Safflower	Sweet corn
5. Twenty six months after PURSUIT application:
  - All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.
6. Forty months after PURSUIT application:
  - Sugarbeets

DO NOT plant other rotational crops prior to 26 months after a PURSUIT application.

\* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT shall be assumed by the user.

### PRECAUTIONS

In the event of a crop loss due to weather, resistant/tolerant field corn hybrids, soybeans, edible beans (lima, dry), southern peas or peanut, can be replanted if PURSUIT was applied alone at rates labelled for these crops. If PURSUIT was tank-mixed with other herbicides, such as atrazine, the label restrictions for these herbicides must also be followed.

There should be an interval of at least 45 days between an application of PURSUIT and corn harvest (silage, fodder, or grain). DO NOT graze or feed treated corn, forage, silage, fodder, or grain for at least 45 days after an application of PURSUIT.

DO NOT apply products containing nicosulfuron (Accent), primisulfuron (Beacon) to field corn the same year as PURSUIT or injury to follow crops may occur.

All soil insecticides, including labelled banded or in-furrow applications, may be used in combination with Pioneer imidazolinone resistant (IR) corn hybrids.

Imidazolinone tolerant hybrids from other seed companies may occasionally exhibit injury symptoms when soil insecticides are used in combination with PURSUIT. USE banded applications of COUNTER® 20CR® or THIMET® in combination with PURSUIT application; DO NOT USE COUNTER soil insecticide products in-furrow with imidazolinone tolerant hybrids. American Cyanamid has not tested all hybrids in which the imidazolinone tolerance trait is claimed and cannot be responsible for factors which are beyond its control, such as growing conditions, environmental conditions, grower practices and the specific genetics of each hybrid which may influence tolerance. Growers should contact the seed company for information regarding specific hybrid tolerance to herbicide and insecticide applications.

This label must be in the possession of the user at the time of herbicide application.

For additional information regarding the use of PURSUIT herbicide, call telephone no. (800) 942-0500.

**EPA Reg. No. 241-310**

**FOR USE IN SUCCULENT PEAS, DRY EDIBLE PEAS,  
LENTILS, AND CHICKPEAS**

PURSUIT can be used on succulent peas, dry edible peas (including chickpeas) and lentils in Washington (Adams, Asotin, Columbia, Garfield, Spokane, Walla Walla and Whitman counties), Idaho (Benewah, Clearwater, Idaho, Kootenai, Latah, Lewis, and Nez Perce counties), Oregon (Union, Umatilla and Wallowa counties) and Montana only.

**THIS LABEL EXPIRES DECEMBER 31, 1993**

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

**GENERAL INFORMATION**

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to peas and lentils. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

**DO NOT** apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application.

**USE PURSUIT ONLY** if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans. Plant peas or lentils at least 1/2 inch deep to reduce risk of crop injury.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

**DISCLAIMER**

**THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.**

**REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.**

\*Registered Trademark of American Cyanamid Company

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

**DO NOT** apply PURSUIT in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT applications must be drained and thoroughly cleaned with water before being used to apply other products.

**DO NOT** make more than one application of PURSUIT per year.

Allow at least 30 days between application and harvest of succulent peas.

Allow at least 60 days between application and harvest of dry edible peas, chickpeas, and lentils.

When PURSUIT is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

(see reverse side)

PURSUIT may be tank-mixed with Sencor DF<sup>1</sup> or Lexone DF<sup>2</sup> to assist in the control of lambsquarters or mayweed-chamomile (dogfennel). Refer to the Sencor or Lexone label for proper application rates and restrictions.

## WEEDS CONTROLLED


PURSUIT applied at the broadcast rate of 3 ounces per acre preplant incorporated or preemergence will control:

Buckwheat, wild	Nightshade, black	Pigweed, redroot
Kochia	Nightshade, Eastern black	Shepherdspurse
Mustard, wild	Nightshade, hairy	Thistle, Russian

## ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT as recommended in this label:

- Anytime  
Chickpeas  
Lentils  
Peas  
Peanuts  
Soybeans
- Four months after PURSUIT application:  
Barley  
Edible beans (dry, lima, snap)  
Rye  
Wheat
- Nine and one-half months after PURSUIT application:  
Field corn  
Field corn grown for seed\*  
Tobacco
- Eighteen months after PURSUIT application:  
Alfalfa  
Cotton  
Lettuce  
Oats  
Popcorn  
Safflower  
Sorghum  
Sweet Corn
- Twenty-six months after PURSUIT application:  
All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.
- Forty months after PURSUIT application:  
Sugar beets



In the event of a crop loss, peas, lentils or chickpeas may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

\*Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into field treated previously with PURSUIT shall be assumed by the user.

CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.

<sup>1</sup>Sencor is a trademark of the Farnam Company of Parkersburg, West Virginia.  
<sup>2</sup>Lexone is a trademark of E. I. duPont de Nemours and Company, Inc.

 **CYANAMID**  
American Cyanamid Company  
Agricultural Division  
Crop Protection Chemicals Department  
Wayne, NJ 07470 ©1992

**EPA Reg. No. 241-310**

**FOR USE IN GREAT NORTHERN, RED KIDNEY AND PINTO  
TYPE DRY BEANS**

**FOR DISTRIBUTION AND USE ONLY IN NEW MEXICO**

**THIS LABEL EXPIRES DECEMBER 31, 1993**

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN  
THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

**GENERAL INFORMATION**

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

**DO NOT** apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application. **DO NOT** apply PURSUIT postemergence after crop has begun to flower or crop injury may result.

**USE PURSUIT ONLY** if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

**DISCLAIMER**

**THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.**

**REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.**

\*Dash is a trademark of BASF Corporation

\*Registered trademark of American Cyanamid Company

(see reverse side)

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

**DO NOT** apply PURSUIT in any manner not specifically described in this label.


Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for post applications.

**DO NOT** use crop oils, petroleum oils, liquid fertilizers or Dash<sup>1</sup>.

**DO NOT** make more than one application of PURSUIT per year. **DO NOT** apply PURSUIT after June 30. Allow at least 60 days between application and harvest.



restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

## APPLICATION RATE AND TIMING

**Preemergence Applications.** Apply PURSUIT at the broadcast rate of 3 ounces per acre immediately after, or up to 1 day after planting. PURSUIT may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide. DO NOT apply PURSUIT on sand or loamy sand soils.

**Postemergence Applications.** Apply PURSUIT at 3 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

**DO NOT APPLY PURSUIT POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.**

## WEEDS CONTROLLED

PURSUIT applied at the broadcast rate of 3 ounces per acre preemergence or early postemergence will control:

Mustard, wild	Nightshade, hairy
Nightshade, black	Pigweed, redroot
Nightshade, Eastern black	

Postemergence applications of 3 ounces per acre must be to weeds less than 2 inches tall for best results.

## ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT as recommended in this label:

1. Anytime
  - Soybeans
  - Peanuts
  - Edible bean types listed on this label (Refer to heading on page 1).

Peas  
Rye  
Wheat

3. Nine and one-half months after PURSUIT application:
  - Field corn
  - Field corn grown for seed\*
  - Tobacco
4. Eighteen months after PURSUIT application:
  - Alfalfa
  - Cotton
  - Lettuce
  - Oats
  - Popcorn
  - Safflower
  - Sorghum
  - Sweet Corn
5. Twenty-six months after PURSUIT application:
  - All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.
6. Forty months after PURSUIT application:
  - Sugar beets

in the event of a crop loss Great Northern, red kidney or pinto beans may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

\* Several seed companies have tested a wide range of in-breds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT shall be assumed by the user.

**CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.**

**EPA Reg. No. 241-310**

**FOR USE IN LIMA BEANS ONLY**

**FOR DISTRIBUTION AND USE IN NEW JERSEY, DELAWARE  
AND MARYLAND ONLY**

**THIS LABEL EXPIRES DECEMBER 31, 1993**

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN  
THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

**GENERAL INFORMATION**

PURSUIT application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT only to the legume vegetables specifically listed in this label. DO NOT apply PURSUIT in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. DO NOT apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

\*Dash is a trademark of BASF Corporation  
\*Registered Trademark of American Cyanamid Company

(see reverse side)

**DISCLAIMER**

**THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.**

**REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.**

**DIRECTIONS FOR USE**


It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply PURSUIT in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for postemergence applications. DO NOT use crop oils, petroleum oils, liquid fertilizers or Dash<sup>1</sup>.



neulous of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

**DO NOT** apply PURSUIT after June 30.

Allow at least 30 days between application and harvest.

**Preplant Incorporated Applications.** Apply PURSUIT at the broadcast rate of 2 ounces per acre to sand and loamy sand soils, or 3 ounces per acre to all other soils within 1 week before planting. Applied preplant incorporated, PURSUIT may be tank-mixed with a registered grass herbicide.

**Preemergence Applications.** Apply PURSUIT at the broadcast rate of 2 ounces per acre to sand and loamy sand soils, or 3 ounces per acre to all other soils immediately after, or up to 1 day after planting. PURSUIT may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

**Postemergence Applications.** Apply PURSUIT at 2 ounces per acre. A nonionic surfactant must be added to the spray solution. A nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

**DO NOT APPLY PURSUIT POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.**

## WEEDS CONTROLLED

PURSUIT applied to low organic matter soils at the broadcast rate of 2 ounces per acre preplant incorporated, preemergence, or postemergence will control light to moderate infestations of:

Nightshade, black  
Nightshade, Eastern black  
Pigweed, redroot

PURSUIT applied at the broadcast rate of 3 ounces per acre preplant incorporated or preemergence will control:

Mustard, wild  
Nightshade, black  
Nightshade, Eastern black  
Nightshade, hairy  
Pigweed, redroot

Soybeans  
Peanuts  
Resistant Field Corn  
Edible bean and pea types listed on this label (Refer to heading on page 1)

2. Two months after PURSUIT application:  
Snap beans (DE, MD, PA and VA only)
3. Four months after PURSUIT application:  
Barley  
Edible bean and pea types not listed on this label (Refer to heading on page 1)  
Rye  
Wheat
4. Nine and one-half months after PURSUIT application:  
Field corn  
Field corn grown for seed\*  
Tobacco
5. Eighteen months after PURSUIT application:  
Alfalfa                      Oats                      Sorghum  
Cotton                      Popcorn                      Sweet Corn  
Lettuce                      Safflower
6. Twenty-six months after PURSUIT application:  
All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.
7. Forty months after PURSUIT application:  
Sugar beets

In the event of a crop loss, the bean or pea crop may be replanted.

**DO NOT** feed treated forage, hay, or straw to livestock.

\* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT shall be assumed by the user.

**CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.**

11/27/92

**PURSUIT® HERBICIDE  
FOR USE IN LIMA BEANS IN NEW JERSEY, DELAWARE AND MARYLAND  
Description of Changes**

This supplemental label is a composite of language approved previously in a supplemental label on March 5, 1992 (exhibit 3) and in the full product label on March 19, 1990 (exhibit 4) with some minor statements added for clarification.

A. This supplemental label is restricted to the use of PURSUIT in lima beans in New Jersey, Delaware and Maryland only (Chart 1). The label includes an expiration date of December 31, 1993.

B. GENERAL INFORMATION, DISCLAIMER AND DIRECTIONS FOR USE

These paragraphs were previously approved in supplemental labeling accepted by EPA on March 5, 1992, and highlighted in exhibit 3, page 1.

In the third paragraph under DIRECTIONS FOR USE, the word "sugarbeets" has been changed to the correctly spelled "sugar beets."

C. APPLICATION INSTRUCTIONS

These paragraphs were previously approved in supplemental labeling accepted by EPA on March 5, 1992, and highlighted in exhibit 3, page 2 for preplant incorporated and preemergence applications.

A 2 ounce postemergence application was added. This application timing is allowed on English peas in the full product label, exhibit 4, page 27.

In addition, the warning that reads, "DO NOT APPLY PURSUIT POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY MAY RESULT" has been expanded. The phrase "(REDUCED CROP GROWTH AND/OR DELAYED MATURITY)" has been inserted after "CROP INJURY" to clarify the type of injury that may occur.

D. WEEDS CONTROLLED

Weeds listed under 2 ounce rates were previously approved in supplemental labeling accepted by EPA on March 5, 1992, and highlighted in exhibit 3, page 3, and in the full label, highlighted in exhibit 4, page 24. Weeds listed under 3 ounce rates were previously approved in supplemental labeling accepted by EPA on March 5, 1992, and highlighted in exhibit 3, page 3.

E. ROTATIONAL CROP GUIDELINE

Some of the language included in the Rotational Crop Guideline was previously approved in supplemental labeling accepted by EPA on March 5, 1992, and highlighted in exhibit 3, page 5.

In addition, "ROTATIONAL CROP RESTRICTIONS" was changed to "ROTATIONAL CROP GUIDELINE" and the rotational crop intervals were made more restrictive. An eighteen-month rotational interval was added allowing cotton, lettuce, oats, popcorn, safflower, sorghum, and sweet corn eighteen months after PURSUIT application. The twenty-six month interval was changed to "All crops not listed

elsewhere in this ROTATIONAL CROP GUIDELINE." A forty-month interval was also added allowing sugar beets to be planted forty months after PURSUIT application. The sentence "DO NOT plant other rotational crops prior to 18 months after a PURSUIT application." was removed.

12 of 32



EPA Reg. No. 241-310

**FOR USE IN GREAT NORTHERN AND RED KIDNEY TYPE DRY BEANS**

**FOR DISTRIBUTION AND USE ONLY IN COLORADO, KANSAS,  
NEBRASKA AND WYOMING**

**PURSUIT MAY BE APPLIED TO THE LISTED CROPS WEST OF U.S. 81 ONLY.**

**THIS LABEL EXPIRES DECEMBER 31, 1993**

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

**GENERAL INFORMATION**

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

**DO NOT** apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application. **DO NOT** apply PURSUIT postemergence after crop has begun to flower or crop injury may result.

**USE PURSUIT ONLY** if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

**DISCLAIMER**

**THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.**

**REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.**

\*Dash is a trademark of BASF Corporation  
\*Registered Trademark of American Cyanamid Company

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

**DO NOT** apply PURSUIT in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for post applications.

**DO NOT** use crop oils, petroleum oils, liquid fertilizers or Dash<sup>1</sup>.

**DO NOT** make more than one application of PURSUIT per year. **DO NOT** apply PURSUIT after June 30. Allow at least 60 days between application and harvest.

(see reverse side)

ACCEPTED  
DEC 15 1993  
241-310

... and pre-emergence. For most usages should be  
... ed. PURSUIT cannot be mixed with any product con-  
... taining a label prohibiting such mixtures.

## APPLICATION RATE AND TIMING

**Pre-emergence Applications.** Apply PURSUIT at the broad-  
... rate of 3 ounces per acre immediately after, or up to 1 day  
... after planting. PURSUIT may be applied in a tank-mix with a  
... registered grass herbicide or applied pre-emergence following a  
... preplant incorporated application of a registered grass herbicide.  
... DO NOT apply PURSUIT on sand or loamy sand soils.

**Postemergence Applications.** Apply PURSUIT at 3 ounces  
... per acre. A nonionic surfactant must be added to the spray  
... solution. The nonionic surfactant must contain at least 80%  
... active ingredient and should be used at a rate of 2 pints per 100  
... gallons of spray mixture.

**DO NOT APPLY PURSUIT POSTEMERGENCE BE-  
... FORE CROP HAS AT LEAST ONE TRUE LEAF OR  
... CROP INJURY (REDUCED CROP GROWTH AND/OR  
... DELAYED MATURITY) MAY RESULT.**

## WEEDS CONTROLLED

PURSUIT applied at the broadcast rate of 3 ounces per acre  
... pre-emergence or early post-emergence will control:

Mustard, wild	Nightshade, hairy
Nightshade, black	Pigweed, redroot
Nightshade, Eastern black	

Post-emergence applications of 3 ounces per acre must be to  
... weeds less than 2 inches tall for best results.

## ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT  
... as recommended in this label:

1. Anytime  
Soybeans  
Peanuts  
Edible bean types listed on this label (Refer to heading  
... on page 1).

PE-17151 Rev. 1 11/12/92 SD

Rye  
Wheat

3. Nine and one-half months after PURSUIT application:  
Field corn  
Field corn, grown for seed\*  
Tobacco
4. Eighteen months after PURSUIT application:  
Alfalfa  
Cotton  
Lettuce  
Oats  
Popcorn  
Safflower  
Sorghum  
Sweet Corn
5. Twenty-six months after PURSUIT application:  
All crops not listed in this ROTATIONAL CROP  
GUIDELINE.
6. Forty months after PURSUIT application:  
Sugar beets

In the event of a crop loss Great Northern or red kidney beans  
... may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

\* Several seed companies have tested a wide range of in-  
... bred for sensitivity to PURSUIT soil residues and have  
... reported good crop safety. However, due to the proprie-  
... tary nature of seed production, American Cyanamid has  
... not been given access to the inbred data. Growers are  
... directed to contact the seed company for information and  
... recommendations regarding the planting of corn grown  
... for seed in fields treated with PURSUIT the previous  
... year. Since growing conditions, environmental condi-  
... tions and grower practices are beyond the control of  
... American Cyanamid Company, all risks and consequences  
... associated with planting seed corn inbreds into fields  
... treated previously with PURSUIT shall be assumed by  
... the user.

CONSULT THE PURSUIT LEAFLET LABEL FOR  
... OTHER PRECAUTIONS.

 **CYANAMID**  
American Cyanamid Company  
Agricultural Division  
Crop Protection Chemicals Department  
Wayne, NJ 07470 ©1992

**EPA Reg. No. 241-310**

**FOR PREEMERGENCE USE IN SNAP BEANS GROWN  
IN NEW JERSEY ONLY**

**THIS LABEL EXPIRES DECEMBER 31, 1993**

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN  
THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

**GENERAL INFORMATION**

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to snap beans. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

**DO NOT** apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application.

**USE PURSUIT ONLY** if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

**DISCLAIMER**

**THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.**

**REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.**

\*Registered Trademark of American Cyanamid Company

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

**DO NOT** apply PURSUIT in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT applications must be drained and thoroughly cleaned with water before being used to apply other products.


**DO NOT** apply PURSUIT after June 20.

Allow at least 30 days between application and harvest.

**DO NOT** make more than one application of PURSUIT per year.

When PURSUIT is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

(see reverse side)



**Preemergence Applications.** For suppression of pigweed, nightshade and common purslane, apply PURSUIT at the broadcast rate of 1.5 ounces per acre immediately after, or up to 1 day after planting. PURSUIT may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

## ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT as recommended in this label:

1. Anytime
  - Soybeans
  - Peanuts
  - Edible beans and peas
2. Four months after PURSUIT application:
  - Barley
  - Rye
  - Wheat
3. Nine and one-half months after PURSUIT application:
  - Field corn
  - Field corn grown for seed\*
  - Tobacco
4. Eighteen months after PURSUIT application:

Alfalfa	Popcorn
Cotton	Safflower
Lettuce	Sorghum
Oats	Sweet Corn
5. Twenty-six months after PURSUIT application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.
6. Forty months after PURSUIT application:
  - Sugar beets

In the event of a crop loss, snap beans may be replanted.

**DO NOT** feed treated forage, hay, or straw to livestock.

\* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions, and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT shall be assumed by the user.

**CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.**

PE-17005 Rev. 1 11/13/92 SD

 **CYANAMID**  
American Cyanamid Company  
Agricultural Division  
Crop Protection Chemicals Department  
Wayne, NJ 07470 ©1992

1807-22

**PURSUIT<sup>®</sup> HERBICIDE**  
**MINOR CHANGES SPECIFIC TO THE REGION 1 SUPPLEMENTAL LABEL**

- A. Under USE AREA MAP, the Delmarva peninsula of Delaware, Maryland and Virginia has been excluded from the shaded area.
  
- B. The sentence under the map has been changed to read: "Use area includes: South Dakota east of U.S. 81, Nebraska east of U.S. 81, Kansas east of U.S. 81, Maryland and Virginia excluding the Delmarva peninsula, and all other states or portions of states in the preceding map."

1807-22

**EPA Reg. No. 241-310**

**FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE, CRANBERRY, AND SMALL WHITE TYPE DRY BEANS, LIMA BEANS, AND SOUTHERN AND ENGLISH PEAS ONLY**

**REFER TO THE MAP ON PAGE 6 FOR GEOGRAPHICAL USE AREA  
READ LABEL CAREFULLY - USE INSTRUCTIONS VARY FOR EACH GROUP  
THIS LABEL EXPIRES DECEMBER 31, 1993**

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

**GENERAL INFORMATION**

PURSUIT application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

**DO NOT** apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application.

**DO NOT** apply PURSUIT postemergence after crop has begun to flower or crop injury may result.

**USE PURSUIT ONLY** if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

**DISCLAIMER**

**THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.**

**REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.**

**DIRECTIONS FOR USE**


It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

**DO NOT** apply PURSUIT in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for postemergence applications. **DO NOT** use crop oils, petroleum oils, liquid fertilizers or Dash<sup>1</sup>.



<sup>1</sup>Dash is a trademark of BASF Corporation  
\*Registered Trademark of American Cyanamid Company

with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

## LIMA BEANS

**DO NOT** apply PURSUIT after June 30.

Allow at least 30 days between application and harvest.

**Preplant Incorporated Applications.** Apply PURSUIT at the broadcast rate of 2 ounces per acre to sand and loamy sand soils, or 3 ounces per acre to all other soils within 1 week before planting. Applied preplant incorporated, PURSUIT may be tank-mixed with a registered grass herbicide.

**Preemergence Applications.** Apply PURSUIT at the broadcast rate of 2 ounces per acre to sand and loamy sand soils, or 3 ounces per acre to all other soils immediately after, or up to 1 day after planting. PURSUIT may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

### NAVY, GREAT NORTHERN, RED KIDNEY, CRANBERRY, BLACK TURTLE, AND SMALL WHITE TYPE DRY BEANS

**DO NOT** apply to Domino variety black turtle beans.

**DO NOT** apply PURSUIT to sand or loamy sand soils.

**DO NOT** apply PURSUIT after June 20.

Allow at least 60 days between application and harvest.

**Preplant Incorporated Applications.** Apply PURSUIT within 1 week before planting. Applied preplant incorporated, PURSUIT may be tank-mixed with a registered grass herbicide. Refer to the following table for use rate.

Soil Type	Use Rate (oz/acre)
Sands, loamy sands	DO NOT USE
Sandy loams and all other soils with $\leq$ 2% O.M.	2
All other soils with $>$ 2% O.M.	3

## ENGLISH PEAS

**DO NOT** apply PURSUIT to sand or loamy sand soils.

Allow at least 30 days between application and harvest.

**Preplant Incorporated Applications.** Apply PURSUIT at the broadcast rate of 3 ounces per acre up to 1 week before planting. **DO NOT** apply PURSUIT to English peas that have been or will be treated with trifluralin.

**Preemergence Applications.** Apply PURSUIT at the broadcast rate of 3 ounces per acre immediately after, or up to 1 day after planting. **DO NOT** apply PURSUIT if English peas have been treated with trifluralin.

**Postemergence Applications.** Apply PURSUIT at 3 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture. **DO NOT APPLY PURSUIT POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.** If English peas have been treated with trifluralin, PURSUIT may be applied ONLY at 2 ounces per acre for suppression of black nightshade. The use of trifluralin prior to PURSUIT application may increase the likelihood and severity of crop injury.

## SOUTHERN PEAS

Allow at least 30 days between application and harvest.

**Preplant Incorporated Applications.** Apply PURSUIT at the broadcast rate of 4 ounces per acre within 1 week before planting. Applied preplant incorporated, PURSUIT may be tank-mixed with a registered grass herbicide.

**Preemergence Applications.** Apply PURSUIT at the broadcast rate of 4 ounces per acre immediately after, or up to 1 day after planting. PURSUIT may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

**Early Postemergence Applications.** Apply PURSUIT at the broadcast rate of 4 ounces per acre when peas are at least 3 inches in height. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

**WEEDS CONTROLLED**

**PURSUIT applied to low organic matter soils at the broadcast rate of 2 ounces per acre preplant incorporated or preemergence will control light to moderate infestations of:**

- Nightshade, black
- Nightshade, Eastern black
- Pigweed, redroot

**PURSUIT applied at the broadcast rate of 3 ounces per acre preplant incorporated, preemergence or early post-emergence will control:**

- Mustard, wild
- Nightshade, black
- Nightshade, Eastern black
- Nightshade, hairy
- Pigweed, redroot

**Postemergence applications of 3 ounces per acre must be made to weeds less than 2 inches tall for best results.**

When applied as directed at the broadcast rate of 4 ounces per acre, PURSUIT will control or reduce competition from the weeds listed below:

NOTE: C = Control, R = Reduced Competition

(The number under Maximum Leaf Stage indicates the MAXIMUM number of leaves at which weeds should be sprayed post-emergence.)

**BROADLEAF WEEDS**

WEEDS CONTROLLED	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Anoda, spurred	C	2	1 - 2
Artichoke, Jerusalem		8	6 - 10
Buffalobur	C**		
Bristly starbur		2	1 - 2
Carpetweed	C		
Cocklebur, common	C**	8	1 - 8
Galinsoga	C		
Jimsonweed	C*	4	1 - 3
Kochia	C	4	1 - 3
Lambsquarters	C*	2	1 - 2
Mallow, Venice	R		
Morningglory,			
entireleaf	R	2	1 - 2
ivyleaf	R	2	1 - 2
pitted	R	2	1 - 2
smallflower	C	4	1 - 3
tall	R	2	1 - 2
Mustard sp.	C	4	1 - 3
Nightshade,			
black	C	4	1 - 3
Eastern black	C	4	1 - 3
hairy	C	4	1 - 3
Pigweed,			
palmer	C	8	1 - 8
redroot	C	8	1 - 8
smooth	C	8	1 - 8
spiny	C	8	1 - 8
Poinsettia, wild	C		
Puncturevine	C		
Purslane, common	C		
Pusley, Florida	C		
Sida, prickly	C*		

**BROADLEAF WEEDS (Continued)**

WEEDS CONTROLLED	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Ragweed,			
common	R	4	1 - 3
giant	R	4	1 - 3
Sage, barnyard		R	1 - 3
Smartweed,			
ladysthumb	C	4	1 - 3
Pennsylvania	C	4	1 - 3
Spurge,			
prostrate	C	4	1 - 3
spotted	C	4	1 - 3
flower, common	C*	4	1 - 3
velvetleaf	C*	4	1 - 3
Thistle, Canada		R	1 - 3
Waterhemp, tall	C	4	1 - 3

\*When soil applied, common lambsquarters, jimsonweed, prickly sida, velvetleaf and common sunflower are more consistently controlled by preplant incorporated treatments.

\*\*Use soil applications for light to moderate infestations only. Must be preplant incorporated for best results.

**GRASS WEEDS\***

WEEDS CONTROLLED	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Barnyardgrass	R	3	1 - 3
Crabgrass,			
large	R	3	1 - 3
smooth	R	3	1 - 3
Cupgrass, woolly		3**	1 - 3
Foxtail,			
giant	C	6	1 - 6
green	C	3	1 - 3
giant green	C	3	1 - 3
robust purple	C	3	1 - 3
robust white	C	3	1 - 3
yellow	C	3	1 - 3
Goosegrass	R		
Johnsongrass,			
seedling	C	6	1 - 8
rhizome		R	1 - 8
Panicum,			
fall	R		
Texas	R		

\*When soil applied to grasses, more consistent control can be obtained from preplant incorporated treatments.

\*\*PURSUIT controls emerged woolly cupgrass only.

GRASS WEEDS (Continued)

WEEDS CONTROLLED	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Red rice		3	1 - 3
Shattercane	R	6	1 - 8
Signalgrass, broadleaf	R	4	1 - 8
<b>SEDGES</b>			
Nutsedge,			
purple	R	R	1 - 3
yellow	R	R	1 - 3

DO NOT count cotyledon leaves when determining weed stage of growth.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT as recommended in this label:

1. Anytime
  - Soybeans
  - Peanuts
  - Resistant Field Corn
  - Edible bean and pea types listed on this label (Refer to heading on page 1)
2. Two months after PURSUIT application:
  - Snap beans (DE, MD, PA and VA only)
3. Four months after PURSUIT application:
  - Barley
  - Edible bean and pea types not listed on this label (Refer to heading on page 1)
  - Rye
  - Wheat
4. Nine and one-half months after PURSUIT application:
  - Field corn
  - Field corn grown for seed\*
  - Tobacco
5. Eighteen months after PURSUIT application:
 

Alfalfa	Popcorn
Cotton	Safflower
Lettuce	Sorghum
Oats	Sweet Corn
6. Twenty-six months after PURSUIT application:
  - All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.
7. Forty months after PURSUIT application:
  - Sugar beets

\* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT shall be assumed by the user.

### ROTATIONAL CROP GUIDELINE (Continued)


In the event of a crop loss, the bean or pea crop may be replanted.

**DO NOT** feed treated forage, hay, or straw to livestock.

**CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.**


### USE AREA MAP

PURSUIT can be applied to the edible legume vegetables listed only in the states or parts of states shaded in the following map:



Use area includes: South Dakota east of U.S. 81, Nebraska east of U.S. 81, Kansas east of U.S. 81, Maryland and Virginia excluding the De'marva peninsula, and all other states or portions of states shaded in the preceding map.

FB-17087 Rev. 1 10/12/92 SD



25 of 2

**PURSUIT® HERBICIDE  
FOR USE IN SOUTH DAKOTA (REGION 7)**

This supplemental label is a composite of language approved previously in supplemental labels on March 5, 1992 (exhibit 11) with one minor statement added for clarification.

A. This supplemental label is restricted to the use of PURSUIT in navy, great northern, red kidney, black turtle, cranberry and small white type dry beans, lima beans, dry edible peas, lentils and chickpeas in South Dakota.

**B. GENERAL INFORMATION, DISCLAIMER AND DIRECTIONS FOR USE**

These paragraphs were previously approved in supplemental labeling accepted by EPA on March 5, 1992, and highlighted in exhibit 11, page 1.

The following more restrictive statement was added under DIRECTIONS FOR USE: "Edible beans must receive a minimum of twelve inches combined rainfall and irrigation between planting and harvest."

**C. APPLICATION INSTRUCTIONS**



These paragraphs were previously approved in supplemental labeling accepted by EPA on March 5, 1992, and highlighted in exhibit 11, page 2 for navy, great northern, red kidney, black turtle, cranberry, and small white type dry beans and lima beans; and page 12 for dry edible peas, lentils and chickpeas.

**D. WEEDS CONTROLLED**

Weeds listed here were previously approved in supplemental labeling accepted by EPA on March 5, 1992, and highlighted in exhibit 11, page 3.

**E. ROTATIONAL CROP GUIDELINE**

Some of the language included in the Rotational Crop Guideline was previously approved in supplemental labeling accepted by EPA on March 5, 1992, and highlighted in exhibit 11, page 5. Other minor changes outlined in exhibit 1 have also been incorporated into this supplemental label.



EPA Reg. No. 241-310

**FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE, CRANBERRY  
AND SMALL WHITE TYPE DRY BEANS, LIMA BEANS,  
DRY EDIBLE PEAS, LENTILS, AND CHICKPEAS**

**FOR DISTRIBUTION AND USE ONLY WITHIN SOUTH DAKOTA**

**PURSUIT may be applied to the listed crops west of U.S. 81 only**

**THIS LABEL EXPIRES DECEMBER 31, 1993**

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN  
THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

### **GENERAL INFORMATION**

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

**DO NOT** apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application.

**USE PURSUIT ONLY** if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

### **DISCLAIMER**

**THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.**

**REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.**

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

**DO NOT** apply PURSUIT in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT applications must be drained and thoroughly cleaned with water before being used to apply other products.

**Edible beans must receive a minimum of twelve inches combined rainfall and irrigation between planting and harvest.**

**DO NOT** apply to Domino variety black turtle beans.

**DO NOT** make more than one application of PURSUIT per year.

Allow at least 30 days between application and harvest of lima beans. Allow at least 60 days between application and harvest of dry beans, dry edible peas, lentils and chickpeas.

When PURSUIT is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

\*Registered Trademark of American Cyanamid Company

the PURSUIT leaflet label for incorporation instructions.

**Preemergence Applications.** Apply PURSUIT at the broadcast rate of 3 ounces per acre after planting, but prior to crop emergence. If weeds are present at the time of application, a burndown herbicide such as Roundup<sup>1</sup> or Gramoxone Extra<sup>2</sup> should be tank-mixed with PURSUIT to control emerged weeds.

## WEEDS CONTROLLED

PURSUIT applied at the broadcast rate of 3 ounces per acre preplant incorporated or preemergence will control:

Mustard, wild	Nightshade, hairy
Nightshade, black	Pigweed, redroot
Nightshade, Eastern black	

## ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT as recommended in this label:


1. Anytime
  - Chickpeas
  - Lentils
  - Dry edible peas
  - Peanuts
  - Edible beans (dry and lima listed on this label)
  - Soybeans
2. Four months after PURSUIT application:
  - Barley
  - Edible beans (dry, snap) and peas not listed on this label
  - Rye
  - Wheat
3. Nine and one-half months after PURSUIT application:
  - Field corn
  - Field corn grown for seed\*
  - Tobacco
4. Eighteen months after PURSUIT application:

Alfalfa	Oats	Sorghum
Cotton	Popcorn	Sweet Corn
Lettuce	Safflower	
5. Twenty-six months after PURSUIT application:
  - All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.

\*Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT shall be assumed by the user.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

**CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.**



<sup>1</sup>Roundup is a trademark of Monsanto Agricultural Products Company  
<sup>2</sup>Gramoxone Extra is a trademark of ICI Americas, Inc.



**PURSUIT® HERBICIDE  
FOR USE IN MICHIGAN (REGION 8)**

This supplemental label is a composite of language approved previously in supplemental labels on March 5, 1992 (exhibit 11) and in the full product label on March 16, 1990 (exhibit 12).

A. This supplemental label is restricted to the use of PURSUIT in navy, great northern, red kidney, black turtle, cranberry and small white type dry beans in Michigan.

B. GENERAL INFORMATION, DISCLAIMER AND DIRECTIONS FOR USE  
These paragraphs were previously approved in supplemental labeling accepted by EPA on March 5, 1992, and highlighted in exhibit 11, page 1.


C. APPLICATION INSTRUCTIONS  
These paragraphs were previously approved in supplemental labeling accepted by EPA on March 5, 1992, and highlighted in exhibit 11, page 2 for preplant incorporated and preemergence applications.

Postemergence applications were previously approved in the full label and highlighted in exhibit 12, page 26.

D. WEEDS CONTROLLED  
Weeds listed under 2 ounce rates were previously approved in supplemental labeling accepted by EPA on March 5, 1992, and highlighted in exhibit 11, page 3 and in the full label, highlighted in exhibit 12, page 24. Weeds listed under 3 ounce rates were previously approved in supplemental labeling accepted by EPA on March 5, 1992, and highlighted in exhibit 11, page 3.

E. ROTATIONAL CROP GUIDELINE  
Some of the language included in the Rotational Crop Guideline was previously approved in supplemental labeling accepted by EPA on March 5, 1992, and highlighted in exhibit 11, page 5. Other minor changes outlined in exhibit 1 have also been incorporated into this supplemental label.

**ACCEPTED**  
MAY 15 1992  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 241-310



**EPA Reg. No. 241-310**

**FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE,  
CRANBERRY, AND SMALL WHITE TYPE DRY BEANS ONLY**

**FOR DISTRIBUTION AND USE IN MICHIGAN ONLY  
(EXCLUDING UPPER PENINSULA)**

**THIS LABEL EXPIRES DECEMBER 31, 1993**

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN  
THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

#### **GENERAL INFORMATION**

PURSUIT application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

**DO NOT** apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application.

**DO NOT** apply PURSUIT postemergence after crop has begun to flower or crop injury may result.

**USE PURSUIT ONLY** if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

#### **DISCLAIMER**

**THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.**

<sup>1</sup>Dash is a trademark of BASF Corporation

\*Registered Trademark of American Cyanamid Company

**REFER TO THE PURSUIT LEAFLET LABEL FOR  
ADDITIONAL DISCLAIMER STATEMENTS.**

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

**DO NOT** apply PURSUIT in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for postemergence applications. **DO NOT** use crop oils, petroleum oils, liquid fertilizers or Dash<sup>1</sup>.

**DO NOT** make more than one application of PURSUIT per year.

Refer to the PURSUIT leaflet label for approved combination treatments. When PURSUIT is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

(See reverse side)

**Preplant Incorporated Applications.** Apply PURSUIT within 1 week before planting. Applied preplant incorporated, PURSUIT may be tank-mixed with a registered grass herbicide. Refer to the following table for use rate.

Soil Type	Use Rate (oz/acre)
Sands, loamy sands	DO NOT USE
Sandy loams and all other soils with $\leq$ 2% O.M.	2
All other soils with $>$ 2% O.M.	3

**Preemergence Applications.** Apply PURSUIT at the rates shown in the preceding table immediately after, or up to 1 day after planting. PURSUIT may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

**Postemergence Applications.** Apply PURSUIT at 2 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

**DO NOT APPLY PURSUIT POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.**

### WEEDS CONTROLLED

PURSUIT applied to low organic matter soils at the broadcast rate of 2 ounces per acre preplant incorporated, preemergence, or postemergence will control light to moderate infestations of:

Nightshade, black  
Nightshade, Eastern black  
Pigweed, redroot

PURSUIT applied at the broadcast rate of 3 ounces per acre preplant incorporated or preemergence will control:

Mustard, wild  
Nightshade, black  
Nightshade, Eastern black  
Nightshade, hairy  
Pigweed, redroot

**Resistant Field Corn**  
Edible bean and pea types listed on this label (Refer to heading on page 1)

- Two months after PURSUIT application:  
Snap beans (DE, MD, PA and VA only)
- Four months after PURSUIT application:  
Barley  
Edible bean and pea types not listed on this label (Refer to heading on page 1)  
Rye  
Wheat
- Nine and one-half months after PURSUIT application:  
Field corn  
Field corn grown for seed\*  
Tobacco
- Eighteen months after PURSUIT application:  
Alfalfa                      Oats                      Sorghum  
Cotton                      Popcorn                      Sweet Corn  
Lettuce                      Safflower
- Twenty-six months after PURSUIT application:  
All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.
- Forty months after PURSUIT application:  
Sugar beets

\* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT shall be assumed by the user.


**CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.**

3/10/92

**PURSUIT® HERBICIDE  
FOR USE IN WISCONSIN (REGION 9)**

This supplemental label is a composite of language previously approved in supplemental labels accepted by EPA on March 5, 1992 (exhibit 11).

- A. This supplemental label is restricted to the use of PURSUIT in navy, great northern, red kidney, black turtle, cranberry and small white type dry beans, lima beans, southern and English peas, and white lupins in Wisconsin only, and addresses only a rotational crop interval for snap beans.
  
- B. GENERAL INFORMATION, DISCLAIMER AND DIRECTIONS FOR USE  
These paragraphs were previously approved in supplemental labeling accepted by EPA on March 5, 1992, and highlighted in exhibit 11, page 1.
  
- C. ROTATIONAL CROP GUIDELINE  
The two-month rotational crop interval for snap beans was previously approved in supplemental labeling accepted by EPA on March 5, 1992, and highlighted in exhibit 11, page 5. Other minor changes outlined in exhibit 1 have also been incorporated into this supplemental label.



EPA Reg. No. 241-310

**FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE,  
CRANBERRY, SMALL WHITE TYPE DRY BEANS, LIMA BEANS,  
SOUTHERN AND ENGLISH PEAS, AND WHITE LUPINS**

**FOR DISTRIBUTION AND USE IN WISCONSIN ONLY**

**THIS LABEL EXPIRES DECEMBER 31, 1993**

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN  
THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**GENERAL INFORMATION**

PURSUIT application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

**DO NOT** apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application.

**DO NOT** apply PURSUIT postemergence after crop has begun to flower or crop injury may result.

**USE PURSUIT ONLY** if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

**DISCLAIMER**

**THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.**

**REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.**

**ROTATIONAL CROP GUIDELINE**

The following crops may be planted after applying PURSUIT as recommended:

1. Two months after PURSUIT application:

Snap beans

PE. 11/12/92 SD

 **CYANAMID**  
American Cyanamid Company  
Agricultural Division  
Crop Protection Chemicals Department  
Wayne, NJ 07470 ©1992

