

PM 25- 241-310

1 of 32

**PURSUIT® HERBICIDE SUPPLEMENTAL LABELING**

EPA Reg. No. 241-310

FOR USE IN SOYBEANS

**TANK-MIXTURE OF PURSUIT® + PINNACLE<sup>1</sup> HERBICIDES  
FOR POSTEMERGENCE CONTROL OF COMMON LAMBSQUARTERS IN SOYBEANS**

THIS LABEL EXPIRES DECEMBER 31, 1993

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT LEAFLET LABEL BEFORE USING. SEE THE PURSUIT LEAFLET LABEL FOR MIXING AND APPLICATION INFORMATION, WEEDS CONTROLLED AND ROTATIONAL CROP RESTRICTIONS.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system.

**GENERAL INFORMATION**

For optimal weed control management, apply a soil applied grass herbicide such as PROWL®, TRI-4® or Treflan<sup>2</sup> followed by PURSUIT postemergence. If common lambsquarters are not adequately controlled by the soil applied treatment, Pinnacle herbicide may be tank-mixed with PURSUIT for additional activity.

The addition of Pinnacle to PURSUIT may cause severe injury and/or stunting to soybeans. The USER ASSUMES ALL RISKS AND CONSEQUENCES associated with applications of this tank mixture to soybeans.

**ACCEPTED**  
~~with COMMENTS~~  
in EPA Letter Dated:

11/30/93

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, this product is registered under EPA Reg. No.

(See reverse side)

241-310

\*Registered Trademarks of American Cyanamid Company

<sup>1</sup>Trademark of E. I. duPont de Nemours and Company

<sup>2</sup>Trademark of DowElanco Company

2 of 2

### APPLICATION RATES

When tank-mixing Pinnacle with PURSUIT, use the following rates:

PURSUIT - 4 ounces per acre

Pinnacle - 0.125 ounce per acre

Add to the spray mixture:

Non-ionic surfactant - 1 quart per 100 gallons (0.25% v/v)

AND

Liquid nitrogen based fertilizer (such as 28%N, 32%N, or 10-34-0) at the rate of 1-2 quarts per acre.

NOTE: DO NOT include a crop oil concentrate in this mixture.

### WEEDS CONTROLLED

This tank mixture will improve the activity of PURSUIT on common lambsquarters. Refer to the PURSUIT label for additional weeds controlled.

### RESTRICTIONS

Apply to 1-3 trifoliolate stage soybeans only.


Refer to both labels for information on methods of application, additional restrictions, precautions and rotational crop guidelines. Follow the more restrictive label.

This label must be in the possession of the user at the time of herbicide application.

PE- 12/92

American Cyanamid Company  
Agricultural Division  
Crop Protection Chemicals Department  
Wayne, NJ 07470 ©1992

2 of 30



**PURSUIT® HERBICIDE SUPPLEMENTAL LABELING**

EPA Reg. No. 241-310

FOR USE IN SOYBEANS

**TANK-MIXTURE OF PURSUIT® + PINNACLE<sup>1</sup> HERBICIDES  
FOR POSTEMERGENCE CONTROL OF COMMON LAMBSQUARTERS IN SOYBEANS**

**FOR USE ONLY IN THE STATES OF MINNESOTA AND MICHIGAN  
(SOUTH OF HIGHWAY 210 IN MINNESOTA)**

THIS LABEL EXPIRES DECEMBER 31, 1993

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT LEAFLET LABEL BEFORE USING. SEE THE PURSUIT LEAFLET LABEL FOR MIXING AND APPLICATION INFORMATION, WEEDS CONTROLLED AND ROTATIONAL CROP RESTRICTIONS.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system.

**GENERAL INFORMATION**

For optimal weed control management, apply a soil applied grass herbicide such as PROWL®, TRI-4® or Treflan<sup>2</sup> followed by PURSUIT postemergence. If common lambsquarters are not adequately controlled by the soil applied treatment, Pinnacle herbicide may be tank-mixed with PURSUIT for additional activity.

The addition of Pinnacle to PURSUIT may cause severe injury and/or stunting to soybeans. The USER ASSUMES ALL RISKS AND CONSEQUENCES associated with applications of this tank mixture to soybeans.

(See reverse side)

®Registered Trademarks of American Cyanamid Company

<sup>1</sup>Trademark of E. I. duPont de Nemours and Company

<sup>2</sup>Trademark of DowElanco Company

4 of 30

### APPLICATION RATES

When tank-mixing Pinnacle with PURSUIT, use the following rates:

PURSUIT - 4 ounces per acre

Pinnacle - 0.125 ounce per acre

Add to the spray mixture:

Non-ionic surfactant - 1 quart per 100 gallons (0.25% v/v)

AND

Liquid nitrogen based fertilizer (such as 28%N, 32%N, or 10-34-0) at the rate of 1-2 quarts per acre.

NOTE: DO NOT include a crop oil concentrate in this mixture.

### WEEDS CONTROLLED

This tank mixture will improve the activity of PURSUIT on common lambsquarters. Refer to the PURSUIT label for additional weeds controlled.

### RESTRICTIONS

Apply to 1-3 trifoliolate stage soybeans only.


Refer to both labels for information on methods of application, additional restrictions, precautions and rotational crop guidelines. Follow the more restrictive label.

This label must be in the possession of the user at the time of herbicide application.

PE- 11/92

American Cyanamid Company  
Agricultural Division  
Crop Protection Chemicals Department  
Wayne, NJ 07470 ©1992

5 of 32



**PURSUIT® HERBICIDE SUPPLEMENTAL LABELING**

EPA Reg. No. 241-310

**FOR USE IN SOYBEANS  
 IN NORTH DAKOTA AND NORTHERN MINNESOTA  
 (SEE USE AREA BELOW)**

**THIS LABEL EXPIRES DECEMBER 31, 1993**

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT LEAFLET LABEL BEFORE USING. SEE THE PURSUIT LEAFLET LABEL FOR ADDITIONAL USE AREAS, MIXING AND APPLICATION INFORMATION, WEEDS CONTROLLED AND ROTATIONAL CROP RESTRICTIONS. THE ROTATIONAL CROP RESTRICTIONS IN THIS SUPPLEMENTAL LABEL SUPERSEDE THE RESTRICTIONS LISTED IN THE LEAFLET LABEL.

**GENERAL INFORMATION**

PURSUIT herbicide is a broad spectrum herbicide that provides season long control of many weeds. Under some conditions (such as heavy texture soil, high organic matter, low pH or low rainfall) PURSUIT may cause injury to subsequent planted crops. Sugar beets are particularly sensitive to PURSUIT residues in soil. PURSUIT is not recommended in crop rotations that include sugar beets. See the ROTATIONAL CROP section of this supplemental label for rotations to other sensitive crops.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system.

**USE AREA**

PURSUIT may be used in North Dakota and in Minnesota north of Highway #210.

\*Registered Trademark of American Cyanamid Company

APPLICATION RATES

Apply PURSUIT at 3 ounces per acre postemergence only.

WEEDS CONTROLLED

<u>Weeds controlled</u>	<u>Maximum Leaf Stage</u>	<u>(inches)</u>
Nightshade, black	4	1-3
Eastern black hairy	4	1-3
Kochia	4	1-3
Pigweed, redroot	4	1-4
Mustard species	4	1-3

POSTEMERGENCE APPLICATIONS OF PURSUIT REQUIRE THE ADDITION OF AN ADJUVANT AND A LIQUID FERTILIZER SOLUTION.

I. ADJUVANTS

SURFACTANTS: Use a non-ionic surfactant containing at least 80% active ingredient. Apply the surfactant at the rate of 1 qt./100 gal.

OR

CROP OIL CONCENTRATE: Instead of a surfactant, a petroleum or vegetable seed based oil concentrate (such as SUN-IT II™) may be used. SUN-IT II is recommended when weeds are under moisture or temperature stress. Use SUN-IT II at the rate of 1.5 to 2 pts./acre (use the higher rate when weeds are at the maximum label size). Other crop oil concentrates should be used at 1.5 pts./A.

AND

II. LIQUID FERTILIZER SOLUTION

Recommended nitrogen based fertilizers include liquid fertilizers (such as 26%N, 32%N or 10-34-0) at the rate of 1-2 qts./acre. Use the higher rate when weeds are under moisture or temperature stress. Instead of a liquid fertilizer, spray grade ammonium sulfate may be used at the rate of 4 lbs./acre.

70732

### ROTATIONAL CROPS

The following crops may be planted after applying PURSUIT at the recommended rate in soybeans.

1. Anytime  
Soybeans
2. Nine and one-half months after PURSUIT application:  
Field Corn  
Edible beans and peas  
wheat
3. Eighteen months after PURSUIT application:  
Barley  
Oats  
Rye
4. Twenty six months after PURSUIT application:  
Canola (rapeseed)  
Carrots  
Celery  
Cole crops  
Garlic  
Flax  
Onions  
Potatoes  
Spinach  
Sunflower  
Tomatoes
5. Forty months after PURSUIT application.  
Sugar beets

DO NOT plant other rotational crops prior to 18 months after a PURSUIT application.


This label must be in the possession of the user at the time of herbicide application.

PE-17132 Rev.1 11/92

American Cyanamid Company  
Agricultural Division  
Crop Protection Chemicals Department  
Wayne, NJ 07470 ©1992

**PURSUIT® HERBICIDE**  
**MINOR CHANGES SPECIFIC TO THE REGION 1 SUPPLEMENTAL LABEL**

- A. Under USE AREA MAP, the Delmarva peninsula of Delaware, Maryland and Virginia has been excluded from the shaded area.
  
- B. The sentence under the map has been changed to read: "Use area includes: South Dakota east of U.S. 81, Nebraska east of U.S. 81, Kansas east of U.S. 81, Maryland and Virginia excluding the Delmarva peninsula, and all other states or portions of states in the preceding map."




EPA Reg. No. 241-310

**FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE, CRANBERRY, AND SMALL WHITE TYPE DRY BEANS, LIMA BEANS, AND SOUTHERN AND ENGLISH PEAS ONLY**

**REFER TO THE MAP ON PAGE 6 FOR GEOGRAPHICAL USE AREA**

**READ LABEL CAREFULLY - USE INSTRUCTIONS VARY FOR EACH CROP**

**THIS LABEL EXPIRES DECEMBER 31, 1993**



**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

### GENERAL INFORMATION

PURSUIT application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT only to the legume vegetables specifically listed in this label. DO NOT apply PURSUIT in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. DO NOT apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

### DISCLAIMER

**THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.**

**REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.**

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply PURSUIT in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for postemergence applications. DO NOT use crop oils, petroleum oils, liquid fertilizers or Dash<sup>1</sup>.

<sup>1</sup>Dash is a trademark of BASF Corporation  
\*Registered Trademark of American Cyanamid Company

with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

## LIMA BEANS

**DO NOT** apply PURSUIT after June 30.

Allow at least 30 days between application and harvest.

**Preplant Incorporated Applications.** Apply PURSUIT at the broadcast rate of 2 ounces per acre to sand and loamy sand soils, or 3 ounces per acre to all other soils within 1 week before planting. Applied preplant incorporated, PURSUIT may be tank-mixed with a registered grass herbicide.

**Preemergence Applications.** Apply PURSUIT at the broadcast rate of 2 ounces per acre to sand and loamy sand soils, or 3 ounces per acre to all other soils immediately after, or up to 1 day after planting. PURSUIT may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

## NAVY, GREAT NORTHERN, RED KIDNEY, CRANBERRY, BLACK TURTLE, AND SMALL WHITE TYPE DRY BEANS

**DO NOT** apply to Domino variety black turtle beans.

**DO NOT** apply PURSUIT to sand or loamy sand soils.

**DO NOT** apply PURSUIT after June 30.

Allow at least 60 days between application and harvest.

**Preplant Incorporated Applications.** Apply PURSUIT within 1 week before planting. Applied preplant incorporated, PURSUIT may be tank-mixed with a registered grass herbicide. Refer to the following table for use rate.

Soil Type	Use Rate (oz/acre)
Sands, loamy sands	<b>DO NOT USE</b>
Sandy loams and all other soils with $\leq$ 2% O.M.	2
All other soils with $>$ 2% O.M.	3

plant incorporated application of a registered grass herbicide.

## ENGLISH PEAS

**DO NOT** apply PURSUIT to sand or loamy sand soils.

Allow at least 30 days between application and harvest.

**Preplant Incorporated Applications.** Apply PURSUIT at the broadcast rate of 3 ounces per acre up to 1 week before planting. **DO NOT** apply PURSUIT to English peas that have been or will be treated with trifluralin.

**Preemergence Applications.** Apply PURSUIT at the broadcast rate of 3 ounces per acre immediately after, or up to 1 day after planting. **DO NOT** apply PURSUIT if English peas have been treated with trifluralin.

**Postemergence Applications.** Apply PURSUIT at 3 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture. **DO NOT APPLY PURSUIT POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.** If English peas have been treated with trifluralin, PURSUIT may be applied **ONLY** at 2 ounces per acre for suppression of black nightshade. The use of trifluralin prior to PURSUIT application may increase the likelihood and severity of crop injury.

## SOUTHERN PEAS

Allow at least 30 days between application and harvest.

**Preplant Incorporated Applications.** Apply PURSUIT at the broadcast rate of 4 ounces per acre within 1 week before planting. Applied preplant incorporated, PURSUIT may be tank-mixed with a registered grass herbicide.

**Preemergence Applications.** Apply PURSUIT at the broadcast rate of 4 ounces per acre immediately after, or up to 1 day after planting. PURSUIT may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

**Early Postemergence Applications.** Apply PURSUIT at the broadcast rate of 4 ounces per acre when peas are at least 3 inches in height. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

**WEEDS CONTROLLED**

PURSUIT applied to low organic matter soils at the broadcast rate of 2 ounces per acre preplant incorporated or preemergence will control light to moderate infestations of:

- Nightshade, black
- Nightshade, Eastern black
- Pigweed, redroot

PURSUIT applied at the broadcast rate of 3 ounces per acre preplant incorporated, preemergence or early post-emergence will control:

- Mustard, wild
- Nightshade, black
- Nightshade, Eastern black
- Nightshade, hairy
- Pigweed, redroot

Postemergence applications of 3 ounces per acre must be made to weeds less than 2 inches tall for best results.

When applied as directed at the broadcast rate of 4 ounces per acre, PURSUIT will control or reduce competition from the weeds listed below:

NOTE: C = Control, R = Reduced Competition

(The number under Maximum Leaf Stage indicates the MAXIMUM number of leaves at which weeds should be sprayed postemergence.)

**BROADLEAF WEEDS**

WEEDS CONTROLLED	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Anoda, spurred	C	2	1 - 2
Artichoke, Jerusalem		8	6 - 10
Buffalobur	C**		
Bristly starbur		2	1 - 2
Carpetweed	C		
Cocklebur, common	C**	8	1 - 8
Galinsoga	C		
Jimsonweed	C*	4	1 - 3
Kochia	C	4	1 - 3
Lambsquarters	C*	2	1 - 2
Mallow, Venice	R		
Morningglory,			
entireleaf	R	2	1 - 2
ivyleaf	R	2	1 - 2
pitted	R	2	1 - 2
smallflower	C	4	1 - 3
tall	R	2	1 - 2
Mustard sp.	C	4	1 - 3
Nightshade,			
black	C	4	1 - 3
Eastern black	C	4	1 - 3
hairy	C	4	1 - 3
Pigweed,			
palmer	C	8	1 - 8
redroot	C	8	1 - 8
smooth	C	8	1 - 8
spiny	C	8	1 - 8
Poinsettia, wild	C		
Puncturevine	C		
Purslane, common	C		
Pusley, Florida	C		
Sida, prickly	C*		

**BROADLEAF WEEDS (Continued)**

WEEDS CONTROLLED	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Ragweed,			
common	R	4	1 - 3
giant	R	4	1 - 3
Sage, barnyard		R	1 - 3
Smartweed,			
ladysthumb	C	4	1 - 3
Pennsylvania	C	4	1 - 3
Spurge,			
prostrate	C	4	1 - 3
spotted	C	4	1 - 3
Sunflower, common	C*	4	1 - 3
Velvetleaf	C*	4	1 - 3
Thistle, Canada		R	1 - 3
Waterhemp, tall	C	4	1 - 3

\*When soil applied, common lambsquarters, jimsonweed, prickly sida, velvetleaf and common sunflower are more consistently controlled by preplant incorporated treatments.

\*\*Use soil applications for light to moderate infestations only. Must be preplant incorporated for best results.

**GRASS WEEDS\***

WEEDS CONTROLLED	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Barnyardgrass	R	3	1 - 3
Crabgrass,			
large	R	3	1 - 3
smooth	R	3	1 - 3
Cupgrass, woolly		3**	1 - 3
Foxtail,			
giant	C	6	1 - 6
green	C	3	1 - 3
giant green	C	3	1 - 3
robust purple	C	3	1 - 3
robust white	C	3	1 - 3
yellow	C	3	1 - 3
Goosegrass	R		
Johnsongrass,			
seedling	C	6	1 - 8
rhizome		R	1 - 8
Panicum,			
fall	R		
Texas	R		

\*When soil applied to grasses, more consistent control can be obtained from preplant incorporated treatments.

\*\*PURSUIT controls emerged woolly cupgrass only.

GRASS WEEDS (Continued)

WEEDS CONTROLLED	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Red rice		3	1 - 3
Shattercane	R	6	1 - 8
Signalgrass, broadleaf	R	4	1 - 8
<b>SEDGES</b>			
Nutsedge,			
purple	R	R	1 - 3
yellow	R	R	1 - 3

DO NOT count cotyledon leaves when determining weed stage of growth.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT as recommended in this label:

1. Anytime
  - Soybeans
  - Peanuts
  - Resistant Field Corn
  - Edible bean and pea types listed on this label (Refer to heading on page 1)
2. Two months after PURSUIT application:
  - Snap beans (DE, MD, PA and VA only)
3. Four months after PURSUIT application:
  - Barley
  - Edible bean and pea types not listed on this label (Refer to heading on page 1)
  - Rye
  - Wheat
4. Nine and one-half months after PURSUIT application:
  - Field corn
  - Field corn grown for seed\*
  - Tobacco
5. Eighteen months after PURSUIT application:
 

Alfalfa	Popcorn
Cotton	Safflower
Lettuce	Sorghum
Oats	Sweet Corn
6. Twenty-six months after PURSUIT application:
  - All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.
7. Forty months after PURSUIT application:
  - Sugar beets

\* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT shall be assumed by the user.

### ROTATIONAL CROP GUIDELINE (Continued)


In the event of a crop loss, the bean or pea crop may be replanted.

**DO NOT** feed treated forage, hay, or straw to livestock.

**CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.**

### USE AREA MAP

PURSUIT can be applied to the edible legume vegetables listed only in the states or parts of states shaded in the following map:



Use area includes: South Dakota east of U.S. 81, Nebraska east of U.S. 81, Kansas east of U.S. 81, Maryland and Virginia excluding the Delmarva peninsula, and all other states or portions of states shaded in the preceding map.

**EPA Reg. No. 241-310**

**FOR PREEMERGENCE USE IN SNAP BEANS GROWN  
IN NEW JERSEY ONLY**

**THIS LABEL EXPIRES DECEMBER 31, 1993**

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN  
THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

**GENERAL INFORMATION**

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to snap beans. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

**DO NOT** apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application.

**USE PURSUIT ONLY** if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

**DISCLAIMER**

**THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.**

**REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.**

\*Registered Trademark of American Cyanamid Company

(see reverse side)

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

**DO NOT** apply PURSUIT in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.


Spray equipment used for PURSUIT applications must be drained and thoroughly cleaned with water before being used to apply other products.

**DO NOT** apply PURSUIT after June 20.

Allow at least 30 days between application and harvest.

**DO NOT** make more than one application of PURSUIT per year.

When PURSUIT is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.



**Preemergence Applications.** For suppression of pigweed, nightshade and common purslane, apply PURSUIT at the broadcast rate of 1.5 ounces per acre immediately after, or up to 1 day after planting. PURSUIT may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

## ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT as recommended in this label:

1. Anytime
  - Soybeans
  - Peanuts
  - Edible beans and peas
2. Four months after PURSUIT application:
  - Barley
  - Rye
  - Wheat
3. Nine and one-half months after PURSUIT application:
  - Field corn
  - Field corn grown for seed\*
  - Tobacco
4. Eighteen months after PURSUIT application:

Alfalfa	Popcorn
Cotton	Safflower
Lettuce	Sorghum
Oats	Sweet Corn
5. Twenty-six months after PURSUIT application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.
6. Forty months after PURSUIT application:
  - Sugar beets

In the event of a crop loss, snap beans may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

\* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT shall be assumed by the user.

**CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.**

PE-17065 Rev. 1 11/12/92 SD



EPA Reg. No. 241-310

**FOR USE IN RED KIDNEY TYPE DRY BEANS, WHITE LUPINS,  
LIMA BEANS, AND ENGLISH PEAS ONLY**

**FOR DISTRIBUTION AND USE ONLY IN ILLINOIS, ICWA, AND WISCONSIN**

**READ LABEL CAREFULLY - USE INSTRUCTIONS VARY FOR EACH GROUP**

**THIS LABEL EXPIRES DECEMBER 31, 1993**

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN  
THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

**GENERAL INFORMATION**

PURSUIT application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT only to the legume vegetables specifically listed in this label. DO NOT apply PURSUIT in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. DO NOT apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application. DO NOT apply PURSUIT postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

**DISCLAIMER**

**THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.**

**REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply PURSUIT in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for post applications.

DO NOT use crop oils, petroleum oils, liquid fertilizers or Dash<sup>1</sup>.

DO NOT make more than one application of PURSUIT per year.

<sup>1</sup>Dash is a trademark of BASF Corporation

<sup>2</sup>Registered Trademark of American Cyanamid Company

(see reverse side)

**ACCEPTED**

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under  
EPA Reg. No. 241-310

DO NOT apply PURSUIT after June 30.

DO NOT apply PURSUIT to white lupins grown on sand or loamy sand soils.

Allow at least 30 days between application and harvest of lima beans.

Allow at least 60 days between application and harvest of red kidney beans or white lupins.

**Preplant Incorporated Applications.** Apply PURSUIT at the broadcast rate of 3 ounces per acre within 1 week before planting. Applied preplant incorporated, PURSUIT may be tank-mixed with a registered grass herbicide.

**Preemergence Applications.** Apply PURSUIT at the broadcast rate of 3 ounces per acre immediately after, or up to 1 day after planting. PURSUIT may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

## ENGLISH PEAS

Allow at least 30 days between application and harvest.

**Preplant Incorporated Applications.** Apply PURSUIT at the broadcast rate of 3 ounces per acre up to 1 week before planting. DO NOT apply PURSUIT to English peas that have been or will be treated with trifluralin.

**Preemergence Applications.** Apply PURSUIT at the broadcast rate of 3 ounces per acre immediately after, or up to 1 day after planting. DO NOT apply PURSUIT if English peas have been treated with trifluralin.

**Postemergence Applications.** Apply PURSUIT at 3 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

**DO NOT APPLY PURSUIT POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.** If English peas have been treated with trifluralin, PURSUIT may be applied ONLY at 2 ounces per acre for suppression of black nightshade. The use of trifluralin prior to PURSUIT application may increase the likelihood and severity of crop injury.

PH-17090 Rev. 1 11/12/92 SD

## ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT as recommended in this label:

1. Anytime  
Soybeans  
Peanuts  
Edible bean and pea types listed on this label (Refer to heading on page 1).
2. Four months after PURSUIT application:  
Barley  
Edible bean and pea types not listed on this label (Refer to heading on page 1).  
Rye  
Wheat
3. Nine and one-half months after PURSUIT application:  
Field corn  
Field corn grown for seed\*  
Tobacco
4. Eighteen months after PURSUIT application:  
Alfalfa      Oats      Sorghum  
Cotton      Popcorn      Sweet Corn  
Lettuce      Safflower
5. Twenty-six months after PURSUIT application:  
All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.
6. Forty months after PURSUIT application:  
Sugar beets

In the event of a crop loss, red kidney beans, white lupins, lima beans, or English peas may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

\* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT shall be assumed by the user.

CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.

EPA Reg. No. 241-310

**ACCEPTED**

**FOR USE IN SUCCULENT PEAS, DRY EDIBLE PEAS,  
LENTILS, AND CHICKPEAS**

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 241-310

PURSUIT can be used on succulent peas, dry edible peas (including chickpeas) and lentils in Washington (Adams, Asotin, Columbia, Garfield, Spokane, Walla Walla and Whitman counties), Idaho (Benewah, Clearwater, Idaho, Kootenai, Latah, Lewis, and Nez Perce counties), Oregon (Union, Umatilla and Wallowa counties) and Montana only.

**THIS LABEL EXPIRES DECEMBER 31, 1993**

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

**GENERAL INFORMATION**

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to peas and lentils. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

**DO NOT** apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application.

**USE PURSUIT ONLY** if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans. Plant peas or lentils at least 1/2 inch deep to reduce risk of crop injury.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

**DISCLAIMER**

**THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.**

**REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.**

\*Registered Trademark of American Cyanamid Company

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

**DO NOT** apply PURSUIT in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT applications must be drained and thoroughly cleaned with water before being used to apply other products.

**DO NOT** make more than one application of PURSUIT per year.

Allow at least 30 days between application and harvest of succulent peas.

Allow at least 60 days between application and harvest of dry edible peas, chickpeas, and lentils.

When PURSUIT is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

(see reverse side)

emergence.

PURSUIT may be tank-mixed with Sencor DF<sup>1</sup> or Lexone DF<sup>2</sup> to assist in the control of lambsquarters or mayweed-chamomile (dogfennel). Refer to the Sencor or Lexone label for proper application rates and restrictions.

## WEEDS CONTROLLED

PURSUIT applied at the broadcast rate of 3 ounces per acre preplant incorporated or preemergence will control:

Buckwheat, wild	Nightshade, black	Pigweed, redroot
Kochia	Nightshade, Eastern black	Shepherdspurse
Mustard, wild	Nightshade, hairy	Thistle, Russian

## ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT as recommended in this label:

- Anytime
  - Chickpeas
  - Lentils
  - Peas
  - Peanuts
  - Soybeans
- Four months after PURSUIT application:
  - Barley
  - Edible beans (dry, lima, snap)
  - Rye
  - Wheat
- Nine and one-half months after PURSUIT application:
  - Field corn
  - Field corn grown for seed\*
  - Tobacco
- Eighteen months after PURSUIT application:
  - Alfalfa
  - Cotton
  - Lettuce
  - Oats
  - Popcorn
  - Safflower
  - Sorghum
  - Sweet Corn
- Twenty-six months after PURSUIT application:
  - All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.
- Forty months after PURSUIT application:
  - Sugar beets

In the event of a crop loss, peas, lentils or chickpeas may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

\*Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into field treated previously with PURSUIT shall be assumed by the user.

CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.

<sup>1</sup>Sencor is a trademark of the Parke Company of Parke-Davis Division, Bayer OTC, Leverkusen  
<sup>2</sup>Lexone is a trademark of E. I. duPont de Nemours and Company, Inc.

PE-17086 Rev 1 11/12/92 SD

 **CYANAMID**  
American Cyanamid Company  
Agricultural Division  
Crop Protection Chemicals Department  
Wayne, NJ 07470 ©1992

21 of 2

EPA Reg. No. 241-310

**FOR USE IN GREAT NORTHERN, RED KIDNEY AND PINTO  
TYPE DRY BEANS**

**FOR DISTRIBUTION AND USE ONLY IN NEW MEXICO**

**THIS LABEL EXPIRES DECEMBER 31, 1993**

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN  
THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

**GENERAL INFORMATION**

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. DO NOT apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application. DO NOT apply PURSUIT postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

**DISCLAIMER**

**THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.**

**REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.**

\*Dash is a trademark of BASF Corporation  
\*Registered Trademark of American Cyanamid Company

(see reverse side)

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply PURSUIT in any manner not specifically described in this label.


Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for post applications.

DO NOT use crop oils, petroleum oils, liquid fertilizers or Dash<sup>1</sup>.

DO NOT make more than one application of PURSUIT per year. DO NOT apply PURSUIT after June 30. Allow at least 60 days between application and harvest.



restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

## APPLICATION RATE AND TIMING

**Preemergence Applications.** Apply PURSUIT at the broadcast rate of 3 ounces per acre immediately after, or up to 1 day after planting. PURSUIT may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide. **DO NOT** apply PURSUIT on sand or loamy sand soils.

**Postemergence Applications.** Apply PURSUIT at 3 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

**DO NOT APPLY PURSUIT POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.**

## WEEDS CONTROLLED

PURSUIT applied at the broadcast rate of 3 ounces per acre preemergence or early postemergence will control:

Mustard, wild	Nightshade, hairy
Nightshade, black	Pigweed, redroot
Nightshade, Eastern black	

Postemergence applications of 3 ounces per acre must be to weeds less than 2 inches tall for best results.

## ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT as recommended in this label:

1. Anytime
  - Soybeans
  - Peanuts
  - Edible bean types listed on this label (Refer to heading on page 1).

Peas  
Rye  
Wheat

3. Nine and one-half months after PURSUIT application:
  - Field corn
  - Field corn grown for seed\*
  - Tobacco
4. Eighteen months after PURSUIT application:
  - Alfalfa
  - Cotton
  - Lettuce
  - Oats
  - Popcorn
  - Safflower
  - Sorghum
  - Sweet Corn
5. Twenty-six months after PURSUIT application:
  - All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.
6. Forty months after PURSUIT application:
  - Sugar beets

In the event of a crop loss Great Northern, red kidney or pinto beans may be replanted.

**DO NOT** feed treated forage, hay, or straw to livestock.

\* Several seed companies have tested a wide range of in-breds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT shall be assumed by the user.

**CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.**

EPA Reg. No. 241-310

**FOR USE IN GREAT NORTHERN AND RED KIDNEY TYPE DRY BEANS**

**FOR DISTRIBUTION AND USE ONLY IN COLORADO, KANSAS,  
NEBRASKA AND WYOMING**

**PURSUIT MAY BE APPLIED TO THE LISTED CROPS WEST OF U.S. 81 ONLY.**

**THIS LABEL EXPIRES DECEMBER 31, 1993**

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN  
THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

**GENERAL INFORMATION**

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

**DO NOT** apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application. **DO NOT** apply PURSUIT postemergence after crop has begun to flower or crop injury may result.

**USE PURSUIT ONLY** if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

**DISCLAIMER**

**THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.**

**REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.**

\*Dash is a trademark of BASF Corporation  
\*Registered Trademark of American Cyanamid Company

(see reverse side)

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

**DO NOT** apply PURSUIT in any manner not specifically described in this label.


Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for post applications.

**DO NOT** use crop oils, petroleum oils, liquid fertilizers or Dash<sup>1</sup>

**DO NOT** make more than one application of PURSUIT per year. **DO NOT** apply PURSUIT after June 30. Allow at least 60 days between application and harvest.



exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

## APPLICATION RATE AND TIMING

**Preemergence Applications.** Apply PURSUIT at the broadcast rate of 3 ounces per acre immediately after, or up to 1 day after planting. PURSUIT may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide. **DO NOT** apply PURSUIT on sand or loamy sand soils.

**Postemergence Applications.** Apply PURSUIT at 3 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

**DO NOT APPLY PURSUIT POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.**

## WEEDS CONTROLLED

PURSUIT applied at the broadcast rate of 3 ounces per acre preemergence or early postemergence will control:

Mustard, wild	Nightshade, hairy
Nightshade, black	Pigweed, redroot
Nightshade, Eastern black	

Postemergence applications of 3 ounces per acre must be to weeds less than 2 inches tall for best results.

## ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT as recommended in this label:

1. Anytime
  - Soybeans
  - Peanuts
  - Edible bean types listed on this label (Refer to heading on page 1).

Rye  
Wheat

3. Nine and one-half months after PURSUIT application:
  - Field corn
  - Field corn grown for seed\*
  - Tobacco
4. Eighteen months after PURSUIT application:

Alfalfa	Popcorn
Cotton	Safflower
Lettuce	Sorghum
Oats	Sweet Corn
5. Twenty-six months after PURSUIT application:
  - All crops not listed in this ROTATIONAL CROP GUIDELINE.
6. Forty months after PURSUIT application:
  - Sugar beets

In the event of a crop loss Great Northern or red kidney beans may be replanted.

**DO NOT** feed treated forage, hay, or straw to livestock.

\* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT shall be assumed by the user.

**CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.**



**CYANAMID**

American Cyanamid Company  
Agricultural Division  
Crop Protection Chemicals Department  
Wayne, NJ 07470 ©1992

**PURSUIT® HERBICIDE  
FOR USE IN SOUTH DAKOTA (REGION 7)**

This supplemental label is a composite of language approved previously in supplemental labels on March 5, 1992 (exhibit 11) with one minor statement added for clarification.

A. This supplemental label is restricted to the use of PURSUIT in navy, great northern, red kidney, black turtle, cranberry and small white type dry beans, lima beans, dry edible peas, lentils and chickpeas in South Dakota.

B. GENERAL INFORMATION, DISCLAIMER AND DIRECTIONS FOR USE  
These paragraphs were previously approved in supplemental labeling accepted by EPA on March 5, 1992, and highlighted in exhibit 1., page 1.

The following more restrictive statement was added under DIRECTIONS FOR USE: "Edible beans must receive a minimum of twelve inches combined rainfall and irrigation between planting and harvest."

C. APPLICATION INSTRUCTIONS  
These paragraphs were previously approved in supplemental labeling accepted by EPA on March 5, 1992, and highlighted in exhibit 11, page 2 for navy, great northern, red kidney, black turtle, cranberry, and small white type dry beans and lima beans; and page 12 for dry edible peas, lentils and chickpeas.

D. WEEDS CONTROLLED  
Weeds listed here were previously approved in supplemental labeling accepted by EPA on March 5, 1992, and highlighted in exhibit 11, page 3.

E. ROTATIONAL CROP GUIDELINE  
Some of the language included in the Rotational Crop Guideline was previously approved in supplemental labeling accepted by EPA on March 5, 1992, and highlighted in exhibit 11, page 5. Other minor changes outlined in exhibit 1 have also been incorporated into this supplemental label.

EPA Reg. No. 241-310

**FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE, CRANBERRY  
AND SMALL WHITE TYPE DRY BEANS, LIMA BEANS,  
DRY EDIBLE PEAS, LENTILS, AND CHICKPEAS**

**FOR DISTRIBUTION AND USE ONLY WITHIN SOUTH DAKOTA**

**PURSUIT may be applied to the listed crops west of U.S. 81 only**

**THIS LABEL EXPIRES DECEMBER 31, 1993**

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN  
THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

### GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. DO NOT apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application.

USE PURSUIT ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

### DISCLAIMER

**THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.**

**REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.**

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

\*Registered Trademark of American Cyanamid Company

(see reverse side)

DO NOT apply PURSUIT in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT applications must be drained and thoroughly cleaned with water before being used to apply other products.


Edible beans must receive a minimum of twelve inches combined rainfall and irrigation between planting and harvest.

DO NOT apply to Domino variety black turtle beans.

DO NOT make more than one application of PURSUIT per year.

Allow at least 30 days between application and harvest of lima beans. Allow at least 60 days between application and harvest of dry beans, dry edible peas, lentils and chickpeas.

When PURSUIT is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.



the PURSUIT leaflet label for incorporation instructions.

**Preemergence Applications.** Apply PURSUIT at the broadcast rate of 3 ounces per acre after planting, but prior to crop emergence. If weeds are present at the time of application, a burn-down herbicide such as Roundup<sup>1</sup> or Gramoxone Extra<sup>2</sup> should be tank-mixed with PURSUIT to control emerged weeds.

## WEEDS CONTROLLED

PURSUIT applied at the broadcast rate of 3 ounces per acre preplant incorporated or preemergence will control:

Mustard, wild	Nightshade, hairy
Nightshade, black	Pigweed, redroot
Nightshade, Eastern black	

## ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT as recommended in this label:

1. Anytime
  - Chickpeas
  - Lentils
  - Dry edible peas
  - Peanuts
  - Edible beans (dry and lima listed on this label)
  - Soybeans
2. Four months after PURSUIT application:
  - Barley
  - Edible beans (dry, snap) and peas not listed on this label
  - Rye
  - Wheat
3. Nine and one-half months after PURSUIT application:
  - Field corn
  - Field corn grown for seed\*
  - Tobacco
4. Eighteen months after PURSUIT application:

Alfalfa	Oats	Sorghum
Cotton	Popcorn	Sweet Corn
Lettuce	Safflower	
5. Twenty-six months after PURSUIT application:
  - All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.

\*Several seed companies have tested a wide range of in-breds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT shall be assumed by the user.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

**CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.**

<sup>1</sup>Roundup is a trademark of Monsanto Agricultural Products Company  
<sup>2</sup>Gramoxone Extra is a trademark of ICI Americas, Inc.



**CYANAMID**

American Cyanamid Company  
Agriculture Division  
Crop Protection Chemicals Department  
Wayne, NJ 07470 ©1990

28942

**PURSUIT® HERBICIDE  
FOR USE IN MICHIGAN (REGION 8)**

This supplemental label is a composite of language approved previously in supplemental labels on March 5, 1992 (exhibit 11) and in the full product label on March 16, 1990 (exhibit 12).

- A. This supplemental label is restricted to the use of PURSUIT in navy, great northern, red kidney, black turtle, cranberry and small white type dry beans in Michigan.
- B. GENERAL INFORMATION, DISCLAIMER AND DIRECTIONS FOR USE  
These paragraphs were previously approved in supplemental labeling accepted by EPA on March 5, 1992, and highlighted in exhibit 11, page 1.
- C. APPLICATION INSTRUCTIONS  
These paragraphs were previously approved in supplemental labeling accepted by EPA on March 5, 1992, and highlighted in exhibit 11, page 2 for preplant incorporated and preemergence applications.  
  
Postemergence applications were previously approved in the full label and highlighted in exhibit 12, page 26.
- D. WEEDS CONTROLLED  
Weeds listed under 2 ounce rates were previously approved in supplemental labeling accepted by EPA on March 5, 1992, and highlighted in exhibit 11, page 3 and in the full label, highlighted in exhibit 12, page 26. Weeds listed under 3 ounce rates were previously approved in supplemental labeling accepted by EPA on March 5, 1992, and highlighted in exhibit 11, page 3.
- E. ROTATIONAL CROP GUIDELINE  
Some of the language included in the Rotational Crop Guideline was previously approved in supplemental labeling accepted by EPA on March 5, 1992, and highlighted in exhibit 11, page 5. Other minor changes outlined in exhibit 1 have also been incorporated into this supplemental label.

EPA Reg. No. 241-310

**ACCEPTED**  
FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE,  
CHERRY, AND SMALL WHITE TYPE DRY BEANS ONLY

FOR DISTRIBUTION AND USE IN MICHIGAN ONLY  
(EXCLUDING UPPER PENINSULA)

THIS LABEL EXPIRES DECEMBER 31, 1993

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 241-310

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN  
THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

### GENERAL INFORMATION

PURSUIT application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT only to the legume vegetables specifically listed in this label. DO NOT apply PURSUIT in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. DO NOT apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

### DISCLAIMER

**THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.**

\*Dash is a trademark of BASF Corporation

\*Registered Trademark of American Cyanamid Company

REFER TO THE PURSUIT LEAFLET LABEL FOR  
ADDITIONAL DISCLAIMER STATEMENTS.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply PURSUIT in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for postemergence applications. DO NOT use crop oils, petroleum oils, liquid fertilizers or Dash<sup>1</sup>.

DO NOT make more than one application of PURSUIT per year.

Refer to the PURSUIT leaflet label for approved combination treatments. When PURSUIT is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

(See reverse side)

**Preplant Incorporated Applications.** Apply PURSUIT within 1 week before planting. Applied preplant incorporated, PURSUIT may be tank-mixed with a registered grass herbicide. Refer to the following table for use rate.

Soil Type	Use Rate (oz/acre)
Sands, loam & sands	DO NOT USE
Sandy loams and all other soils with $\leq$ 2% O.M.	2
All other soils with $>$ 2% O.M.	3

**Preemergence Applications.** Apply PURSUIT at the rates shown in the preceding table immediately after, or up to 1 day after planting. PURSUIT may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

**Postemergence Applications.** Apply PURSUIT at 2 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

**DO NOT APPLY PURSUIT POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.**

### WEEDS CONTROLLED

PURSUIT applied to low organic matter soils at the broadcast rate of 2 ounces per acre preplant incorporated, preemergence, or postemergence will control light to moderate infestations of:

Nightshade, black  
Nightshade, Eastern black  
Pigweed, redroot

PURSUIT applied at the broadcast rate of 3 ounces per acre preplant incorporated or preemergence will control:

Mustard, wild  
Nightshade, black  
Nightshade, Eastern black  
Nightshade, hairy  
Pigweed, redroot

**Resistant Field Corn**  
Edible bean and pea types listed on this label (Refer to heading on page 1)

- Two months after PURSUIT application:  
Snap beans (DE, MD, PA and VA only)
- Four months after PURSUIT application:  
Barley  
Edible bean and pea types not listed on this label (Refer to heading on page 1)  
Rye  
Wheat
- Nine and one-half months after PURSUIT application:  
Field corn  
Field corn grown for seed\*  
Tobacco
- Eighteen months after PURSUIT application:  
Alfalfa                      Oats                      Sorghum  
Cotton                      Popcorn                      Sweet Corn  
Lettuce                      Safflower
- Twenty-six months after PURSUIT application:  
All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.
- Forty months after PURSUIT application:  
Sugar beets

\* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT shall be assumed by the user.

**CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.**

**PURSUIT® HERBICIDE  
FOR USE IN WISCONSIN (REGION 9)**

This supplemental label is a composite of language previously approved in supplemental labels accepted by EPA on March 5, 1992 (exhibit 11).

- A. This supplemental label is restricted to the use of PURSUIT in navy, great northern, red kidney, black turtle, cranberry and small white type dry beans, lima beans, southern and English peas, and white lupins in Wisconsin only, and addresses only a rotational crop interval for snap beans.
  
- B. GENERAL INFORMATION, DISCLAIMER AND DIRECTIONS FOR USE  
These paragraphs were previously approved in supplemental labeling accepted by EPA on March 5, 1992, and highlighted in exhibit 11, page 1.
  
- C. ROTATIONAL CROP GUIDELINE  
The two-month rotational crop interval for snap beans was previously approved in supplemental labeling accepted by EPA on March 5, 1992, and highlighted in exhibit 11, page 5. Other minor changes outlined in exhibit 1 have also been incorporated into this supplemental label.

EPA Reg. No. 241-310

**FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE,  
CRANBERRY, SMALL WHITE TYPE DRY BEANS, LIMA BEANS,  
SOUTHERN AND ENGLISH PEAS, AND WHITE LUPINS**

**FOR DISTRIBUTION AND USE IN WISCONSIN ONLY**

**THIS LABEL EXPIRES DECEMBER 31, 1993**

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN  
THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### **GENERAL INFORMATION**

PURSUIT application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

**DO NOT** apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application.

**DO NOT** apply PURSUIT postemergence after crop has begun to flower or crop injury may result.

**USE PURSUIT ONLY** if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

### **DISCLAIMER**

**THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.**


**REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.**

### **ROTATIONAL CROP GUIDELINE**

The following crops may be planted after applying PURSUIT as recommended:

1. Two months after PURSUIT application:

Snap beans



 **CYANAMID**  
American Cyanamid Company  
Agricultural Division  
Crop Protection Chemicals Department  
Wayne, NJ 07470 ©1992