

EPA Reg. No. 241-310

**FOR USE IN ENGLISH PEAS ONLY**

**FOR DISTRIBUTION AND USE ONLY ON THE DELMARVA PENINSULA OF  
MARYLAND, DELAWARE, AND VIRGINIA**

**READ LABEL CAREFULLY**

**THIS LABEL EXPIRES DECEMBER 31, 1993**

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN  
THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

**GENERAL INFORMATION**

PURSUIT application methods and rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

**DO NOT** apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application.

**DO NOT** apply PURSUIT postemergence after crop has begun to flower or crop injury may result.

**USE PURSUIT ONLY** if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

\*Dash is a trademark of BASF Corporation

\*\*Registered Trademark of American Cyanamid Company

**BEST AVAILABLE COPY**

(see reverse side)

**DISCLAIMER**

**THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.**

**REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

**DO NOT** apply PURSUIT in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for postemergence applications. **DO NOT** use crop oils, petroleum oils, liquid fertilizers or Dash<sup>1</sup>.

**ACCEPTED**

MAR 26 1993

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 241-310

another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

DO NOT apply PURSUIT to sand or loamy sand soils.

Allow at least 30 days between application and harvest.

**Preplant Incorporated Applications.** Apply PURSUIT at the broadcast rate of 3 ounces per acre up to 1 week before planting. DO NOT apply PURSUIT to English peas that have been or will be treated with trifluralin.

**Preemergence Applications.** Apply PURSUIT at the broadcast rate of 3 ounces per acre immediately after, or up to 1 day after planting. DO NOT apply PURSUIT if English peas have been treated with trifluralin.

## WEEDS CONTROLLED

PURSUIT applied at the broadcast rate of 3 ounces per acre preplant incorporated or preemergence will control:

Mustard, wild	Nightshade, hairy
Nightshade, black	Pigweed, redroot
Nightshade, Eastern black	

## ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT as recommended in this label:

- Anytime
  - Soybeans
  - Peanuts
  - Resistant Field Corn
  - English peas and lima beans
- Two months after PURSUIT application:
  - Snap beans

- Nine and one-half months after PURSUIT application:
  - Field corn
  - Field corn grown for seed\*
  - Tobacco

- Eighteen months after PURSUIT application:
  - Alfalfa
  - Cotton
  - Lettuce
  - Oats
  - Popcorn
  - Safflower
  - Sorghum
  - Sweet Corn

- Twenty-six months after PURSUIT application:
  - All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.

- Forty months after PURSUIT application:
  - Sugar beets

\* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT shall be assumed by the user.

In the event of a crop loss, the bean or pea crop may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

**CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.**