

PM 25 241-310

1048

Pursuit[®]	
herbicide	
FOR USE IN SOYBEANS AND PEANUTS	
ACTIVE INGREDIENT	
INERT INGREDIENTS	
TOTAL	
EPA Reg. No. 241-310	EPA Est. No. 241-PR-02
KEEP OUT OF REACH OF CHILDREN	
CAUTION! PRECAUTION!	
PRECAUCION AL USUARIO	In case of an emergency endangering life or property involving this product, call collect day or night, area code 201 835 3100
STATEMENT OF PRACTICAL TREATMENT	
IF ON SKIN	
IF IN EYES	
IF INHALED	
CYANAMID American Cyanamid Company	

ACCEPTED
MAY 11 1993
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 241-310

BEST AVAILABLE COPY

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

CAUTION!

Harmful if inhaled or absorbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, or areas where surface water is present, or to intertidal areas below the mean high water mark. DO NOT contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of pesticide application.

Observe all cautions and limitations on this label and on the labels of products used in combination with PURSUIT. DO NOT use PURSUIT other than in accordance with the instructions set forth on this label. The use of PURSUIT not consistent with this label may result in injury to crops. Keep containers closed to avoid spills and contamination.

DO NOT apply this product through any type of irrigation system.

STORAGE AND DISPOSAL

PROHIBITIONS: KEEP FROM FREEZING. DO NOT store below 32°F. DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, by incineration or, if allowed by State and local authorities by burning. If burned, stay out of smoke.

DISCLAIMER

The label instructions for the use of this product reflect the opinion of experts based on research and field use. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the use of, or application of the product contrary to label instructions, all of which are beyond the control of American Cyanamid Company. All such risks shall be assumed by the user.

American Cyanamid Company shall not be responsible for losses or damages resulting from use of this product in any manner not

set forth on this label. User assumes all risks associated with the use of this product in any manner not specifically set forth on this label.

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use thereon described when used in accordance with the directions for use, subject to the risks referred to above. CYANAMID DOES NOT MAKE OR AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED AND EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

BUYER'S EXCLUSIVE REMEDY AND AMERICAN CYANAMID'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL BE LIMITED TO REFUND OF THE PURCHASE PRICE OF PURSUIT. In no case shall Cyanamid or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product.

Uses With Other Products (Tank-mixes)

If this product is used in combination with any other product except as specifically recommended in writing by American Cyanamid Company, then American Cyanamid Company shall have no liability for any loss, damage or injury arising out of its use in any such combination not so specifically recommended. If used in combination recommended by American Cyanamid Company, the liability of American Cyanamid Company shall in no manner extend to any damage, loss or injury not directly caused by the inclusion of the American Cyanamid Company product in such combination use, and in any event shall be limited to return of the account of the purchase price of the American Cyanamid Company product.

GENERAL INFORMATION

Entry into treated fields is not permitted without protective clothing until sprays have dried.

PURSUIT can be applied as an early preplant, preplant incorporated, preemergence, at ground-cracking (peanuts) or early postemergence. The application method of choice will depend on the anticipated weed spectrum and the preference of the applicator. The mode of weed killing activity involves uptake of PURSUIT by weed roots and/or foliage and rapid translocation to the growing points. Adequate soil moisture is important for optimum PURSUIT activity. When adequate soil moisture is present, PURSUIT will provide residual control of susceptible germinating weeds; activity on established weeds will depend on the weed species and the location of its root system in the soil.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

Occasionally, internode shortening and/or temporary yellowing of crop plants may occur following PURSUIT applications. These effects occur infrequently and are temporary. Normal growth and

appearance should resume within 1 to 2 weeks. These effects will not affect crop yields.

Replanting: If replanting is necessary in a field previously treated with PURSUIT, the field may be replanted to soybeans or peanuts. Rework the soil no deeper than the treated zone. Do not apply a second treatment of PURSUIT.

MIXING INSTRUCTIONS

Fill the spray tank one-half to three-quarters full with clean water. Use a calibrated measuring device to measure the required amount of PURSUIT. Add PURSUIT to the spray tank while agitating. Fill the remainder of the tank with water.

POSTEMERGENCE APPLICATIONS OF PURSUIT REQUIRE THE ADDITION OF AN ADJUVANT AND A LIQUID FERTILIZER SOLUTION.

1. ADJUVANTS

SURFACTANTS:

Use a non-ionic surfactant containing at least 80% active ingredient. Apply the surfactant at the rate of 1 qt./100 gal.

OR

CROP OIL CONCENTRATE:

Instead of a surfactant, a petroleum or vegetable seed based crop oil concentrate (such as SUN-IT II[®]) may be used. SUN-IT II is recommended when weeds are under moisture or temperature stress. Use SUN-IT II or other crop oil concentrates at the rate of 1.5 to 2 pts./acre (use the higher rate when weeds are at the maximum label size).

AND

2. LIQUID FERTILIZER SOLUTION

Recommended nitrogen based fertilizers include liquid fertilizers (such as 28%N, 32%N or 10-34-0) at the rate of 1-2 gals./acre. Use the higher rate when weeds are under moisture or temperature stress. Instead of a liquid fertilizer, spray grade ammonium sulfate may be used at the rate of 2.5 lbs./acre.

NOTE: Liquid fertilizer solutions are not required in PURSUIT applications in use areas south of Interstate Highway 40.

When tank-mixing PURSUIT with recommended herbicides, add the other herbicide to the spray tank first and make sure it is thoroughly mixed before adding PURSUIT. When tank-mixing Bronco[®], Roundup[®], Gramoxone Extra[®], or Starline[®] with PURSUIT herbicide and PROWL[®] herbicide, trillerain, Lasso[®] or Dual[®] add the products in the following order: (1) PROWL, trillerain, Lasso or Dual, (2) Roundup, Gramoxone Extra or Starline, (3) PURSUIT, (4) non-ionic surfactant or crop oil concentrate, and (5) liquid fertilizer.

If dicamba sprayer contamination is possible, refer to the appropriate dicamba label (Banvel, Markman) for sprayer cleanup directions. Clean the sprayer prior to PURSUIT applications to avoid dicamba injury to soybeans.

¹ Bronco, Lasso and Roundup are trademarks of Monsanto Agricultural Products Company.

² Gramoxone Extra and Starline are trademarks of ICI Americas, Inc.

³ Dual is a trademark of Ciba Geigy Corporation.

⁴ Registered trademark of American Cyanamid Company.

⁵ SUN-IT II is a trademark of Agco, Inc.

To avoid injury to sensitive applications must be drained before used to apply other products. When PURSUIT is used in combination with other products, the more restrictive label restrictions should be followed. PURSUIT has a label prohibiting such use.

SPRAYING

DO NOT apply when wind speed may be carried to sensitive crops and sugarbeets.

GROUND APPLICATIONS:

Uniformly apply with proper gallons of water per acre. A spray tank should be calibrated to ensure thorough coverage. To ensure thorough coverage, apply PURSUIT at the rate of 1 qt./100 gal. Adjust the boom height according to the manufacturer's tips for postemergence application. Avoid overlaps when spraying.

PURSUIT APPLICATION

PURSUIT may be applied to applying PURSUIT with a Spray tank. The maximum size listed in dependent upon good spray calibration to deliver the recommended adequate spray coverage. When applying PURSUIT with gallons per acre of spray solution for optimum coverage.

AERIAL APPLICATION:


PURSUIT herbicide may be applied by air to peanuts.

Uniformly apply with proper gallons of water per acre. Use of a non-ionic surfactant and weed control. Apply a non-ionic gallons of spray mixture per acre. (See instructions under EMERGENCE).

To avoid injury to sensitive crops to the following SPECIAL AERIAL APPLICATIONS:

- Nozzle height above ground
- Nozzles must be pointed at a 45 degree angle of the nozzle
- To minimize wing-tip be located any closer than distance from the center

BEST AVAILABLE COPY



- Use a maximum spray pressure of 40 psi.
- A buffer zone must be established between the area to be sprayed and sensitive crops.
- DO NOT spray when wind velocity is greater than 5 mph. Coarse sprays (large droplets) are less likely to drift.

Applicator is responsible for any loss or damage which results from spraying PURSUIT in a manner other than recommended in this label. In addition, applicator must follow all applicable state and local regulations and ordinances in regard to spraying.

APPLICATION INFORMATION

POSTEMERGENCE

PURSUIT is effective in controlling weeds in conservation tillage as well as in conventional production systems. Apply PURSUIT herbicide as an early postemergence treatment when weeds are actively growing and before they exceed a height of 3 inches, unless otherwise indicated. Delay application until the majority of the weeds are at the recommended growth stage. Application timing should be based on weed size and not crop growth stage. Apply PURSUIT to crops and weeds that are actively growing.

An adjuvant (either a surfactant or a crop oil concentrate) and a liquid fertilizer must be added to the spray solution for optimum weed control activity. See the ADDITIVE section under MIXING INSTRUCTIONS for specific instructions.

When PURSUIT is applied postemergence, absorption will occur through both the roots and foliage. Susceptible weeds stop growing and either die or are not competitive with the crop. PURSUIT not only controls many existing broadleaf and grass weeds when applied postemergence, it also provides control of susceptible weeds that may emerge after application.

For maximum weed control, cultivate 7 - 10 days following a postemergence PURSUIT application. This timely cultivation will enhance residual weed control, especially under dry conditions.

If air temperatures reach or stay below 40° for 10 or more hours, delay a PURSUIT application for 48 hours from the time temperatures increase above 40°. Unusually cool temperatures (40° or less) reduces photosynthesis and transpiration. This reduces the uptake and translocation (and effectiveness) of PURSUIT in weeds.

Use a soil applied grass herbicide (such as PROWL®) to control grass weeds not on the PURSUIT label and to enhance the control of certain broadleaf weeds such as common lambsquarters. Refer to the PROWL (or other grass herbicide) label for specific use recommendations, rates and precautions.

PURSUIT should be applied a minimum of one hour before rainfall or overhead irrigation.

NO-TILL/MINIMUM TILLAGE AND DOUBLECROP SOYBEANS:

PURSUIT controls existing weeds and provides residual control of most weeds when applied early postemergence in no-till, minimum tillage and doublecrop soybean production systems. The application may be applied either before or after emergence of the soybeans. (Refer to the WEEDS CONTROLLED POSTEMERGENCE chart for weeds controlled and recommended weed size)

If PURSUIT is applied prior to emergence of the crop, and weeds exceed the recommended size, a contact herbicide should be added to PURSUIT to enhance control. (See instructions for NO-TILL OR REDUCED TILLAGE under the PREEMERGENCE section of this label).

SOIL APPLICATIONS

PURSUIT provides effective weed control in conservation tillage systems designed to meet conservation compliance requirements. PURSUIT can be applied as an early preplant, preplant incorporated, or preemergence treatment in soybeans. It can also be applied in conventional, minimum tillage and no-till production systems. The application method of choice will depend on the anticipated weed spectrum and the preference of the applicator.

- Adequate soil moisture is required for optimum activity. Rainfall or overhead irrigation is necessary to move PURSUIT into the weed germination zone. The amount of rainfall or irrigation required following application depends on existing soil moisture, soil texture and organic matter content. Sufficient water to moisten the soil to a depth of 2 inches is normally adequate. If adequate moisture is not received within 7 days after treatment, a cultivation is recommended to control escaped weeds. When adequate moisture is received after dry conditions, PURSUIT will provide residual control of susceptible germinating weeds; activity on established weeds will depend on the weed species and the location of its root system in the soil.
- PURSUIT controls weeds by uptake by weed roots and translocation to the growing points where it stops weed growth. Susceptible weeds may emerge, growth will stop and the weeds will either die or are not competitive with the crop.

SOIL APPLICATIONS WITH LIQUID FERTILIZERS

PURSUIT can be applied to the soil in liquid fertilizers, alone or in combination with PROWL, Infratrilin, Lasso or Dual. Follow all PURSUIT label recommendations regarding incorporation, timing of application, special instructions and precautions. Apply treatments in 20 or more gallons of liquid fertilizer per acre with ground equipment. Always test the compatibility of PURSUIT with the liquid fertilizer before mixing in the spray tank.

PREEMERGENCE (SURFACE APPLICATIONS)

PURSUIT offers flexibility in that it can be utilized in all production tillage systems. It can be applied prior to planting (up to 45 days prior to planting), at planting in conventional, reduced tillage or no-till production systems, or after planting and before crop emergence.

NO-TILL OR REDUCED TILLAGE:

Apply PURSUIT treatments before, during or after planting but before crop emergence. To ensure thorough coverage, use a minimum of 20 gallons of water per acre. Use higher gallonage for fields with dense vegetation or heavy crop residues.

For maximum grass control, tank-mix PURSUIT with PROWL, Lasso or Dual. To kill existing vegetation, Gramoxone Extra, Starfire or Roundup may be tank-mixed with PURSUIT alone or in combination with PROWL, Lasso or Dual. Gramoxone Extra, Starfire or Roundup should be deleted from the tank-mixture if vegetation is absent at the time of application.

NOTE: Adjust planters to ensure adequate seed coverage

PREPLANT INCORPORATED APPLICATIONS

PURSUIT may be applied following land preparation and should be thoroughly incorporated to a depth of 1 to 2 inches. When applied to beds,

maintain PURSUIT in the surface 1 to 2 inches of the finished beds. Application may be made up to 45 days prior to planting soybeans. Mechanical incorporation can be achieved by one of the following:

- (a) Disk harrow set to cut 3 to 4 inches deep and operated at 4 to 6 mph with the second pass at an angle to the first.
- (b) PTO-driven equipment (tillers, cultivators, hoes) set to cut 2 inches deep and operated one time at 4 mph or less.
- (c) Rolling cultivator set to cut 2 to 4 inches deep and operated two times at 6 to 8 mph with the second pass at an angle to the first. Use only on coarse textured soils.
- (d) Field cultivator with 3 or 4 rows of sweeps (Do not use chisel points), spaced at intervals of 7 inches or less and staggered so that no soil is left unturned, set to cut 3 inches deep, and operated two times at more than 5 mph with the second pass at an angle to the first.
- (e) LELY-ROTERRA® set and operated according to the manufacturer's directions to thoroughly incorporate PURSUIT to a depth of 1 to 3 inches.
- (f) Do-All set and operated according to the manufacturer's directions to incorporate PURSUIT 1 to 3 inches deep and operated at 4 to 6 mph.

If soybeans are planted on beds, apply and incorporate after bed formation using PTO-driven equipment or a rolling cultivator.

When PURSUIT is soil applied for control of nutsedge in peanuts, incorporate with two passes of the incorporation implement. Make the second pass at an offset angle to the first pass to minimize the potential for streaking.


CROPS

SOYBEANS - DIRECTIONS FOR USE

PURSUIT HERBICIDE USE AREA (SOYBEANS) (NOT FOR USE IN CALIFORNIA)

PURSUIT may be applied to soybeans only in the states or parts of states shaded in the map below:

(In Minnesota PURSUIT may be applied south of state highway 210. In Texas PURSUIT may be applied east of state highway 83).



USE RATE (4 OUNCES PER ACRE)

Apply PURSUIT at a broadcast rate of 4 ounces per acre (1/4 pint) for all methods of application: early preplant, preplant incorporated, preemergence, and postemergence (including minimum and no-till). At this broadcast rate, one gallon of PURSUIT will treat 32 acres of soybeans.

NOTE: Only one application of PURSUIT may be made during the season.

WEEDS CONTROLLED

When applied as directed, PURSUIT will control or reduce competition from the weeds listed below. Refer to the ADDITIVES section on page 2 for recommendations when weeds are at the maximum recommended growth stage, or are under stress.

NOTE: C = Control, R = Reduced Competition

(The number under Maximum Leaf Stage indicates the MAXIMUM number of leaves at which weeds should be sprayed postemergence.)

Weeds controlled	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Arnica, spurred	C	2	1-2
Artichoke, Jerusalem		8	6-10
Butterbur	C*		
Bristly starbur		2	1-2
Carpetweed	C		
Cocklebur, common**	C**	8	1-8
Gambusia	C		
Jimsonweed	C*	4	1-3
Kochia	C	4	1-3
Lambsquarters, common	C*	2	1-2
Mallow, Venice	R		
Marshelder	C	4	1-3
Morning glory			
entireleaf	R	2	1-2
ivyleaf	R	2	1-2
pitted	R	2	1-2
smallflower	C	4	1-3
tall	R	2	1-2
Mustard sp.	C	4	1-3

* LELY-ROTERRA is a trademark of Lely

® Registered Trademark of American Cyanamid Company

ACCEPTED
MAY 11 1993
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 241-310

BEST AVAILABLE COPY

BEST AVAILABLE COPY

Weeds Controlled	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Nightshade			
black	C	4	1-3
Eastern black	C	4	1-3
hairy	C	4	1-3
Pigweed			
palmer	C	8	1-8
redroot	C	8	1-8
smooth	C	8	1-8
spiny	C	8	1-8
Poinsettia, wild	C		
Puncturevine	C		
Purslane, common	C		
Pusley, Florida	C		
Sida, prickly	C		
Ragweed,			
common	R	4	1-3
giant	R	4	1-3
Sage, barnyard	R		1-3
Smartweed			
ladythumb	C	4	1-3
Pennsylvania	C	4	1-3
Spurge			
prostrate	C	4	1-3
spotted	C	4	1-3
Sunflower, common	C	4	1-3
Velvetleaf	C	4	1-3
Thistle, Canada	R		1-3
Waterhemp, tall	C	4	1-3

* When soil applied, common lambsquarters, smoothstem, prickly sida, velvetleaf and common sunflower are more consistently controlled by preplant incorporated treatments.
 ** Must be preplant incorporated for best results. Cultivation and/or a postemergence herbicide may be required for season-long results.

GRASS WEEDS*

Weeds Controlled	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Barnyardgrass	R	3	1-3
Crabgrass, large	R	3	1-3
Crabgrass, smooth	R	3	1-3
Crabgrass, woolly		3**	1-3
Foxtail,			
giant	C	6	1-6
green	C	3	1-3
yellow	C	3	1-3
Goosegrass	R		
Johnsongrass,			
seedling	C	6	1-8
rhizome		R	1-8
Millet, wild proso	R	R	1-3
Panicum,			
fall	R		
Texas	R		
Red rice		3	1-3
Shattercane	R	6	1-8
Signalgrass, broadleaf	R	4	1-8
Sorghum, alnum	R	6	1-3

SEDGES

Nutsedge			
purple	R	R	1-3
yellow	R	R	1-3

* When soil applied to grasses, more consistent control can be obtained from preplant incorporated treatments.
 ** PURSUIT controls emerged weedy crabgrass only.
 Do not count cotyledon leaves when determining weed stage of growth. PURSUIT is active against many broadleaf and grass species. However, when heavy grass or common lambsquarters pressure is anticipated, PURSUIT should be used in combination with a registered soil-applied grass herbicide (See HERBICIDE COMBINATIONS section).

HERBICIDE COMBINATIONS

GRASS WEEDS

When heavy grass pressure or common lambsquarters is anticipated, PURSUIT should be used in combination with one of the following registered soil-applied grass herbicides: PROWL herbicide, trifluralin, Lasso or Dual.

When applied as directed, PURSUIT preplant incorporated or preemergence combination treatments with PROWL, trifluralin, Lasso or Dual will control the weeds listed in following table, in addition to those controlled by PURSUIT alone.

GRASSES	PROWL ¹	trifluralin ²	Lasso ³	Dual ⁴
Barnyardgrass	X	X	X	X
Crabgrass, smooth	X	X	X	X
Crabgrass, large	X	X	X	X
Crowfootgrass	X	X		
Goosegrass	X	X	X	X
Millet, wild proso	X	X		
Panicum, fall	X	X	X	X
Panicum, Texas	X	X		
Sandbur, field	X	X		
Shattercane	X	X		
Signalgrass, broadleaf	X	X	X	X
Witchgrass	X	X	X	X

¹ Preplant incorporated tank-mixture applications of PURSUIT plus PROWL will suppress the growth of setgrass, wild proso millet and rhizome johnsongrass.
² Preplant incorporated treatments only.

A selective postemergence grass herbicide such as Select⁵, Poast Plus⁶, Fusilade 2000⁷, Fusivar⁸, Assure⁹ or Option¹⁰ may be mixed with PURSUIT to control volunteer corn or grasses not controlled by PURSUIT. For best results use SUN-IT II or a crop oil concentrate AND liquid fertilizer with grass herbicide tank mixtures. In some cases the activity of the grass herbicide may be reduced when mixed with PURSUIT. The reduction in activity may be overcome by delaying the application of the postemergence grass herbicide 7 days following the application of PURSUIT. If the grass herbicide is applied first, wait 3 days before applying PURSUIT. Refer to the respective grass herbicide label for recommended application rate, weed size and restrictions.

BROADLEAF WEEDS

Broadleaf herbicides that can be tank-mixed with PURSUIT include Basagran¹¹, Blazer¹², Bronco¹³, Cobra¹⁴, Galaxy¹⁵, Gramoxone Extra¹⁶, Roundup¹⁷, Storm¹⁸, or Reflex¹⁹. Certain herbicides should not be applied with PURSUIT (see Page 9 for restrictions).

PURSUIT may be applied either preplant incorporated or preemergence in tank-mix combination with metribuzin (Sencor²⁰ or Laxone²¹) or preemergence with linuron (Lorox²² or Linex²³). PURSUIT may also be applied preplant incorporated, followed by a preemergence application of metribuzin or linuron. The tank-mix or sequential application of PURSUIT and metribuzin will enhance the control of common cocklebur and common ragweed in addition to the weeds controlled by a soil application of

PURSUIT. The addition of linuron to PURSUIT will enhance the control of common ragweed.

PEANUTS - DIRECTIONS FOR USE

PURSUIT USE AREA (PEANUTS) (NOT FOR USE IN CALIFORNIA)

PURSUIT may be used on peanuts in any state where peanuts are grown (except California).

PURSUIT HERBICIDE USE RATE (4 OUNCES PER ACRE)

Apply PURSUIT at a broadcast rate of 4 ounces per acre (1/4 pint) for all methods of application (except sequential²⁴ see below) preplant incorporated, preemergence, ground-checking and postemergence. At this broadcast rate, one gallon of PURSUIT will treat 32 acres of peanuts. PURSUIT may also be applied in a sequential application (See page 8). Apply 2 ounces in a soil application (preplant incorporated or preemergence) followed by 2 ounces applied at ground-check or postemergence. NOTE: Do not apply more than 4 ounces of PURSUIT during the growing season.

WEEDS CONTROLLED

When applied as directed, PURSUIT will control or reduce competition from the weeds listed below. Refer to the ADDITIVES section on page 2 for recommendations when weeds are at the maximum recommended growth stage, or are under stress.

NOTE: C = Control, R = Reduced Competition.
 (The number under Maximum Leaf Stage indicates the MAXIMUM number of leaves at which weeds should be sprayed postemergence.)

BROADLEAF WEEDS

Weeds Controlled	SOIL APPLIED	AT-CRACK	POSTEMERGENCE	
			Maximum Leaf Stage	Size (inches)
Anoda, spurred	C	C	2	1-2
Butterbur	C	C		
Bristly starbur			2	1-2
Carpetweed	C	C		
Cocklebur, common	C	C	8	1-8
Devilclaw	C	C		

¹ Cobra and Select are trademarks of Valent Chemical Co.
² Basagran, Blazer, Storm, and Poast Plus are trademarks of BASF Corporation.
³ Fusilade 2000, Fusivar, Gramoxone Extra and Reflex are trademarks of ICI Americas, Inc.
⁴ Assure, Galaxy, Lasso and Lorox are trademarks of E. I. du Pont de Nemours & Co., Inc.
⁵ Option is a trademark of Hoechst-Roussel Agri-Vet Company.
⁶ Bronco and Roundup are trademarks of Monsanto Agricultural Products Co.
⁷ Sencor is a trademark of Mobay Corporation.
⁸ Linex is a trademark of Giffen Ag Products Co., Inc.

6 of 8

BEST AVAILABLE COPY

ROTATIONAL CROPS

The following rotational crops may be planted after applying PURSUIT at recommended rate:

Anytime

- Lima beans
- Southern peas
- Soybeans
- Peanuts
- Resistant field corn

Four months after PURSUIT application:

- Rye
- Wheat
- Edible beans and peas (other than lima beans and Southern peas)

Nine and one-half months after PURSUIT application:

- Barley
- Field corn
- Field corn grown for seed*
- Tobacco

Eighteen months after PURSUIT application:

- Alfalfa
- Cotton
- Lettuce
- Oats
- Popcorn
- Safflower
- Sorghum
- Sweet corn

Twenty six months after PURSUIT application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDE-LINE.

Forty months after PURSUIT application:

- Suparbeets

* Several seed companies have tested a wide range of hybrids for sensitivity to PURSUIT and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the hybrid data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn hybrids into fields treated previously with PURSUIT shall be assumed by the user.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

PRECAUTIONS

If soybeans are fallow irrigated, till the soil prior to planting winter wheat or barley. The beds should be broken up and the soil mixed with tillage equipment set to cut 4-8 inches deep.

Only rotational crops harvested at maturity may be used for feed or food.

In the event of a crop loss due to weather, soybeans, peanuts or resistant field corn can be replanted. DO NOT work the soil deeper than 2 inches.

There should be an interval of at least 85 days between an application of PURSUIT and soybean or peanut harvest.

PURSUIT applications should be made before soybean bloom.

DO NOT graze or feed treated soybean forage, hay or straw to livestock.

DO NOT graze or feed treated peanut forage, vines, hay or straw to livestock.

Except where stated on a supplemental label or in this label, DO NOT apply to soybeans products containing chlorimuron ethyl (Classic[®], Canopy[®], Zenith[®], Lorax Plus[®], Preview[®], etc.) imazamox (SCEPTER[®], SQUADRON[®], TRI-SCEPTER[®], SCEPTER[®] O.T.[®], SCEPTER[®] 700G) or products containing mazathapyr (PURSUIT, PURSUIT[®] PLUS, PASSPORT[®]) the same year as PURSUIT or injury to follow crops may occur.

NOTE: Classic may be applied postemergence to peanuts following a PURSUIT application. Refer to the Classic label for specific use recommendations.

DO NOT apply PURSUIT[®] PLUS or PASSPORT[®] to peanuts the same year as PURSUIT.

DO NOT tank-mix PURSUIT with clomazone containing herbicides (Command[®] or Commence[®]). PURSUIT may be applied postemergence following a soil application of Command or Commence.

For additional information regarding the use of PURSUIT herbicide, call telephone no. 800-842-0500.

[®] Classic, Canopy, Zenith, Lorax Plus and Preview are trademarks of E. I. duPont de Nemours and Company.

[®] Command and Commence are trademarks of FMC Corporation.

[®] Registered Trademarks of American Cyanamid Company.



AMERICAN CYANAMID COMPANY
Agricultural Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1992

INTEG[™] is protected by the following Patents: U.S.A. 4680080, 4592572, 4675062, Canada 422780, and others



FOR USE IN SOYBEANS AND PEANUTS

Application Timing

Application Rate

Application Method

KEEP OUT OF REACH OF CHILDREN CAUTION!

For additional information regarding the use of PURSUIT herbicide, call telephone no. 800-842-0500.



STATEMENT OF PRACTICAL TREATMENT

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF IN EYES: Flush with plenty of water.

IF INHALED: Remove victim to fresh air.

In case of an emergency endangering life or property involving this product, call collect, day or night, area code 201-845-3100

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

CAUTION

Harmful if inhaled or absorbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, or areas where surface water is present, or to intertidal areas below the mean high water mark. DO NOT contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Observe all cautions and limitations on this label and on the labels of products used in combination with PURSUIT. The use of PURSUIT not consistent with this label may result in injury to crops. Keep containers closed to avoid spills and contamination. DO NOT apply this product through any type of irrigation system. See complete Directions For Use inside label.

STORAGE AND DISPOSAL

PROHIBITIONS: KEEP FROM FREEZING.

DO NOT store below 32°F.

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, by incineration or, if allowed by State and local authorities by burning. If burned, stay out of smoke.

See Disclaimer Inside Leaflet Label.

ACCEPTED
MAY 11 1993
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 241-310

**FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE,
CRANBERRY, AND SMALL WHITE TYPE DRY BEANS ONLY**

**FOR DISTRIBUTION AND USE IN MICHIGAN ONLY MAY 11 1993
(EXCLUDING UPPER PENINSULA)**

THIS LABEL EXPIRES DECEMBER 31, 1993

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 241-310

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

GENERAL INFORMATION

PURSUIT application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT postemergence after crop has begun to flower or crop injury may result.

USE PURSUIT ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.

*Dash is a trademark of BASF Corporation

*Registered Trademark of American Cyanamid Company

**REFER TO THE PURSUIT LEAFLET LABEL FOR
ADDITIONAL DISCLAIMER STATEMENTS.**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply PURSUIT in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for postemergence applications. **DO NOT** use crop oils, petroleum oils, liquid fertilizers or Dash¹.

DO NOT make more than one application of PURSUIT per year.

Refer to the PURSUIT leaflet label for approved combination treatments. When PURSUIT is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

(See reverse side)