

BEST AVAILABLE COPY**FOR USE ON FIELD CORN
(APPLY ONLY ON IMI-CORN HYBRIDS)**

DEC 2 - 1993

**USE DIRECTIONS FOR THE USE OF PURSUIT FOR WEED CONTROL
IN IMI-CORN HYBRIDS**

Under the Federal Insecticide, Fungicide, and Plant Disease Act, Pursuit for Weeds is registered under the

241-310

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT LEAFLET LABEL BEFORE USING. SEE THE PURSUIT LEAFLET LABEL FOR PRECAUTIONS, WORKER PROTECTION STANDARD REQUIREMENTS, APPLICATION INFORMATION, USES WITH OTHER PRODUCTS AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product through any type of irrigation system.

- utilize tank-mixes or sequential applications of different mode-of-action herbicides to control the same target weeds
- combine cultivation with herbicide treatments

See your Cyanamid representative for additional information.

Replanting: If replanting is necessary in a field previously treated with PURSUIT, the field may be replanted to IMI-Corn, soybeans, peanuts, lima beans or southern peas. Rework the soil no deeper than the treated zone. DO NOT apply a second treatment of PURSUIT herbicide.

GENERAL INFORMATION

Apply PURSUIT herbicide only on selected field corn hybrids (IMI-Corn) warranted by the seed company to possess resistance/tolerance to direct application of PURSUIT (for example: Pioneer IR hybrids). DO NOT apply PURSUIT to corn hybrids which lack genetic resistance/tolerance to PURSUIT herbicide. Contact your chemical dealer, seed supplier or American Cyanamid to obtain information regarding IMI-Corn hybrids.

PURSUIT can be applied as an early preplant, preplant incorporated, preemergence or postemergence treatment on IMI-Corn hybrids.

Crops growing under stressful environmental conditions can exhibit various injury symptoms which may be more pronounced if herbicides are used. Corn plants treated with PURSUIT may exhibit yellowing on new growth. Such effects occur infrequently and are temporary. Normal growth and appearance should resume within 1 to 2 weeks.

Within any weed population, there may be the potential for the occurrence of weeds resistant to herbicides when compounds that affect the same site of action are used continuously on the same acre. Therefore, any multiple year weed management program should include, when local agronomic practices allow, as many of the following practices as possible:

- use scouting and weed thresholds to determine the need for herbicide treatment

MIXING INSTRUCTIONS

When tank-mixing PURSUIT with recommended herbicides, add the other herbicides and adjuvants in the following order, while agitating:

1. Fill spray tank 1/4 to 1/2 full with clean water.
2. Add PURSUIT and thoroughly mix.
3. Add other aqueous solution products.
4. Add soluble packet products and thoroughly mix.
5. Add WP (wettable powder), DG (dispersible granule), DF (dry flowable), or LF (liquid flowable) formulations.
6. Add EC (emulsifiable concentrate) products.
7. Add surfactant or crop oil to the spray tank.
8. Add liquid fertilizer.
9. While agitating, fill the remainder of the tank with water.

USE RATE**(4 OUNCES PER ACRE)**

Apply PURSUIT at a broadcast rate of 4 ounces per acre (1/4 pint) for all methods of application: early preplant, preplant incorporated, preemergence (including minimum and no-till), and postemergence. At this broadcast rate, one gallon of PURSUIT herbicide will treat 32 acres of IMI-Corn.

NOTE: Only one application of PURSUIT may be made during the growing season.

POSTEMERGENCE

Apply PURSUIT herbicide as a postemergence treatment to IMI-Corn when crop and weeds are actively growing. Apply PURSUIT before weeds exceed a height of 3 inches, unless otherwise indicated (see PURSUIT leaflet label). More restrictive crop growth stage limitations of tank-mix partners must be followed.

ADDITIVES:

POSTEMERGENCE APPLICATIONS OF PURSUIT REQUIRE THE ADDITION OF AN ADJUVANT AND A LIQUID FERTILIZER¹ SOLUTION.

I. ADJUVANTS

SURFACTANTS: Use a non-ionic surfactant containing at least 80% active ingredient. Apply the surfactant at the rate of 1 quart per 100 gallons of spray solution

OR

CROP OIL CONCENTRATE: Instead of a surfactant, a petroleum or vegetable seed oil based crop oil concentrate (such as SUN-IT II™) may be used. SUN-IT II is recommended when weeds are under moisture or temperature stress. Use SUN-IT II or other methylated seed oils at the rate of 1.5-2 pints per acre (use the higher rate when weeds are at the maximum label size), or use a crop oil concentrate at 2 pints per acre.

AND

II. LIQUID FERTILIZER SOLUTION

Recommended nitrogen based fertilizers include liquid fertilizers (such as 28%N, 32%N or 10-34-0) at the rate of 1-2 quarts per acre. Use the higher rate when weeds are under moisture or temperature stress. Instead of a liquid fertilizer, spray grade ammonium sulfate may be used at the rate of 2.5 lbs per acre.

DO NOT apply PURSUIT herbicide with liquid fertilizer as a carrier for postemergence applications.

¹PURSUIT applications may be made with a non-ionic surfactant only (liquid fertilizer not required) in the states of Arkansas, Bootheel of Missouri, Tennessee, South Carolina, Louisiana, Mississippi, Alabama, and Georgia.

²In some cases the grass activity of PURSUIT will be reduced when used in combination with atrazine, Buctril, Banvel, Basagran, Clarity, or Marksman.

³Some corn leaf burn may result with Buctril or atrazine postemergence combinations with PURSUIT.

⁴Applications of Banvel or Clarity herbicide to corn during periods of rapid growth may result in temporary leaning

MT™ or Dual™. Follow all PURSUIT label recommendations regarding incorporation, timing of application, special instructions and precautions. Apply treatments in 20 or more gallons of liquid fertilizer per acre with ground equipment. Always test the compatibility of PURSUIT herbicide with the liquid fertilizer before mixing in the spray tank.

SEQUENTIAL HERBICIDE COMBINATIONS WITH PURSUIT

PURSUIT FOLLOWING PREPLANT APPLICATION OF:

2,4-D	Frontier™
Bicep™, Bicep Lite™	Gramoxone Extra™
Bullet™	Lasso EC
Bronco™	Lasso MT
Dual	Roundup™
Extrazine II™	

PURSUIT FOLLOWING PREPLANT INCORPORATED APPLICATION OF:

Atrazine ²	Extrazine II
Bicep, Bicep Lite	Frontier
Bullet	Lariat™
Dual	Lasso EC, Lasso MT
Eradicane Extra™	Sutan+™

PURSUIT FOLLOWING PREEMERGENCE APPLICATION OF:

Atrazine	Clarity™, ²
Banvei™, ²	Dual
Bicep, Bicep Lite	Extrazine
Bladex™ 4L	Frontier
Bladex 90DF	Lariat
Bronco	Lasso EC, Lasso MT
Bullet	Marksman™, ²
	PROWL™, ³ 3.3EC

TANK-MIX HERBICIDE COMBINATIONS WITH PURSUIT

PREPLANT APPLICATION OF PURSUIT TANK MIXED WITH:

2,4-D	Gramoxone Extra
Bronco	Lasso EC
Dual	Lasso MT
Frontier	Roundup

BEST AVAILABLE COPY

3045

PREPLANT INCORPORATED APPLICATION OF PURSUIT TANK-MIXED WITH:

Atrazine	Extrazine
Bicep, Bicep Lite	Frontier
Bullet	Lariat
Dual	Lasso EC, Lasso MT
Eradicane Extra	Sutan+

PREEMERGENCE APPLICATION OF PURSUIT TANK-MIXED WITH:

Atrazine	Clarity
Banvel	Dual
Bicep, Bicep Lite	Extrazine
Bladex 4L	Frontier
Bladex 90DF	Lariat
Bronco	Lasso EC, Lasso MT
Bullet	Marksman
	PROWL 3.3EC

POSTEMERGENCE APPLICATIONS OF PURSUIT TANK-MIXED WITH:

Atrazine ³	Frontier
Banvel ⁴	Lasso EC, Lasso MT
Basagran ^{™,2}	Marksman
Buctril ^{™,2,3}	PROWL 3.3EC
Clarity ⁴	Sencor DF [™]
Dual	Sencor 4L [™]

DO NOT use crop oil concentrates as adjuvants in PURSUIT combinations with Banvel, Buctril, Clarity, Marksman, Sencor DF or Sencor 4L.

DO NOT tank-mix PURSUIT herbicide with Accent[™] or Beacon[™].

PURSUIT is active against many broadleaf and grass species. However, for long term weed management, alternate mode of action herbicides are recommended with PURSUIT. The application of a soil applied grass herbicide underlay will control grass weeds not on the PURSUIT herbicide label and enhance the control of certain broadleaf weeds such as common lambsquarters.

When PURSUIT is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions.

AERIAL APPLICATIONS

Uniformly apply with properly calibrated aerial equipment in 5 or more gallons of water per acre. When applied POSTEMERGENCE, the addition of a non-ionic surfactant AND fertilizer solution are required for optimum weed control. Apply a non-ionic surfactant at the rate of 1 quart per 100 gallons of spray mixture OR a crop oil concentrate at the rate of 1.5 - 2 pints per acre AND a liquid fertilizer at the rate of 1 quart per acre. (See

instructions under APPLICATION INSTRUCTIONS - POSTEMERGENCE).

To avoid injury to sensitive crops from drift, aerial applicators must adhere to the following SPECIAL AERIAL USE DIRECTIONS AND PRECAUTIONS.

- Nozzle height above ground must be a maximum of 10 feet.
- Nozzles must be pointed toward the rear of the aircraft. The downward angle of the nozzle should not be greater than 20 degrees.
- To minimize wing-tip vortex roll, nozzles or spray boom must not be located any closer to end of wing or rotor than three-fourths the distance from the center of the aircraft.
- Use a maximum spray pressure of 40 psi.
- A buffer zone must be established between the area to be sprayed and sensitive crops.
- DO NOT spray when wind velocity is greater than 5 mph. Coarse sprays (larger droplets) are less likely to drift.

Applicator is responsible for any loss or damage which results from spraying PURSUIT in a manner other than recommended in this label. In addition, applicator must follow all applicable state and local regulations and ordinances in regard to spraying.

When applied as directed at the broadcast rate of 4 ounces per acre, PURSUIT herbicide will control or reduce competition from the weeds listed below.

NOTE: C = Control, R = Reduced Competition

BROADLEAF WEEDS

Weeds Controlled	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (Inches)
Alligator weed		4	1 - 3
Anoda, spurred	C	2	1 - 2
Artichoke, Jerusalem		8	6 - 10
Buffalobur	C*	R	1 - 3
Bristly starbur		2	1 - 2
Carpetweed	C		
Cocklebur, common	R	8	1-8
Galinsoga	C		
Jimsonweed	C*	4	1 - 3
Kochia	C	4	1 - 3
Lambsquarters, common	C*	R	1 - 2
Mallow, Venice	R		
Marshelder	C	4	1 - 3
Momingglory,			
entireleaf	R	2	1 - 2
ivyleaf	R	2	1 - 2
pitted	R	2	1 - 2
smallflower	C	4	1 - 3
tall	R	2	1 - 2
Mustard sp.	C	4	1 - 3
Nightshade,			
black	C	4	1 - 3
Eastern black	C	4	1 - 3
hairy	C	4	1 - 3

BROADLEAF WEEDS (Continued)

Weeds Controlled	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (Inches)
Pigweed,			
palmer	C	8	1 - 8
redroot	C	8	1 - 8
smooth	C	8	1 - 8
spiny	C	8	1 - 8
Poinsettia, wild	C		
Puncturevine	C		
Purslane, common	C		
Pusley, Florida	C		
Sida, prickly	C*		
Ragweed,			
common	R	4	1 - 3
giant	R	4	1 - 3
Sage, barnyard		R	1 - 3
Smartweed,			
ladysthumb	C	4	1 - 3
Pennsylvania	C	4	1 - 3
Spurge,			
prostrate	C	4	1 - 3
spotted	C	4	1 - 3
Sunflower	C*	4	1 - 3
Velvetleaf	C*	4	1 - 3
Thistle, Canada		R	1 - 3
Waterhemp, tall	C	4	1 - 3

*When PURSUIT is soil applied, these weeds are more consistently controlled by preplant incorporated treatments.

DO NOT count cotyledon leaves when determining weed stage of growth.

GRASS WEEDS

Weeds Controlled	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (Inches)
Barnyardgrass	R	3	1 - 3
Crabgrass,			
large	R	3	1 - 3
smooth	R	3	1 - 3
Cupgrass, woolly		3	1 - 3
Foxtail,			
giant	C	6	1 - 6
green	C	3	1 - 3
giant green	C	3	1 - 3
Goosegrass	R		
Johnsongrass,			
seedling	C	6	1 - 8
rhizome	R		1 - 8
Millet, wild proso	R	R	1 - 3
Panicum,			
fall	R		
Texas	R		

GRASS WEEDS (Continued)

Weeds Controlled	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (Inches)
Red rice		3	1 - 3
Sandbur, field	R	R	≤1
Shattercane	R	6	1 - 8
Signalgrass, broadleaf	R	4	1 - 8
Sorghum alnum	R	6	1 - 3

SEDGES

Nutsedge,			
purple	R	R	1 - 3
yellow	R	R	1 - 3

When PURSUIT is applied to grasses, preplant incorporated treatments of PURSUIT are more consistent.

(The number under Maximum Leaf Stage indicates the MAXIMUM number of leaves at which weeds should be sprayed postemergence.)

ROTATIONAL CROPS

Refer to the PURSUIT leaflet label for further rotational crop restrictions.

The following rotational crops may be planted after applying PURSUIT herbicide at the recommended rate in corn:

- Anytime

Lima beans	Peanuts
Southern peas	IMI-Corn
Soybeans	
- Four months after PURSUIT application:
 - Rye
 - Wheat
 - Edible beans and peas (other than lima beans and southern peas)
- Eight and one-half months after PURSUIT application:
 - Field corn
 - Field corn grown for seed*
- Nine and one-half months after PURSUIT application:
 - Barley
 - Tobacco
- Eighteen months after PURSUIT application:
 - Alfalfa
 - Cotton
 - Lettuce
 - Oats
 - Popcorn
 - Safflower
 - Sorghum
 - Sweet corn
- Twenty six months after PURSUIT application:
 - All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.

BEST AVAILABLE COPY

5075

7. Forty months after PURSUIT application:

Sugarbeets

DO NOT plant other rotational crops prior to 26 months after a PURSUIT herbicide application.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT shall be assumed by the user.

PRECAUTIONS

In the event of a crop loss due to weather, IMI-Corn hybrids, soybeans, lima beans, southern peas or peanuts can be replanted if PURSUIT herbicide was applied alone at rates labelled for these crops. If PURSUIT was tank-mixed with other herbicides, such as atrazine, the label restrictions for these herbicides must also be followed.

There should be an interval of at least 45 days between an application of PURSUIT and corn harvest (silage, fodder, or grain). DO NOT graze or feed treated corn, forage, silage, fodder, or grain for at least 45 days after an application of PURSUIT.

All soil insecticides, including labelled banded or in-furrow applications, may be used in combination with Pioneer imidazolinone resistant (IK) corn hybrids.

Imidazolinone tolerant hybrids from other seed companies may occasionally exhibit injury symptoms when soil insecticides are used in combination with PURSUIT. USE banded applications of COUNTER® 15G, COUNTER® 20CR® or THIMET® in combination with PURSUIT herbicide application; DO NOT USE COUNTER 15G soil insecticide in-furrow with imidazolinone tolerant hybrids. American Cyanamid has not tested all hybrids in which the imidazolinone tolerance trait is claimed and cannot be responsible for factors which are beyond its control, such as growing conditions, environmental conditions, grower practices and the specific genetics of each hybrid which may influence tolerance. Growers should contact the seed company for information regarding specific hybrid tolerance to herbicide and insecticide applications.

This label must be in the possession of the user at the time of herbicide application.

For additional information regarding the use of PURSUIT herbicide, call telephone no. (800) 942-0500.

PE-17150 Rev. 3 8/1/93 SD

- ®Registered Trademark of American Cyanamid Company.
- ™SUN-IT II is a Trademark of Agsco, Inc.
- ™Buctril is a Trademark of Rhone Poulenc Ag Company.
- ™Banvel, Clarity, Frontier, and Marksman are Trademarks of Sandoz Crop Protection Corporation.
- ™Basagran is a Trademark of BASF Corporation.
- ™Beacon, Bicep, Bicep Lite, and Dual are Trademarks of Ciba Geigy Corporation.
- ™Bronco, Bullet, Lariat, Lasso EC, Lasso MT, and Roundup are Trademarks of Monsanto Agricultural Products Company.
- ™Sencor is a Trademark of Mobay Corporation.
- ™Gramoxone Extra, Eradicane Extra, and Sutan+ are Trademarks of ICI Americas, Inc.
- ™Accent, Bladex 4L, Bladex 90DF, and Extrazine are Trademarks of E. I. DuPont de Nemours and Company, Inc.

