

FEB 2 1994

Mrs. Desiree L. Little
American Cyanamid Co.
P.O. Box 400
Princeton, NJ 08540

Dear Mrs. Little:

Subject: Supplemental Label for Use on Pinto Beans
For Distribution and Use Only in Colorado,
Kansas, Wyoming, Nebraska, and Texas
Pursuit® Herbicide
EPA Reg. No. 241-310
Your submission dated January 12, 1992

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

Sincerely yours,

BEST AVAILABLE COPY

Robert J. Taylor
Product Manager 25
Fungicide-Herbicide Branch
Registration Division (H7505C)

RD:RTaylor:eja:88:305-6800

EPA Reg. No. 241-310

FOR POSTEMERGENCE USE IN PINTO TYPE DRY BEANS

**FOR DISTRIBUTION AND USE ONLY IN COLORADO, KANSAS,
WYOMING, NEBRASKA, AND TEXAS**

THIS LABEL EXPIRES DECEMBER 31, 1994

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application.

DO NOT apply PURSUIT postemergence after crop has begun to flower or crop injury may result.

Some varieties of pinto beans may be inherently less tolerant to PURSUIT applications and crop injury could result. Contact the seed company for information and recommendations about postemergence applications of PURSUIT.

USE PURSUIT ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

ACCEPTED
FOR DISTRIBUTION
in Letter Dated:

Registered to: American Cyanamid Company

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, potatoes, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for post applications.

DO NOT make more than one application of PURSUIT per year. **DO NOT** apply PURSUIT after July 15. Allow at least 60 days between application and harvest.

Under the Federal Insecticide, Fungicide, and Plant Antifeedant Act as amended, for the pesticide registered under EPA Reg. No.

241-310

BEST AVAILABLE COPY

exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

APPLICATION RATE AND TIMING

Postemergence Applications. Apply PURSUIT at 3 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

DO NOT use crop oils, petroleum oils, liquid fertilizers or Dash¹.

DO NOT APPLY PURSUIT POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRIFOLIATE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

WEEDS CONTROLLED

PURSUIT applied at the broadcast rate of 3 ounces per acre early postemergence will control:

Mustard, wild	Nightshade, hairy
Nightshade, black	Pigweed, redroot
Nightshade, Eastern black	

Postemergence applications of 3 ounces per acre must be to weeds less than 2 inches tall for best results.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT as recommended in this label:

- Anytime
 - Soybeans
 - Peanuts
 - Great Northern, red kidney, and pinto type dry beans
 - IMI-Corn

Wheat

- Nine and one-half months after PURSUIT application:
 - Field corn
 - Field corn grown for seed*
 - Tobacco
- Eighteen months after PURSUIT application:
 - Alfalfa
 - Cotton
 - Lettuce
 - Oats
 - Popcorn
 - Safflower
 - Sorghum
 - Sweet Corn
- Twenty-six months after PURSUIT application:
 - All crops not listed in this ROTATIONAL CROP GUIDELINE.
- Forty months after PURSUIT application:
 - Sugar beets

In the event of a crop loss Great Northern or red kidney beans may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT shall be assumed by the user.

CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.

BEST AVAILABLE COPY

¹Dash is a trademark of BASF Corporation.

DE XXXXX 11/18/93 50



CYANAMID

American Cyanamid Company
A Division of Eastman-Kodak Company
1000 Park Avenue, New York, N.Y. 10022
© 1993 Eastman-Kodak Company