

PM 25 241-310

10/6

PURSUIT® HERBICIDE SUPPLEMENTAL LABELING

ACCEPTED  
APR 12 1994  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, this pesticide registered under EPA Reg. No. 241-310

EPA Reg. No. 241-310

FOR USE IN SOYBEANS  
IN NORTH DAKOTA AND NORTHERN MINNESOTA  
(SEE USE AREA BELOW)

THIS LABEL EXPIRES DECEMBER 31, 1994

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT LEAFLET LABEL BEFORE USING. SEE THE PURSUIT LEAFLET LABEL FOR ADDITIONAL USE AREAS, MIXING AND APPLICATION INFORMATION, WEEDS CONTROLLED AND ROTATIONAL CROP RESTRICTIONS. THE ROTATIONAL CROP RESTRICTIONS IN THIS SUPPLEMENTAL LABEL SUPERSEDE THE RESTRICTIONS LISTED IN THE LEAFLET LABEL.

GENERAL INFORMATION

PURSUIT herbicide is a broad spectrum herbicide that provides season long control of many weeds. Under some conditions (such as heavy texture soil, high organic matter, low pH or low rainfall) PURSUIT may cause injury to subsequent planted crops. Sugar beets are particularly sensitive to PURSUIT residues in soil. PURSUIT is not recommended in crop rotations that include sugar beets. See the ROTATIONAL CROP section of this supplemental label for rotations to other sensitive crops.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system.

USE AREA

PURSUIT may be used in North Dakota and in Minnesota north of Highway #210.

®Registered Trademark of American Cyanamid Company

2076

**APPLICATION RATES**

Apply PURSUIT at 3 ounces per acre postemergence only.

**BROADLEAF WEEDS**

**WEEDS CONTROLLED**

<u>Weeds controlled</u>	<u>Maximum Leaf Stage</u>	<u>(inches)</u>
Cocklebur, common*	4	1-4
Nightshade, black	4	1-3
Eastern black hairy	4	1-3
Kochia	4	1-3
Pigweed, redroot	4	1-4
Mustard species	4	1-3
Wild oats**	3	1-4

\*For control of common cocklebur, add COBRA<sup>1</sup> herbicide at the rate of 4 ounces per acre to the spray solution.

\*\*PURSUIT will reduce competition from wild oats.

**GRASS WEEDS**

For control of several annual grasses, tank-mix PURSUIT and SELECT<sup>1</sup> herbicides at the following rates:

PURSUIT - 3 oz/Acre  
SELECT - 6-8 oz/Acre\*

\*Refer to the SELECT label for grasses controlled.

**VOLUNTEER CORN**

For control of volunteer corn, tank-mix PURSUIT and SELECT herbicides at the following rates:

PURSUIT - 3 oz/Acre  
SELECT - 4 oz/Acre

Apply the tank-mixture to volunteer corn when 4 - 12 inches tall.

Refer to the COBRA or SELECT labels for additional information, precautions and restrictions. The most restrictive labeling applies when using a tank-mixture.

<sup>1</sup>Trademarks of Valent U.S.A. Corporation

POSTEMERGENCE APPLICATIONS OF PURSUIT REQUIRE THE ADDITION OF AN ADJUVANT AND A LIQUID FERTILIZER SOLUTION.

#### I. ADJUVANTS

**SURFACTANTS:** Use a non-ionic surfactant containing at least 80% active ingredient. Apply the surfactant at the rate of 1 qt./100 gal.

OR

**CROP OIL CONCENTRATE:** Instead of a surfactant, a petroleum or vegetable seed based oil concentrate (such as SUN-IT II™) may be used. SUN-IT II is recommended when weeds are under moisture or temperature stress. Use SUN-IT II or other crop oil concentrates at the rate of 1.5 to 2 pts./acre (use the higher rate when weeds are at the maximum label size).

AND

#### II. LIQUID FERTILIZER SOLUTION

Recommended nitrogen based fertilizers include liquid fertilizers (such as 28%N, 32%N or 10-34-0) at the rate of 1-2 qts./acre. Use the higher rate when weeds are under moisture or temperature stress. Instead of a liquid fertilizer, spray grade ammonium sulfate may be used at the rate of 2.5 lbs./acre.

107W

ROTATIONAL CROPS

The following crops may be planted after applying PURSUIT at the recommended rate in soybeans.

- 1. Anytime
  - IMI-Corn
  - Soybeans
- 2. Nine and one-half months after PURSUIT application:
  - Field Corn
  - Edible beans and peas
  - Wheat
- 3. Eighteen months after PURSUIT application:
  - Barley
  - Oats
  - Rye
- 4. Twenty six months after PURSUIT application:
  - Canola (rapeseed)
  - Carrots
  - Celery
  - Cole crops
  - Garlic
  - Flax
  - Onions
  - Potatoes
  - Spinach
  - Sunflower
  - Tomatoes
- 5. Forty months after PURSUIT application.
  - Sugar beets

DO NOT plant other rotational crops prior to 26 months after a PURSUIT application.

This label must be in the possession of the user at the time of herbicide application.

PE-17132 Rev.2 12/93

American Cyanamid Company  
 Agricultural Division  
 Crop Protection Chemicals Department  
 Wayne, NJ 07470 ©1994


**PURSUIT® HERBICIDE SUPPLEMENTAL LABELING**

EPA Reg. No. 241-310

FOR DISTRIBUTION AND USE ONLY IN THE STATES OF  
IOWA, ILLINOIS, INDIANA AND OHIO

**PLANTING OF POPCORN  
FOLLOWING APPLICATIONS OF PURSUIT HERBICIDE**

This Label Expires December 31, 1994



OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT LEAFLET LABEL BEFORE USING. SEE THE PURSUIT LEAFLET LABEL FOR MIXING AND APPLICATION INFORMATION, AND ROTATIONAL CROP RESTRICTIONS.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system.

**GENERAL INFORMATION/CAUTION**

Some popcorn varieties may be injured when planted at less than 18 months following an application of PURSUIT herbicide. Several popcorn companies have determined their popcorn varieties can be planted with no injury or minimal injury the year following PURSUIT use. Before planting popcorn contact the popcorn company for information and recommendations regarding the tolerance of popcorn varieties planned for fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, **ALL RISKS AND CONSEQUENCES ASSOCIATED WITH PLANTING POPCORN VARIETIES INTO FIELD TREATED PREVIOUSLY WITH PURSUIT SHALL BE ASSUMED BY THE USER.**

Stunting and maturity delay, or other adverse effects may result when popcorn is planted following PURSUIT use.

This label must be in the possession of the user at the time of herbicide application.

**CROP DAMAGE DISCLAIMER**

American Cyanamid does not advocate nor recommend planting popcorn at less than 18 months following an application of PURSUIT. PURSUIT herbicide may cause stunting and delayed maturity to popcorn.

PE-17179 Rev.1 1/94

American Cyanamid Company  
Agricultural Division  
Crop Protection Chemicals Department  
Wayne, NJ 07470 ©1994