

Please read Instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 11-30-93

(A) 	United States Environmental Protection Agency Office of Pesticide Programs (H7505C) Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number  207888
	Application for Pesticide:		

Section I

1. Company/Product Number 241-310	2. EPA Product Manager R. Taylor	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) PURSUIT® herbicide	PM# 25	
5. Name and Address of Applicant (Include ZIP Code) American Cyanamid Company Agricultural Research Center P.O. Box 400 Princeton, NJ 08543-0400 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to:  EPA Reg. No. _____ Product Name _____	

Section II

<input type="checkbox"/> Amendment - Explain below	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - explain below.

**Explanation:** Use additional page(s) if necessary. (For section I and Section II.)

Notification of revision of self-imposed expiration date per PR Notice 95-2. This notification is consistent with the provisions of PR Notice 95-2 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 95-2 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Metal Plastic Glass Paper Other (Specify) _____		
* Certification must be submitted.		If "Yes," Unit Package wgt.      No. per container			
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) of Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner In Which Label Is Affixed To Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Other (_____) <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled					

Section IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Janet M. Overholt	Title Product Registrations Manager	Telephone No. (include Area Code) 609-716-2410
<b>Certification</b> I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Product Registrations Manager	
4. Typed Name Janet M. Overholt	5. Date 2/12/94	



# Supplemental Labeling

2/3

EPA Reg. No. 241-310

## FOR USE IN PEANUTS ONLY

FOR USE ONLY IN NORTH CAROLINA, SOUTH CAROLINA AND VIRGINIA

COTTON PLANTING FOLLOWING PURSUIT APPLICATIONS TO PEANUTS -  
ROTATIONAL CROP GUIDELINE MODIFICATION

THIS LABEL EXPIRES MARCH 30, 1997


OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT LEAFLET LABEL BEFORE USING. SEE THE PURSUIT LEAFLET LABEL FOR USE AREA, MIXING AND APPLICATION INFORMATION, WEEDS CONTROLLED AND ROTATIONAL CROP RESTRICTIONS.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system.

### GENERAL INFORMATION

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.



### ROTATIONAL CROPPING GUIDELINE


Cotton may be planted at the following intervals after a PURSUIT application if the criteria listed are met:

1. Nine and one-half months after an application of PURSUIT
  - PURSUIT applied to peanuts only, and
  - soil texture is sandy loam or loamy sand only, and
  - greater than 16 inches of rainfall and/or irrigation is received following application of PURSUIT through October of the application year.
  - Postemergence applications only. (Refer to the PURSUIT herbicide label for application instructions).
  
2. Eighteen months after an application of PURSUIT
  - If the above criteria are not met, rotation to cotton must be at least 18 months following an application of PURSUIT herbicide.

This label must be in the possession of the user at the time of herbicide application.

FE-17137 Rev. 2 1/96 SD

American Cyanamid Company  
Agricultural Products Division  
Crop Protection Chemicals Division  
Wayne, NJ 07470 ©1996



Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT application must be drained and thoroughly cleaned with water before being used to apply other products.

**DO NOT** make more than one application of PURSUIT per year.

Refer to the PURSUIT leaflet label for approved combination treatments. When PURSUIT is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

**DO NOT** apply PURSUIT after June 30.

Allow at least 30 days between application and harvest.

**Preplant Incorporated Applications.** Apply PURSUIT at the broadcast rate of 2 ounces per acre to sand and loamy sand soils, or 3 ounces per acre to all other soils within 1 week before planting. Applied preplant incorporated, PURSUIT may be tank-mixed with a registered grass herbicide.

**Preemergence Applications.** Apply PURSUIT at the broadcast rate of 2 ounces per acre to sand and loamy sand soils, or 3 ounces per acre to all other soils immediately after, or up to 1 day after planting. PURSUIT may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

**PURSUIT applied to low organic matter soils at the broadcast rate of 2 ounces per acre preplant incorporated or preemergence will control light to moderate infestations of:**

Nightshade, black  
Nightshade, Eastern black  
Pigweed, redroot

**PURSUIT applied at the broadcast rate of 3 ounces per acre preplant incorporated or preemergence will control:**

Mustard, wild  
Nightshade, black  
Nightshade, Eastern black  
Nightshade, hairy  
Pigweed, redroot

### ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT as recommended in this label:

1. Anytime  
Soybeans  
Peanuts  
IMI-Corn™  
Lima beans
2. Two months after PURSUIT application:  
Snap beans
3. Four months after PURSUIT application:  
Barley  
Other types of edible beans and peas  
Rye  
Wheat
4. Eight and one-half months after PURSUIT application:  
Field corn  
Field corn grown for seed\*
5. Nine and one-half months after PURSUIT application:  
Tobacco

....

24/25

**ROTATIONAL CROP GUIDELINE (continued)**

**6. Eighteen months after PURSUIT application:**

- |         |            |
|---------|------------|
| Alfalfa | Safflower  |
| Cotton  | Sorghum    |
| Lettuce | Sunflower  |
| Oats    | Sweet corn |
| Popcorn |            |

**7. Twenty-six months after PURSUIT application:**

- Potatoes

**8. Forty months after PURSUIT application:**

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

\* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT shall be assumed by the user.

In the event of a crop loss, the bean or pea crop may be replanted.

**DO NOT** feed treated forage, hay, or straw to livestock.

**CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.**



EPA Reg. No. 241-310

**FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE,  
CRANBERRY, SMALL WHITE TYPE DRY BEANS, LIMA BEANS,  
SOUTHERN AND ENGLISH PEAS, AND WHITE LUPINS**

**FOR DISTRIBUTION AND USE IN WISCONSIN ONLY**

**THIS LABEL EXPIRES APRIL 30, 1997**

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN  
THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

#### **GENERAL INFORMATION**

PURSUIT application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

**DO NOT** apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application.

**DO NOT** apply PURSUIT postemergence after crop has begun to flower or crop injury may result.

**USE PURSUIT ONLY** if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

#### **DISCLAIMER**

**THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.**

**REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.**

#### **ROTATIONAL CROP GUIDELINE**

The following crops may be planted after applying PURSUIT as recommended:

1. Two months after PURSUIT application:  
Snap beans

\*Registered Trademark of American Cyanamid Company  
PB-17166 Rev. 3 1/23/96 SD



American Cyanamid Company  
Agricultural Products Division  
Crop Protection Chemicals Department  
Wayne, NJ 07470 ©1996

ACCEPTED  
RECOMMENDATIONS  
In EPA Letter Dated

FEB 13 1996

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.  
241-310

~~with comments~~  
In EPA Letter Dated

FEB 13 1996

EPA Reg. No. 241-310

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

## FOR USE IN SOYBEANS

### **PURSUIT PLUS SCEPTER® OR SCEPTER® DG HERBICIDES FOR SUPPRESSION OF VOLUNTEER CORN AND ENHANCED CONTROL OF SUNFLOWERS**

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT LEAFLET LABEL BEFORE USING. SEE THE PURSUIT LEAFLET LABEL FOR WORKER PROTECTION STANDARD REQUIREMENTS, USE AREA, MIXING AND APPLICATION INFORMATION, AND WEEDS CONTROLLED.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system. THIS LABEL MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION. ....

#### GENERAL INFORMATION

PURSUIT may be tank mixed with SCEPTER or SCEPTER DG for suppression of volunteer corn or for enhanced control of common sunflower in soybeans. Apply postemergence to soybeans only. In this label, SCEPTER refers to both formulations SCEPTER and SCEPTER DG. Be sure to refer to the correct label for additional information.

#### USE AREA

The application of PURSUIT plus SCEPTER or SCEPTER DG may be applied to states or portions of states described as Region 2 or Region 3 on the SCEPTER or SCEPTER DG labels, and the following counties in South Dakota: Yankton, Bon Homme, Hutchinson, McCook, Hanson, Davison, Miner, Lake and Kingsbury. Refer to the respective labels for the recommended use area. Do not use this tank mixture in North Dakota or in Minnesota north of state highway #210.

#### APPLICATION RATES

Apply the products at the following rate:

PURSUIT	4 oz/A*
AND	
SCEPTER	2 oz/A (liquid)
OR	
SCEPTER DG	0.53 oz/A**

\*At the rate of 4 ounces per acre, one gallon of PURSUIT will treat 32 acres.

\*\*At the rate of 0.53 ounces per acre, one 14 ounce soluble bag of SCEPTER DG will treat 26.4 acres.



\* Field corn may be planted 9 1/2 months following a postemergence application of PURSUIT when applied in tank-mixture with SCEPTER or SCEPTER DG at the recommended rates unless extreme drought conditions develop (less than 12 inches of rainfall or irrigation is received from two weeks prior to the date of PURSUIT application through November 15 of the same year). If the minimum rainfall requirement is not met, only field corn hybrids (IMI-Corn™) which possess tolerance or resistance to PURSUIT may be planted the spring of the year following a PURSUIT + SCEPTER application.

Use of PURSUIT and SCEPTER herbicides in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of these products and, therefore, rotational crop injury is always possible.

Except as stated above, DO NOT use this tank mixture in the same year as an application of SCEPTER, SCEPTER DG, SQUADRON®, SQUADRON® DG, TRI-SCEPT®, SCEPTER® O.T.®, STEEL™ CP, DETAIL®, PURSUIT®, PURSUIT® DG, PURSUIT® PLUS EC, CONTOUR® or RESOLVE®. Also, DO NOT use the above tank mixture in the same year as an application of a product containing chlorimuron ethyl (e.g. Classic<sup>1</sup>, Canopy<sup>1</sup>, Concert<sup>1</sup>, Gemini<sup>1</sup>, Lorox Plus<sup>1</sup>, Preview<sup>1</sup>, Reliance<sup>1</sup>, Synchrony<sup>1</sup> etc.)

®/™ Trademarks of American Cyanamid Company  
<sup>1</sup>Trademarks of E.I. duPont deNemours & Company, Inc.

PE-xxxxx 1/96 SD



Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

EPA Reg. No. 241-310

241-310 **FOR USE IN SUCCULENT PEAS, DRY EDIBLE PEAS,  
LENTILS, CHICKPEAS, AND LIMA BEANS**

PURSUIT can be used on succulent peas, dry edible peas (including chickpeas) and lentils in Washington (Adams, Asotin, Columbia, Garfield, Spokane, Walla Walla and Whitman counties), Idaho (Benewah, Clearwater, Idaho, Kootenai, Latah, Lewis, and Nez Perce counties), Oregon (Union, Umatilla and Wallowa counties) and Montana only.

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

#### GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

**DO NOT** apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application.

**USE PURSUIT ONLY** if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans. Plant peas, lentils, or lima beans at least 1/2 inch deep to reduce risk of crop injury.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

#### DISCLAIMER

**THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.**

**REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.**

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

**DO NOT** apply by air.

**DO NOT** apply PURSUIT in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT application must be drained and thoroughly cleaned with water before being used to apply other products.

**DO NOT** make more than one application of PURSUIT per year.

Allow at least 30 days between application and harvest of succulent peas and lima beans.

Allow at least 60 days between application and harvest of dry edible peas, chickpeas, and lentils.

When PURSUIT is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

instructions.

**Preemergence Applications.** Apply PURSUIT at the broadcast rate of 3 ounces per acre after planting, but prior to crop emergence.

PURSUIT may be tank-mixed with Sencor DF<sup>1</sup> or Lexone DF<sup>2</sup> to assist in the control of lambsquarters or mayweed-chamomile (dogfennel). Refer to the Sencor or Lexone label for proper application rates and restrictions.

### WEEDS CONTROLLED

PURSUIT applied at the broadcast rate of 3 ounces per acre preplant incorporated or preemergence will control:

Buckwheat, wild	Nightshade, Eastern black
Kochia	Nightshade, hairy
Lambsquarters, common*	Pigweed, redroot
Mustard, wild	Shepherdspurse
Nightshade, black	Thistle, Russian

\*preplant incorporated applications only

### ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT as recommended in this label:

- Anytime:

Chickpeas	Peas	Peanuts
Lentils	Lima beans	Soybeans
IMI-Corn™		
- Four months after PURSUIT application:
  - Barley
  - Dry edible beans
  - Snap beans (3 months)
  - Rye
  - Wheat
- Nine and one-half months after PURSUIT application:
  - Field corn
  - Field corn grown for seed\*\*
  - Tobacco

### 6. Forty months after PURSUIT application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

In the event of a crop loss, peas, lentils, chickpeas, or lima beans may be replanted.

**DO NOT** feed treated forage, hay, or straw to livestock.

\*\* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT shall be assumed by the user.

**CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.**

<sup>1</sup>Sencor is a trademark of the Parent Company of Farbenfabriken Bayer GmbH, Leverkusen

<sup>2</sup>Lexone is a trademark of E. I. duPont de Nemours and Company, Inc.

™Trademark of American Cyanamid Company

PE-17086 Rev. 4 1/10/96 SD



American Cyanamid Company  
Agricultural Products Division  
Crop Protection Chemicals Department  
Wayne, NJ 07470 ©1996

**EPA Reg. No. 241-310**

**FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE,  
CRANBERRY, AND SMALL WHITE TYPE DRY BEANS, LIMA BEANS, AND  
SOUTHERN AND ENGLISH PEAS**

**REFER TO THE MAP ON PAGE 6 FOR GEOGRAPHICAL USE AREA**

**READ LABEL CAREFULLY - USE INSTRUCTIONS VARY FOR EACH GROUP**

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN  
THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

**GENERAL INFORMATION**

PURSUIT application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

**DO NOT** apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application.

**DO NOT** apply PURSUIT postemergence after crop has begun to flower or crop injury may result.

**USE PURSUIT ONLY** if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

**DISCLAIMER**

**THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.**

**REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

**DO NOT** apply by air.

**DO NOT** apply PURSUIT in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for postemergence applications. **DO NOT** use crop oils, petroleum oils, liquid fertilizers or ~~ACCEPTED~~

**with COMMENTS**  
**In EPA Letter Dated**

**FEB 13 1996**

**Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.  
241-310**

<sup>1</sup>Dash is a trademark of BASF Corporation

<sup>2</sup>Registered Trademark of American Cyanamid Company

precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

## LIMA BEANS

**DO NOT** apply PURSUIT after June 30.

Allow at least 30 days between application and harvest.

**Preplant Incorporated Applications.** Apply PURSUIT at the broadcast rate of 2 ounces per acre to sand and loamy sand soils, or 3 ounces per acre to all other soils within 1 week before planting. Applied preplant incorporated, PURSUIT may be tank-mixed with a registered grass herbicide.

**Preemergence Applications.** Apply PURSUIT at the broadcast rate of 2 ounces per acre to sand and loamy sand soils, or 3 ounces per acre to all other soils immediately after, or up to 1 day after planting. PURSUIT may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

## NAVY, GREAT NORTHERN, RED KIDNEY, CRANBERRY, BLACK TURTLE, AND SMALL WHITE TYPE DRY BEANS

**DO NOT** apply to Domino variety black turtle beans.

**DO NOT** apply PURSUIT to sand or loamy sand soils.

**DO NOT** apply PURSUIT after June 20.

Allow at least 60 days between application and harvest.

**Preplant Incorporated Applications.** Apply PURSUIT within 1 week before planting. Applied preplant incorporated, PURSUIT may be tank-mixed with a registered grass herbicide. Refer to the following table for use rate.

Soil Type	Use Rate (oz/acre)
Sands, loamy sands	<b>DO NOT USE</b>
Sandy loams and all other soils with <2% O.M.	2
All other soils with >2% O.M.	3

## ENGLISH PEAS

**DO NOT** apply PURSUIT to sand or loamy sand soils.

Allow at least 30 days between application and harvest.

**Preplant Incorporated Applications.** Apply PURSUIT at the broadcast rate of 3 ounces per acre up to 1 week before planting. **DO NOT** apply PURSUIT to English peas that have been or will be treated with trifluralin.

**Preemergence Applications.** Apply PURSUIT at the broadcast rate of 3 ounces per acre immediately after, or up to 1 day after planting. **DO NOT** apply PURSUIT if English peas have been treated with trifluralin.

**Postemergence Applications.** Apply PURSUIT at 3 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture. **DO NOT APPLY PURSUIT POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.** If English peas have been treated with trifluralin, PURSUIT may be applied ONLY at 2 ounces per acre for suppression of black nightshade. The use of trifluralin prior to PURSUIT application may increase the likelihood and severity of crop injury.

## SOUTHERN PEAS

Allow at least 30 days between application and harvest.

**Preplant Incorporated Applications.** Apply PURSUIT at the broadcast rate of 4 ounces per acre within 1 week before planting. Applied preplant incorporated, PURSUIT may be tank-mixed with a registered grass herbicide.

**Preemergence Applications.** Apply PURSUIT at the broadcast rate of 4 ounces per acre immediately after, or up to 1 day after planting. PURSUIT may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

**Early Postemergence Applications.** Apply PURSUIT at the broadcast rate of 4 ounces per acre when peas are at least 3 inches in height. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

8/25

**WEEDS CONTROLLED**

**PURSUIT applied to low organic matter soils at the broadcast rate of 2 ounces per acre preplant incorporated or preemergence will control light to moderate infestations of:**

- Nightshade, black
- Nightshade, Eastern black
- Pigweed, redroot

**PURSUIT applied at the broadcast rate of 3 ounces per acre preplant incorporated, preemergence, or early postemergence will control:**

- Mustard, wild
- Nightshade, black
- Nightshade, Eastern black
- Nightshade, hairy
- Pigweed, redroot

Postemergence applications of 3 ounces per acre must be made to weeds less than 2 inches tall for best results.

When applied as directed at the broadcast rate of 4 ounces per acre, PURSUIT will control or reduce competition from the weeds listed below:

NOTE: C = Control, R = Reduced Competition

(The number under Maximum Leaf Stage indicates the MAXIMUM number of leaves at which weeds should be sprayed postemergence.)

**BROADLEAF WEEDS**

WEEDS CONTROLLED	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Anoda, spurred	C	2	1 - 2
Artichoke, Jerusalem		8	6 - 10
Buffalobur	C**		
Bristly starbur		2	1 - 2
Carpetweed	C		
Cocklebur, common	C**	8	1 - 8
Galinsoga	C		
Jimsonweed	C*	4	1 - 3
Kochia	C	4	1 - 3
Lambsquarters	C*	R	1 - 2
Mallow, Venice	R		
Morningglory			
entireleaf	R	2	1 - 2
ivyleaf	R	2	1 - 2
pitted	R	2	1 - 2
smallflower	C	4	1 - 3
tall	R	2	1 - 2
Mustard sp.	C	4	1 - 3
Nightshade,			
black	C	4	1 - 3
Eastern black	C	4	1 - 3
hairy	C	4	1 - 3
Pigweed,			
redroot	C	8	1 - 8
smooth	C	8	1 - 8
spiny	C	8	1 - 8
Poinsettia, wild	C		
Puncturevine	C		
Purslane, common	C		
Pusley, Florida	C		
Sida, prickly	C*		

9/25

**BROADLEAF WEEDS (Continued)**

WEEDS CONTROLLED	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Ragweed,			
common	R	4	1-3
giant	R	4	1-3
Sage, barnyard		R	1-3
Smartweed,			
ladysthumb	C	4	1-3
Pennsylvania	C	4	1-3
Spurge,			
prostrate	C	4	1-3
spotted	C	4	1-3
Sunflower, common	C*	4	1-3
Velvetleaf	C*	4	1-3
Thistle, Canada		R	1-3

\*When soil applied, common lambsquarters, jimsonweed, prickly sida, velvetleaf and common sunflower are more consistently controlled by preplant incorporated treatments.

\*\*Use soil applications for light to moderate infestations only. Must be preplant incorporated for best results.

**GRASS WEEDS\***

WEEDS CONTROLLED	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Barnyardgrass	R	3	1-3
Crabgrass,			
large	R	3	1-3
smooth	R	3	1-3
Cupgrass, woolly		3**	1-3
Foxtail,			
giant	C	6	1-6
green	C	3	1-3
robust purple	C	3	1-3
robust white	C	3	1-3
yellow	C	3	1-3
Goosegrass	R		
Johnsongrass,			
seedling	C	6	1-8
rhizome		R	1-8
Panicum,			
fall	R		
Texas	R		

10/25

**GRASS WEEDS\* (Continued)**

WEEDS CONTROLLED	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Red rice		3	1 - 3
Shattercane	R	6	1 - 8
Signalgrass, broadleaf	R	4	1 - 8

**SEDGES**

Nutsedge,			
purple	R	R	1 - 3
yellow	R	R	1 - 3

**DO NOT** count cotyledon leaves when determining weed stage of growth.

\*When soil applied to grasses, more consistent control can be obtained from preplant incorporated treatments.

\*\*PURSUIT controls emerged woolly cupgrass only.

**ROTATIONAL CROP GUIDELINE**

The following crops may be planted after applying PURSUIT as recommended in this label:

1. Anytime:
  - Soybeans
  - Peanuts
  - IMI-Corn™
  - Edible bean and pea types listed on this label (Refer to heading on page 1)
2. Two months after PURSUIT application:
  - Snap beans (DE, MD, PA and VA only)
3. Four months after PURSUIT application:
  - Barley
  - Edible bean and pea types not listed on this label (Refer to heading on page 1)
  - Rye
  - Wheat
4. Eight and one-half months after PURSUIT application
  - Field corn
  - Field corn grown for seed\*
5. Nine and one-half months after PURSUIT application:
  - Tobacco
6. Eighteen months after PURSUIT application:
 

Alfalfa	Safflower
Cotton	Sorghum
Lettuce	Sunflower
Oats	Sweet Corn
Popcorn	

7. Twenty-six months after PURSUIT application:  
Potatoes

8. Forty months after PURSUIT application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

\* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT shall be assumed by the user.

11/25

### ROTATIONAL CROP GUIDELINE (Continued)

In the event of a crop loss, the bean or pea crop may be replanted.


**DO NOT** feed treated forage, hay, or straw to livestock.

**CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.**

#### USE AREA MAP

PURSUIT can be applied to the edible legume vegetables listed only in the states or parts of states shaded in the following map:

Use area includes: South Dakota east of U.S. 81, Nebraska east of U.S. 81, Kansas east of U.S. 81, Maryland and Virginia EXCLUDING THE DELMARVA PENINSULA, and other states or portions of states shaded in the following map.



**EPA Reg. No. 241-310**

**FOR PREEMERGENCE USE IN SNAP BEANS GROWN  
IN NEW JERSEY ONLY**

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN  
THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

**GENERAL INFORMATION**

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to snap beans. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

**DO NOT** apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application.

**USE PURSUIT ONLY** if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

**DISCLAIMER**

**THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.**

**REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.**

\*Registered Trademark of American Cyanamid Company

(see reverse side)

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

**DO NOT** apply by air.

**DO NOT** apply PURSUIT in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT applications must be drained and thoroughly cleaned with water before being used to apply other products.

**DO NOT** apply PURSUIT after June 20.

Allow at least 30 days between application and harvest.

**DO NOT** make more than one application of PURSUIT per year.

When PURSUIT is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing ~~any~~ <sup>ACCEPTED</sup> prohibiting such mixtures.

In EPA Letter Dated

FEB 13 1980

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide

**Preemergence Applications.** For suppression of pigweed, nightshade and common purslane, apply PURSUIT at the broadcast rate of 1.5 ounces per acre immediately after, or up to 1 day after planting. PURSUIT may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

## ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT as recommended in this label:

1. Anytime:
  - Soybeans
  - Peanuts
  - Edible beans and peas
  - IMI-Corn™
2. Four months after PURSUIT application:
  - Barley
  - Rye
  - Wheat
3. Eight and one-half months after PURSUIT application:
  - Field corn
  - Field corn grown for seed\*
4. Nine and one-half months after PURSUIT application:
  - Tobacco
5. Eighteen months after PURSUIT application:

Alfalfa	Safflower
Cotton	Sorghum
Lettuce	Sunflower
Oats	Sweet corn
Popcorn	
6. Twenty-six months after PURSUIT application:
  - Potatoes

**ROTATIONAL CROP GUIDELINE,** a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

In the event of a crop loss, snap beans may be replanted.

**DO NOT** feed treated forage, hay, or straw to livestock.

\* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT shall be assumed by the user.

**CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.**

™Trademark of American Cyanamid Company  
PE-17085 Rev. 4 1/23/96 SD



American Cyanamid Company  
Agricultural Products Division  
Crop Protection Chemicals Department  
Wayne, NJ 07470 ©1996

**EPA Reg. No. 241-310**

**FOR USE IN ENGLISH PEAS ONLY**

**FOR DISTRIBUTION AND USE ONLY ON THE DELMARVA PENINSULA OF  
MARYLAND, DELAWARE, AND VIRGINIA**

**READ LABEL CAREFULLY**

**THIS LABEL EXPIRES APRIL 30, 1997**

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN  
THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

**GENERAL INFORMATION**

PURSUIT application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

**DO NOT** apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application.

**USE PURSUIT ONLY** if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

**DISCLAIMER**

**THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.**

**REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

**DO NOT** apply by air.

**DO NOT** apply PURSUIT in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT application must be drained and thoroughly cleaned with water before being used to apply other products.

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated**

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

241-310

of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

**DO NOT** apply PURSUIT to sand or loamy sand soils.

Allow at least 30 days between application and harvest.

**Preplant Incorporated Applications.** Apply PURSUIT at the broadcast rate of 3 ounces per acre up to 1 week before planting. **DO NOT** apply PURSUIT to English peas that have been or will be treated with trifluralin.

**Preemergence Applications.** Apply PURSUIT at the broadcast rate of 3 ounces per acre immediately after, or up to 1 day after planting. **DO NOT** apply PURSUIT if English peas have been treated with trifluralin.

### WEEDS CONTROLLED

PURSUIT applied at the broadcast rate of 3 ounces per acre preplant incorporated or preemergence will control:

Mustard, wild	Nightshade, hairy
Nightshade, black	Pigweed, redroot
Nightshade, Eastern black	

### ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT as recommended in this label:

1. Anytime:
  - Soybeans
  - Peanuts
  - IMI-Corn™
  - English peas and lima beans
2. Two months after PURSUIT application:
  - Snap beans
3. Four months after PURSUIT application:
  - Barley
  - Other types of edible beans and peas
  - Rye
  - Wheat

™Trademark of American Cyanamid Company  
PE-17181 Rev. 3 1/22/96 SD

6. Eighteen months after PURSUIT application:
  - Alfalfa
  - Cotton
  - Lettuce
  - Oats
  - Popcorn
  - Safflower
  - Sorghum
  - Sunflower
  - Sweet corn

7. Twenty-six months after PURSUIT application:
  - Potatoes

8. Forty months after PURSUIT application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

- \* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT shall be assumed by the user.

In the event of a crop loss, the bean or pea crop may be replanted.

**DO NOT** feed treated forage, hay, or straw to livestock.

**CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.**



American Cyanamid Company  
Agricultural Products Division  
Crop Protection Chemicals Department  
Wayne, NJ 07470 ©1996

**EPA Reg. No. 241-310**

**FOR USE IN RED KIDNEY TYPE DRY BEANS, WHITE LUPINS,  
LIMA BEANS, AND ENGLISH PEAS ONLY**

**FOR DISTRIBUTION AND USE ONLY IN ILLINOIS, IOWA, AND WISCONSIN**

**READ LABEL CAREFULLY - USE INSTRUCTIONS VARY FOR EACH GROUP**

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN  
THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

**GENERAL INFORMATION**

PURSUIT application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

**DO NOT** apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application. **DO NOT** apply PURSUIT postemergence after crop has begun to flower or crop injury may result.

**USE PURSUIT ONLY** if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

**DISCLAIMER**

**THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.**

**REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

**DO NOT** apply by air.

**DO NOT** apply PURSUIT in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT application must be drained and thoroughly cleaned with water before being used to apply other products.

**ACCEPTED  
with COMMENTS  
by EPA Letter Dated**

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.  
241-310

\*Registered Trademark of American Cyanamid Company

**DO NOT** make more than one application of PURSUIT per year.

Refer to the PURSUIT leaflet label for approved combinations. When PURSUIT is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

### **RED KIDNEY TYPE DRY BEANS, WHITE LUPINS AND LIMA BEANS**

**DO NOT** apply PURSUIT after June 30.

**DO NOT** apply PURSUIT to white lupins grown on sand or loamy sand soils.

Allow at least 30 days between application and harvest of lima beans.

Allow at least 60 days between application and harvest of red kidney beans or white lupins.

**Preplant Incorporated Applications.** Apply PURSUIT at the broadcast rate of 3 ounces per acre within 1 week before planting. Applied preplant incorporated, PURSUIT may be tank-mixed with a registered grass herbicide.

**Preemergence Applications.** Apply PURSUIT at the broadcast rate of 3 ounces per acre immediately after, or up to 1 day after planting. PURSUIT may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

### **ENGLISH PEAS**

Allow at least 30 days between application and harvest.

**Preplant Incorporated Applications.** Apply PURSUIT at the broadcast rate of 3 ounces per acre up to 1 week before planting. **DO NOT** apply PURSUIT to English peas that have been or will be treated with trifluralin.

ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

**DO NOT APPLY PURSUIT POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.** For optimum weed control and crop safety, apply PURSUIT prior to 5 nodes before flowering. If English peas have been treated with trifluralin, PURSUIT may be applied **ONLY** at 2 ounces per acre for suppression of black nightshade. The use of trifluralin prior to PURSUIT application may increase the likelihood and severity of crop injury.

### **WEEDS CONTROLLED**

PURSUIT applied at the broadcast rate of 3 ounces per acre preplant incorporated, preemergence, or early postemergence will control:

Mustard, wild	Nightshade, hairy
Nightshade, black	Pigweed, redroot
Nightshade, Eastern black	

Postemergence applications of 3 ounces per acre must be to weeds less than 2 inches tall for best results.

### **ROTATIONAL CROP GUIDELINE**

The following crops may be planted after applying PURSUIT as recommended in this label:

- Anytime:
  - Soybeans
  - Peanuts
  - Edible beans and pea types listed on this label (Refer to heading on page 1).
  - IMI-Corn™
- Four months after PURSUIT application:
  - Barley
  - Edible bean and pea types not listed on this label (Refer to heading on page 1).
  - Rye
  - Wheat

<sup>1</sup>Dash is a trademark of BASF Corporation

™Trademark of American Cyanamid Company

18/25

### ROTATIONAL CROP GUIDELINE (continued)

3. Eight and one-half months after PURSUIT application:

- Field corn
- Field corn grown for seed\*

4. Nine and one-half months after PURSUIT application:

- Tobacco

5. Eighteen months after PURSUIT application:

- |         |            |
|---------|------------|
| Alfalfa | Safflower  |
| Cotton  | Sorghum    |
| Lettuce | Sunflower  |
| Oats    | Sweet corn |
| Popcorn |            |

6. Twenty-six months after PURSUIT application:

- Potatoes

7. Forty months after PURSUIT application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

In the event of a crop loss, red kidney beans, white lupins, lima beans, or English peas may be replanted.

**DO NOT** feed treated forage, hay, or straw to livestock.

Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT shall be assumed by the user.

**CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.**



EPA Reg. No. 241-310

**FOR USE IN GREAT NORTHERN, RED KIDNEY AND PINTO  
TYPE DRY BEANS**

**FOR DISTRIBUTION AND USE ONLY IN NEW MEXICO**

**THIS LABEL EXPIRES APRIL 30, 1997**

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN  
THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

**GENERAL INFORMATION**

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

**DO NOT** apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application. **DO NOT** apply PURSUIT postemergence after crop has begun to flower or crop injury may result.

**USE PURSUIT ONLY** if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

**DISCLAIMER**

**THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.**

**REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

**DO NOT** apply by air.

**DO NOT** apply PURSUIT in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

**ACCEPTED  
~~with COMMENTS~~  
In EPA Letter Dated**

FEB 13 1996

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.  
241-310

\*Registered Trademark of American Cyanamid Company

**DO NOT** use crop oils, petroleum oils, liquid fertilizers or Dash.<sup>1</sup>

**DO NOT** make more than one application of PURSUIT per year. **DO NOT** apply PURSUIT after June 30. Allow at least 60 days between application and harvest.

Refer to the PURSUIT leaflet label for approved combinations. When PURSUIT is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

#### APPLICATION RATE AND TIMING

**Preemergence Applications.** Apply PURSUIT at the broadcast rate of 3 ounces per acre immediately after, or up to 1 day after planting. PURSUIT may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide. **DO NOT** apply PURSUIT on sand or loamy sand soils.

**Postemergence Applications.** Apply PURSUIT at 3 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

**DO NOT APPLY PURSUIT POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.**

Mustard, wild  
Nightshade, black  
Nightshade, Eastern black

Nightshade, hairy  
Pigweed, redroot

Postemergence applications of 3 ounces per acre must be to weeds less than 2 inches tall for best results.

#### ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT as recommended in this label:

1. Anytime:
  - Soybeans
  - Peanuts
  - Edible bean types listed on this label (Refer to heading on page 1).
  - IMI-Corn™
2. Four months after PURSUIT application:
  - Barley
  - Edible bean types not listed on this label (Refer to heading on page 1).
  - Peas
  - Rye
  - Wheat
3. Eight and one-half months after PURSUIT application:
  - Field corn
  - Field corn grown for seed\*
4. Nine and one-half months after PURSUIT application:
  - Tobacco
5. Eighteen months after PURSUIT application:

Alfalfa	Safflower
Cotton	Sorghum
Lettuce	Surflowe.
Oats	Sweet corn
Popcorn	

<sup>1</sup>Dash is a trademark of BASF Corporation

™Trademark of American Cyanamid Company

21/25

**ROTATIONAL CROP GUIDELINE (continued)**

6. Twenty-six months after PURSUIT application:  
Potatoes

7. Forty months after PURSUIT application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

In the event of a crop loss Great Northern, red kidney or pinto beans may be replanted.

**DO NOT** feed treated forage, hay, or straw to livestock.

\* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT shall be assumed by the user.

**CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.**



**EPA Reg. No. 241-310**

**FOR USE IN LIMA BEANS ONLY**

**FOR DISTRIBUTION AND USE ONLY ON THE DELMARVA PENINSULA OF  
MARYLAND, DELAWARE, AND VIRGINIA**

**READ LABEL CAREFULLY**

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN  
THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

**GENERAL INFORMATION**

PURSUIT application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT only to the legume vegetables specifically listed in this label. **DO NOT** apply PURSUIT in geographical regions outside those specifically defined in this label for each type of legume vegetable.

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

**DO NOT** apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application.

**USE PURSUIT ONLY** if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

**DISCLAIMER**

**THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.**

**REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

**DO NOT** apply by air. **ACCEPTED**  
~~with~~ **CONDITIONS**  
In EPA Letter Dated

FEB 13 1966

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended for the pesticide  
registered under EPA Reg. No.

241-310