

PM 25 241-310
Pursuit
herbicide

Supplemental
Labeling

Page 173

ACCEPTED
with COMMENTS
In EPA Letter Dated
APR 14 1997

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
241-310

EPA Reg. No. 241-310

**FOR POSTEMERGENCE USE IN SNAP BEANS GROWN
IN ARKANSAS, MISSOURI, OKLAHOMA, WEST TEXAS AND NEW MEXICO
ONLY
(see USE AREA below)**

**FOR USE ONLY BY GROWERS CONTRACTED BY ALLEN CANNING
COMPANY. THIS LABEL EXPIRES APRIL 30, 1999**

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

NOTICE

AMERICAN CYANAMID COMPANY ("CYANAMID") HAS OBTAINED A REGISTRATION AS A SERVICE TO THOSE GROWERS WHO WISH TO USE PURSUIT HERBICIDE AS AN AID IN CONTROLLING WEEDS IN EDIBLE LEGUME VEGETABLES. CYANAMID DOES NOT WARRANT THE USE OF PURSUIT ON EDIBLE LEGUME VEGETABLES. THE DECISION TO APPLY THIS HERBICIDE MUST BE MADE BY THE GROWER WITH THE UNDERSTANDING THAT REDUCED CROP GROWTH, QUALITY, YIELD AND/OR DELAYED MATURITY MAY RESULT EVEN WHEN DIRECTIONS FOR USE ARE FOLLOWED PRECISELY.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

LIABILITY WAIVER

THIS LABEL IS NOT VALID UNLESS THE CROP OWNER HAS SIGNED A CROP DAMAGE DISCLAIMER AND LIABILITY WAIVER AND INDEMNIFICATION AGREEMENT PURSUANT TO WHICH THE CROP OWNER AGREES TO DEFEND, INDEMNIFY AND HOLD CYANAMID AND ALLEN CANNING COMPANY, THEIR EMPLOYEES, OFFICERS, AGENTS, AND DEALERS, HARMLESS AGAINST ANY AND ALL CLAIMS, CAUSES OF ACTION, FINES, PENALTIES, DAMAGES INCLUDING CONSEQUENTIAL DAMAGES, LOSSES, LIABILITIES, JUDGMENTS AND EXPENSES ARISING OUT OF OR RELATING TO ANY CROP LOSS OR OTHER INJURY TO PERSON OR PROPERTY RESULTING FROM THE USE OF PURSUIT HERBICIDE ON EDIBLE LEGUMES.

USE AREA

For use in the Texas Counties of Bailey, Castro, Lamb, and Parmer, and the New Mexico Counties of Curry and Roosevelt and the states of Arkansas, Missouri and Oklahoma.

GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to snap beans. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. DO NOT apply PURSUIT

if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application.

USE PURSUIT ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT applications must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for post applications.

DO NOT use crop oils, petroleum oils, liquid fertilizers or Dash¹.

Allow at least 30 days between application and harvest.

DO NOT make more than one application of PURSUIT per year. **DO NOT** apply PURSUIT after June 30.

When PURSUIT is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

APPLICATION INSTRUCTIONS

Postemergence Applications. For suppression of pigweed and nightshade, apply PURSUIT at the broadcast rate of 1.5 ounces per acre in a tank-mix combination with Basagran¹. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

Refer to the Basagran label for proper application rates and restrictions. Always use in accordance with the more restrictive label restrictions and precautions. Use only nonionic surfactants as a spray additive.

DO NOT APPLY PURSUIT POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT as recommended in this label:

1. Anytime:
 - Soybeans
 - Peanuts
 - Edible beans and peas
 - IMI-CORN™
2. Four months after PURSUIT application:
 - Alfalfa
 - Rye
 - Wheat
3. Eight and one-half months after PURSUIT application:
 - Field corn
 - Field corn grown for seed*
4. Nine and one-half months after PURSUIT application:
 - Barley
 - Tobacco

¹Trademark of American Cyanamid Company

¹Dash and Basagran are trademarks of BASF Corporation

ROTATIONAL CROP GUIDELINE (continued)

- 5. Eighteen months after PURSUIT application:

Cotton	Safflower
Lettuce	Sorghum
Oats	Sunflower
Popcorn	Sweet corn

- 6. Twenty-six months after PURSUIT application:
 - Potatoes

- 7. Forty months after PURSUIT application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

In the event of a crop loss, snap beans may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT shall be assumed by the user.

CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.