



**Pursuit**  
herbicide

**Supplemental  
Label**

EPA Reg. No. 241-310

**FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY,  
BLACK TURTLE, CRANBERRY, AND PINTO TYPE DRY BEANS,  
AND DRY EDIBLE PEAS ONLY**

**FOR DISTRIBUTION AND USE IN NORTH DAKOTA AND  
NORTHERN MINNESOTA (see USE AREA below)**

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN  
THE PURSUIT LEAFLET LABEL BEFORE USING.**

DISCLAIMER

THIS PRODUCT WHEN USED ON EDIBLE LEGUME VEGETABLE CROPS MAY LEAD TO CROP INJURY, LOSS, OR DAMAGE. CYANAMID RECOMMENDS THAT THE USER AND OR GROWER TEST THIS PRODUCT IN ORDER TO DETERMINE ITS SUITABILITY FOR SUCH INTENDED USE. CYANAMID MAKES THIS PRODUCT AVAILABLE TO THE USER AND OR GROWER SOLELY TO THE EXTENT THE BENEFIT AND UTILITY, IN THE SOLE OPINION OF THE USER AND/OR GROWER, OUTWEIGH THE EXTENT OF POTENTIAL INJURY ASSOCIATED WITH THE USE OF THIS PRODUCT. THE DECISION TO USE OR NOT TO USE THIS HERBICIDE MUST BE MADE BY EACH INDIVIDUAL PURSUIT USER AND/OR GROWER ON THE BASIS OF POSSIBLE CROP INJURY FROM PURSUIT, THE SEVERITY OF WEED INFESTATION, THE COST OF ALTERNATIVE WEED CONTROLS, AND OTHER FACTORS. CYANAMID INTENDS THAT BECAUSE OF THE RISK OF FAILURE TO PERFORM OR CROP DAMAGE THAT ALL SUCH USE IS AT THE USER'S AND/OR GROWER'S RISK.

AMERICAN CYANAMID DISCLAIMS ANY LIABILITY FOR CLAIMS, CAUSES OF ACTION, FINES, PENALTIES, DAMAGES, INCLUDING CONSEQUENTIAL INCIDENTS AND DAMAGES, LOSSES, LIABILITIES, JUDGMENTS, AND EXPENSES ARISING OUT OF OR RELATING TO INJURY TO PERSONS, CROPS, OR PROPERTY RESULTING FROM THE USE OF PURSUIT

HERBICIDE ON EDIBLE LEGUMES CONTRARY TO THE LABEL INSTRUCTIONS. AMERICAN CYANAMID COMPANY WARRANTS ONLY THAT THE MATERIAL CONTAINED HEREIN CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL AND IS REASONABLY FIT FOR THE USE THEREIN DESCRIBED WHEN USED IN ACCORDANCE WITH THE DIRECTIONS FOR USE, SUBJECT TO THE RISKS REFERRED TO ABOVE.

CYANAMID DOES NOT MAKE OR AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED AND EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

BUYER'S EXCLUSIVE REMEDY AND AMERICAN CYANAMID'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF PURSUIT. In no case shall Cyanamid or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product.

**ACCEPTED**

AUG 25 1998

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 241-310

### GENERAL INFORMATION

PURSUIT<sup>®</sup> application methods and use rates vary for different types of edible legume vegetables. Additionally, geographical restrictions apply. Apply PURSUIT only to the legume vegetables specifically listed in this labeling. **DO NOT** apply PURSUIT in geographical regions outside those specifically defined in this labeling for each type of legume vegetable.

Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

**DO NOT** apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application.

**DO NOT** apply PURSUIT postemergence after crop has begun to flower or crop injury may result.

**USE PURSUIT ONLY** if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Under some conditions (such as heavy texture soil, high organic matter, low pH or low rainfall) PURSUIT may cause injury to subsequent planted crops. Sugar beets are particularly sensitive to PURSUIT residues in soil. PURSUIT is not recommended in crop rotations that include sugar beets. See the ROTATIONAL CROP section of this supplemental label for rotations to other sensitive crops.

### USE AREA

PURSUIT may be used in North Dakota and in Minnesota north of highway #210.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

**DO NOT** apply PURSUIT in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, potatoes, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT.

Spray equipment used for PURSUIT application must be drained and thoroughly cleaned with water before being used to apply other products.

Use only nonionic surfactants as a spray additive for postemergence applications. **DO NOT** use crop oils, petroleum oils, liquid fertilizers or Dash.

**DO NOT** make more than one application of PURSUIT per year.

When PURSUIT is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

**DO NOT** apply at any timing to Domino variety black turtle beans.

Pinto varieties UT-111 and Olathe are more sensitive to PURSUIT than other varieties.

**DO NOT** apply PURSUIT after July 10.

**DO NOT** feed treated forage, hay, or straw to livestock.

Allow at least 60 days between application and harvest.

**Postemergence Applications.** Apply PURSUIT at 2 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

**DO NOT APPLY PURSUIT POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRIFOLIATE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.**

### WEEDS CONTROLLED

**PURSUIT applied at the broadcast rate of 2 ounces per acre will control:**

- wild mustard
- black nightshade\*
- eastern black nightshade\*

\*suppression

### ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT as recommended in this label:

1. Anytime:
  - Soybeans
  - IMI-CORN<sup>®</sup> seed hybrids
  - Edible bean and pea types listed on this label (Refer to heading on page 1)

- 2. Four months after PURSUIT application:
  - Alfalfa
  - Wheat
  - Edible bean and pea types not listed on this label (Refer to heading on page 1)
- 3. Eight and one-half months after PURSUIT application:
  - Field corn
  - Field corn grown for seed\*
- 4. Nine and one-half months after PURSUIT application:
  - Barley
- 5. Eighteen months after PURSUIT application:
 

Cotton	Safflower
Lettuce	Sorghum
Oats	Sunflower
Popcorn	Sweet Corn
Rye	
- 6. Twenty-six months after PURSUIT application:
  - Potatoes
  - Flax
- 7. Forty months after PURSUIT application:
 

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a

successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

\* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT shall be assumed by the user.

**CONSULT THE PURSUIT LEAFLET LABEL FOR OTHER PRECAUTIONS.**

<sup>1</sup>Dash is a trademark of BASF Corporation  
<sup>®</sup> Registered trademark of American Cyanamid Company



American Cyanamid Company  
 North America Agricultural Products Division  
 Crop Protection Products Department  
 One Campus Drive  
 Parsippany, NJ 07054 1998

