

241-310

1-11-2001

1/13

Container Front Label

NEWPATH™ herbicide
FOR USE ON CLEARFIELD* RICE
FOR RICE SEED PROPAGATION
IN ARKANSAS, LOUISIANA, MISSISSIPPI, MISSOURI,
TEXAS AND PUERTO RICO

DO NOT USE SEED FOR FOOD OR FEED. HARVESTED SEED MUST BE DESTROYED OR IDENTITY PRESERVED AND STORED SEPARATELY FROM RICE INTENDED FOR FOOD PURPOSES IN A SECURED STORAGE FACILITY UNTIL RELEASED FOR BREEDING PURPOSES OR SEED PRODUCTION. ALL STRAW AND SEED SCREENINGS MUST BE DESTROYED.

ACTIVE INGREDIENT:

Ammonium salt of imazethapyr (±)-2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-5-ethyl-3-pyridinecarboxylic acid**..... 22.87%

INERT INGREDIENTS:..... 77.13%

TOTAL..... 100.00%

**Equivalent to 21.6% (±)-2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-5-ethyl-3-pyridinecarboxylic acid

(1 gallon contains 2.0 pounds of active ingredient as the free acid)

EPA Reg. No. 241-310

U.S. Patent No. 4,798,619

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN

CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
 (If you do not understand this label, find someone to explain it to you in detail).

In case of an emergency endangering life or property involving this product,
 call day or night 800-832-HELP.

FIRST AID

IF ON SKIN OR CLOTHING:

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF IN EYES:

Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF INHALED:

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

See Back for Additional Precautionary Statements

Net Contents: 1 Gallon
 ™/*Trademarks of BASF

1/01
 NVA 2001-05-

BASF Corporation
 26 Davis Drive
 Research Triangle Park, NC 27709

BASF

ACCEPTED
 JAN 11 2001

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 241-310

2/13

Container Back Label

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
CAUTION!**

Harmful if inhaled or absorbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing.

Personal Protective Equipment (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- long-sleeved shirt and long pants
- chemical resistant gloves
- shoes plus socks

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

DO NOT contaminate water when disposing of equipment washwaters.

DO NOT use water from NEWPATH treated field to irrigate food or feed crops which are not registered for use with NEWPATH or imazethapyr herbicides.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40CFR part 170. Refer to leaflet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. See Leaflet for Complete Directions for Use, Groundwater Advisory and Proper Handling Instructions.

STORAGE AND DISPOSAL

PROHIBITIONS:

KEEP FROM FREEZING

DO NOT store below 32°F.

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, by incineration or, if allowed by State and local authorities by burning. If burned, stay out of smoke.

See Leaflet for Disclaimer.

NEWPATH™ herbicide

FOR USE ON CLEARFIELD* RICE

FOR RICE SEED PROPAGATION
IN ARKANSAS, LOUISIANA, MISSISSIPPI, MISSOURI, TEXAS
AND PUERTO RICO

DO NOT USE SEED FOR FOOD OR FEED. HARVESTED SEED MUST BE DESTROYED OR IDENTITY PRESERVED AND STORED SEPARATELY FROM RICE INTENDED FOR FOOD PURPOSES IN A SECURED STORAGE FACILITY UNTIL RELEASED FOR BREEDING PURPOSES OR SEED PRODUCTION. ALL STRAW AND SEED SCREENINGS MUST BE DESTROYED.

THIS LABEL EXPIRES DECEMBER 31, 2002

ACTIVE INGREDIENT:

Ammonium salt of imazethapyr (±)-2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-5-ethyl-3-pyridinecarboxylic acid** 22.87%

INERT INGREDIENTS: 77.13%

TOTAL 100.00%

**Equivalent to 21.6% (±)-2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-5-ethyl-3-pyridinecarboxylic acid (1 gallon contains 2.0 pounds of active ingredient as the free acid)

EPA Reg. No. 241-310

U.S. Patent No. 4,798,619

KEEP OUT OF REACH OF CHILDREN

CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand this label, find someone to explain it to you in detail).

FIRST AID

IF ON SKIN OR

CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

**In case of an emergency endangering life or property involving this product,
call day or night 800-832-HELP.**

See inside for additional precautionary statements

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

CAUTION!

Harmful if inhaled or absorbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing.

Personal Protective Equipment (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- long-sleeved shirt and long pants
- chemical resistant gloves
- shoes plus socks

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

DO NOT contaminate water when disposing of equipment washwaters.

DO NOT use water from NEWPATH treated field to irrigate food or feed crops which are not registered for use with NEWPATH or imazethapyr herbicides.

Groundwater Advisory and Proper Handling Instructions

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

This product may not be mixed or loaded within 50 feet of any wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes or reservoirs. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or properly diked mixing/loading areas.

Operations that involve mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well are prohibited unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or washwater, and rainwater that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above specific minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site. States may have in effect additional requirements regarding wellhead setbacks and operational containment.

DO NOT apply this product through any type of irrigation system.

Product must be used in a manner which will prevent back siphoning in wells, spills or improper disposal of excess pesticide, spray mixtures or rinsates.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

This labeling must be in the possession of the user at the time of pesticide application.

6/13

Observe all cautions and limitations on this label and on the labels of products used in combination with NEWPATH. Do not use NEWPATH other than in accordance with the instructions set forth on this label. The use of NEWPATH not consistent with this label may result in injury to crops. Keep containers closed to avoid spills and contamination.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours. Exception: if the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls
- chemical resistant gloves
- shoes plus socks

STORAGE AND DISPOSAL

PROHIBITIONS:

KEEP FROM FREEZING

DO NOT store below 32°F.

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, by incineration or, if allowed by State and local authorities by burning. If burned, stay out of smoke.

CONTAINER DISPOSAL FOR BULK: Return empty container for reuse.

SEED DISPOSAL: To dispose of out-of-date or otherwise unmarketable seed from plants which have been treated with NEWPATH herbicide, broadcast and lightly incorporate in field soils using disc or other suitable implement. Any resulting crop may be destroyed by chemical or mechanical means. Alternately, seed may be destroyed by deep burial, incineration or landfill disposal.

DISCLAIMER

The label instructions for the use of this product reflect the opinion of experts based on research and field use. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, herbicide resistant plant populations, or the use of, or application of the product contrary to label instructions, all of which are beyond the control of BASF Corporation (BASF). All such risks shall be assumed by the user.

BASF shall not be responsible for losses or damages resulting from use of this product in any manner not set forth on this label. User assumes all risks associated with the use of this product in any manner not specifically set forth on this label.

BASF warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use, subject to the risks referred to above. **BASF DOES NOT MAKE OR AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED AND EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.**

BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF NEWPATH. In no case shall BASF or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product.

Uses With Other Products (Tank-mixes)

If this product is used in combination with any other product except as specifically recommended in writing by BASF, then BASF shall have no liability for any loss, damage or injury arising out of its use in any such combination not so specifically recommended. If used in combination recommended by BASF Corporation, the liability of BASF shall in no manner extend to any damage, loss or injury not directly caused by the inclusion of the BASF Corporation product in such combination use, and in any event shall be limited to return of the amount of the purchase price of the BASF product.

Use On CLEARFIELD Rice

Licensed for use on ATCC 75295, ATCC 97523, PTA-902, PTA-903, PTA-904, PTA-905, PTA-906, PTA-907, or PTA-908 rice and derivatives and progeny. The purchase of this herbicide includes a sublicense to practice the processes claimed by United States Patent Nos. 5,952,553; 5,773,704; 5,773,703; 5,736,629; 5,545,822; and additional pending applications drawn to such rice or the use of such rice lines, by applying this herbicide to fields planted with rice seed purchased in a container bearing the legend "Licensed ATCC 75295, ATCC 97523, PTA-902, PTA-903, PTA-904, PTA-905, PTA-906, PTA-907, or PTA-908 Rice" in full accordance with the directions printed on this label.

GENERAL INFORMATION

NEWPATH herbicide can be applied preplant or preplant incorporated up to 14 days prior to rice planting, preemergence and postemergence for control of susceptible rice "segregates" or undesirable weedy varietal rice plants that do not contain the CLEARFIELD rice tolerance trait in only CLEARFIELD rice (imidazolinone tolerant rice). Apply NEWPATH only on selected rice varieties labeled as "CLEARFIELD" and warranted by the seed company to possess tolerance to direct application of certain imidazolinone herbicides. DO NOT apply NEWPATH to rice varieties which lack tolerance to imidazolinone herbicides as NEWPATH will kill all non-imidazolinone tolerant varieties. This product should not be applied in fields intentionally planted with non-imidazolinone tolerant rice. Contact your seed supplier, chemical dealer or BASF to obtain information regarding imidazolinone tolerant rice varieties.

NEWPATH kills plants by root and/or foliage uptake and rapid translocation to the growing points. Adequate soil moisture is important for optimum NEWPATH activity. When adequate soil moisture is present, NEWPATH will provide residual control of susceptible germinating plants; activity on established plants will depend on the species and the location of its root system in the soil. Activity of NEWPATH on susceptible plants is usually visible in 10-14 days.

Crops growing under stressful environmental conditions can exhibit various injury symptoms which may be more pronounced if herbicides are used. Rice plants treated with NEWPATH may exhibit a slight height reduction. Such effects occur infrequently and are temporary. Normal growth and appearance should resume within 2-4 weeks.

NEWPATH can be applied to CLEARFIELD rice in conventional or conservation tillage and broadcast or drill-seeded (dry planted) systems. The use rate and timing of application may vary with these production systems.

Use of NEWPATH herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible. Under some conditions (such as heavy texture soil, high organic matter or low pH) NEWPATH may cause injury to subsequent planted crops. Vegetable crops, cotton and non-CLEARFIELD rice are sensitive to NEWPATH residues in the soil.

Replanting: If replanting is necessary in a field previously treated with NEWPATH, the field may be replanted to CLEARFIELD rice, soybeans, peanuts or CLEARFIELD corn (imidazolinone resistant/tolerant corn), lima beans or Southern peas. Rework the soil no deeper than the treated zone. Do not apply a second treatment of NEWPATH or other imidazolinone containing product.

MIXING INSTRUCTIONS

POSTEMERGENCE APPLICATIONS OF NEWPATH REQUIRE THE ADDITION OF AN ADJUVANT.

SURFACTANTS: Use a non-ionic surfactant containing at least 80% active ingredient. Apply the surfactant at the rate of 0.25% v/v (1 quart per 100 gallons of spray solution).

9/13

Fill the spray tank one-half to three-quarters full with clean water. Use a calibrated measuring device to measure the required amount of NEWPATH. Add NEWPATH to the spray tank while agitating. Add adjuvants and fill the remainder of the tank with water.

To avoid injury to sensitive crops, spray equipment used for NEWPATH applications must be drained and thoroughly cleaned with water before being used to apply other products.

SPRAYING INSTRUCTIONS

DO NOT apply when wind velocity is greater than 10 mph, when temperature inversion conditions exist, or when spray may be carried to sensitive crops. Sensitive crops include, but are not limited to, leafy vegetables, cotton and non-CLEARFIELD rice varieties.

GROUND APPLICATIONS

Uniformly apply with properly calibrated ground equipment in 10 or more gallons of water per acre. A spray pressure of 20 to 40 psi is recommended.

Adjust the boom height to ensure proper coverage of foliage (according to the manufacturer's recommendation). Use only flat-fan nozzle tips for postemergence applications.

Avoid overlaps when spraying.

AERIAL APPLICATION

NEWPATH herbicide may be applied by air only to CLEARFIELD rice varieties. DO NOT apply by air to other crops.

Uniformly apply with properly calibrated aerial equipment in 5 or more gallons of water per acre. When applied POSTEMERGENCE, the addition of a non-ionic surfactant is required for optimum control. Apply a non-ionic surfactant at the rate of 1 quart per 100 gallons of spray solution. (See instructions under APPLICATION INFORMATION - POSTEMERGENCE).

To avoid injury to sensitive crops from drift, aerial applicators must adhere to the following SPECIAL AERIAL USE DIRECTIONS AND PRECAUTIONS.

- Nozzle height above ground must be a maximum of 10 feet.
- Nozzles must be pointed toward the rear of the aircraft. The downward angle of the nozzle should not be greater than 20 degrees.
- To minimize wing-tip vortex roll, nozzles or spray boom must not be located any closer to end of wing or rotor than three-fourths the distance from the center of the aircraft.
- Use a maximum spray pressure of 40 psi.
- A buffer zone must be established between the area to be sprayed and sensitive crops.
- DO NOT spray when wind velocity is greater than 5 mph. Coarse sprays (larger droplets) are less likely to drift.

Applicator is responsible for any loss or damage which results from spraying NEWPATH in a manner other than recommended in this label. In addition, applicator must follow all applicable state and local regulations and ordinances in regard to spraying.

10/13

APPLICATION INFORMATION

NEWPATH offers flexibility in both timing and method of application. Use this product in dry seeded rice in either conservation or conventional tillage systems. NEWPATH can be applied preplant incorporated, preemergence and postemergence (preflood).

SOIL APPLICATIONS

Preplant Incorporated Application

NEWPATH can be applied as a preplant incorporated treatment up to 14 days prior to rice planting. When applied preplant incorporated, NEWPATH should be thoroughly incorporated (at least single pass with a field cultivator, no disks) to a depth of 1 to 2 inches.

Preemergence Application

NEWPATH can be applied as a preemergence treatment up to 14 days prior to rice planting and prior to rice emergence.

Adequate soil moisture is required for optimum herbicide activation for all methods of soil application. If sufficient levels of precipitation DO NOT occur within 48 hours after application, flood irrigation (flush) is recommended to move NEWPATH into the germination zone for maximum activity. The amount of rainfall or irrigation required following application depends on existing soil moisture, soil texture and organic matter content. Sufficient water to moisten the soil to a depth of 2 inches is normally adequate. When adequate moisture is received after dry conditions, NEWPATH will provide residual control of susceptible germinating plants; activity on established plants will depend on the location of its root system in the soil.

NEWPATH controls plants by root uptake and translocation to the growing points where it inhibits growth. Susceptible plants may emerge but growth will stop and the plants will become noncompetitive with the CLEARFIELD rice.

FOLIAR APPLICATIONS

Postemergence (Preflood) Application

NEWPATH herbicide can be applied postemergence (preflood) when CLEARFIELD rice is in the 3-5 leaf stage. DO NOT apply NEWPATH when rice is in the spike to less than 3 leaf stage. Apply no later than 2 days prior to permanent flood. DO NOT apply into standing water. Moist soils are acceptable.

Apply NEWPATH herbicide as a postemergence treatment when plants are actively growing but before tillering and/or before they exceed a height of 6 inches (4-5 leaf stage). If possible, delay application until the majority of the plants are at the recommended growth stage. Application timing should be within the acceptable rice growth stage.

Do not apply NEWPATH to rice growing under stress induced by adverse conditions such as cool temperatures, saline soil, nutrient deficiency and disease pressure, or to rice when conditions are forecast that stress rice, especially cool temperatures. If applied under these conditions stunting and/or yellowing may occur.

An adjuvant must be added to the spray solution for optimum control activity. See the ADJUVANTS section under MIXING INSTRUCTIONS for specific instructions.

When NEWPATH is applied postemergence, absorption will occur through both the roots and foliage. Susceptible plants stop growing and either die or become noncompetitive with the crop. Activity of NEWPATH on susceptible plants is usually visible in 10-14 days.

NEWPATH should be applied a minimum of one hour before rainfall and/or at least 48 hr prior to permanent flooding.

USE RATE

For control of susceptible rice "segregates" or undesirable weedy varietal rice plants that do not contain the CLEARFIELD rice tolerance trait, sequentially apply NEWPATH at a broadcast rate of 3 fluid ounces of product per acre preplant, preplant incorporated or preemergence followed by a postemergence application of 3 fluid ounces of product per acre applied to CLEARFIELD rice at the 3- to 5-leaf growth stage. Apply 6 fluid ounces per acre during the growing season. Use this product ONLY on CLEARFIELD rice varieties as NEWPATH herbicide will kill all non-imidazolinone tolerant varieties.

INSTRUCTIONS FOR SEED HANDLING, STORAGE AND USE

Seed from treated plants must be held in secure storage until used for breeding or producing imidazolinone tolerant rice seed, or destroyed. Seed from treated plants must be labelled as follows: "Do Not Use for Feed or Food Purposes. Store Away from Feed or Food Stuffs." In addition, label the seed with the "Seed Disposal" statements found in the "Storage and Disposal" section of this label. DO NOT use rice, any rice processed commodities or rice straw treated with NEWPATH for food or feed. Destroy rice straw by incorporating it into the soil, or by composting, followed by incorporating its compost into soil.

12/13

ROTATIONAL CROP GUIDELINE

The following rotational crops may be planted after applying NEWPATH at the recommended rate: (Planting earlier than the recommended interval may result in crop injury).

1. Anytime

| | |
|---------------------------|----------|
| Lima beans | Soybeans |
| Southern peas | Peanuts |
| CLEARFIELD corn hybrids | |
| CLEARFIELD rice varieties | |
2. Four months after NEWPATH application:
 - Alfalfa
 - Rye
 - Wheat
 - Edible beans and peas (other than lima beans and Southern peas)
3. Eight and one-half months after NEWPATH application:
 - Field corn
 - Field corn grown for seed
4. Nine and one-half months after NEWPATH application:
 - Barley
 - Tobacco
5. Eighteen months after NEWPATH application:

| | |
|-----------------------------------|------------|
| Cotton | Safflower |
| Lettuce | Sorghum |
| Oats | Sunflower |
| Popcorn | Sweet corn |
| Rice (non-imidazolinone tolerant) | |
6. Twenty-six months after NEWPATH application.
 - Potatoes
 - Flax
7. Forty months after NEWPATH application.
 - All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE*

*Following forty months after a NEWPATH application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

Use of NEWPATH herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

For additional information regarding the use of NEWPATH herbicide, call telephone no. 800-942-0090.

13/13

1603

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709

BASF

™*Trademarks of BASF
©BASF Corporation
NVA 2001-04-