

FEB 15 2002

Ms. Rebecca L. Johnston
 Manager, Global Regulatory Affairs
 BASF CORPORATION
 P. O. Box 400
 Princeton, NJ 08543-044

SUBJECT: Label Amendment
 PURSUIT DG Herbicide, EPA Reg. No. 241-350
 PURSUIT Herbicide, EPA Reg. No. 241-310
 Your Application Dated January 11 2002

The above mentioned amendment submitted in connection with registration under Federal Insecticide, Fungicide, and Rodenticide Act, as amended is acceptable provided you make the label changes listed below:


1. On page 1 in label title list "peas and beans" instead of "Edible Legumes"
2. On page 27 and 35 add "succulent lima bean".
3. On page 28 move "succulent lima beans" to end of sentence.
4. On pages 32, 33, 34, 36 and 37 add "Do not make more than one application of Pursuit per year".
5. On page 37 specify "Dry or succulent chick peas" - if "dry" change "30" to "60 days" between applications.

CONCURRENCES

SYMBOL ▶	7505C							
SURNAME ▶	MINOR, E.							
DATE ▶	Feb 13, 2002							

A stamped copy is enclosed for your records. Please submit one (1) final printed copy for the above mentioned label before releasing the product for shipment. The amended labeling supersedes all previously accepted ones.

Sincerely,



James Tompkins
Product Manager, (25)
Herbicide Branch
Registration Division (7505C)

Enclosure

3/62

PURSUIT®
herbicide

128923
922

1st pass beans

**FOR USE IN ALFALFA, EDIBLE LEGUMES, FIELD CORN
(Apply Only on CLEARFIELD™ corn hybrids),
PEANUTS, AND SOYBEANS AND PEANUTS**

ACTIVE INGREDIENT:

Ammonium salt of imazethapyr (+)-2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-5-ethyl-3-pyridinecarboxylic acid** 22.87%

INERT INGREDIENTS 77.13%

TOTAL 100.00%

**Equivalent to 21.6% (+)-2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-5-ethyl-3-pyridinecarboxylic acid

(1 gallon contains 2.0 pounds of active ingredient as the free acid)

U.S. Patent No. 4,798,619
EPA Reg. No. 241-310

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN

CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand this label, find someone to explain it to you in detail).

STATEMENT OF PRACTICAL TREATMENT FIRST AID

IF ON SKIN OR

CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF IN EYES:

Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF INHALED:

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

**In case of an emergency endangering life or property involving this product,
call day or night 800-832-HELP.**

ACCEPTED
with COMMENTS
In EPA Letter Dated

See inside for additional precautionary statements

FEB 15 2002
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
241-310

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709

BASF

Net Contents: 1 Gallon (3.78 liters)

®/™ Trademarks of BASF

TABLE OF CONTENTS

4/62

<u>Page</u>	
Ingredients.....	1
First Aid	1
Precautionary Statements	4
Environmental Hazards	4
Directions for Use	5
Agricultural Use Requirements.....	5
Storage and Disposal.....	6
Disclaimer	6
General Information	8
Use Area Restrictions.....	9
Mixing Instructions	9
Tank mix Combinations With Other Herbicides.....	11
Spraying Instructions	11
Ground Applications.....	11
PURSUIT Applications With a Low Volume Sprayer	11
Aerial Applications	13
Application Information.....	13
Postemergence	13
No-Till/Minimum Tillage and Double crop Soybeans	14
Soil Applications.....	14
Soil Applications with Liquid Fertilizers	15
Preemergence (Surface Applications).....	15
No-Till or Reduced Tillage	15
Preplant Incorporated Applications.....	15
Federal Conservation Reserve Program/Agricultural Reserve Program Land.....	16
Directions for Use	16
Alfalfa	
Use Rate	17
Weeds Controlled.....	17
Tank Mixture Combinations With Other Herbicides.....	20
Application Information.....	21
CLEARFIELD corn	
Use Rate	22
Weeds Controlled.....	22
Tank-mix Herbicide Combinations With PURSUIT	25
Use in North Dakota and Northern Minnesota	26
Navy, Great Northern, Red Kidney, Black Turtle, Cranberry, Pinto, Lima, and Small White Type Dry Beans, Lentils, White Lupins, Chickpeas (Garbanzo Beans), Dry Edible Peas, English and Southern Peas in the states east of and including: North Dakota, South Dakota, Wyoming, Colorado and New Mexico (except states east of and including: Connecticut, Massachusetts, and Vermont).	
Application Instructions.....	27
Weeds Controlled.....	28
Red Kidney Beans in California	
Application Rate and Timing	32
Weeds Controlled.....	32
Snap Beans in Alabama, Florida, Georgia, Illinois, Minnesota, Michigan, New Jersey and Wisconsin	
Application Instructions.....	33
Weeds Suppressed.....	33
Snap Beans in Arkansas, Missouri, Oklahoma, Texas and New Mexico	
Application Instructions.....	34
Weeds Suppressed.....	34

Succulent Peas, Dry Edible Peas, Lentils, Chickpeas and Lima Beans in Idaho, Montana, Nevada, Oregon, Utah, and Washington

 Application Rate and Timing 35

 Weeds Controlled..... 36

Chickpeas in Arizona and California

 Application Rate and Timing 37

 Weeds Controlled..... 37

Peanuts

 Use Rate 38

 Weeds Controlled..... 38

 Weeds Controlled by Sequential Applications of PURSUIT..... 41

 Herbicide Combinations 42

Soybeans

 Use Rate 44

 Weeds Controlled..... 44

 Herbicide Combinations 47

 Use in North Dakota and Northern Minnesota 53

Rotational Crop Guideline 54

Exceptions to Rotational Crop Guideline..... 56

Precautions 58

6/62

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

CAUTION!

Harmful if inhaled or absorbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing.

Personal Protective Equipment (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow instructions for category A on an EPA chemical resistant category selection chart.

Applicators and other handlers must wear:

- long-sleeved shirt and long pants
- ~~waterproof~~ chemical-resistant gloves, such as butyl rubber ≥ 14 mils, or natural rubber ≥ 14 mils, or neoprene rubber ≥ 14 mils, or nitrile rubber ≥ 14 mils.
- shoes plus socks

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

DO NOT contaminate water when disposing of equipment washwaters.

Groundwater Advisory and Proper Handling Instructions

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

This product may not be mixed or loaded within 50 feet of any wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes or reservoirs. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or properly diked mixing/loading areas.

Operations that involve mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well are prohibited unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or washwater, and rainwater that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means

7/62

the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest

pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above specific minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site. States may have in effect additional requirements regarding wellhead setbacks and operational containment.

DO NOT apply this product through any type of irrigation system.

Product must be used in a manner which will prevent back siphoning in wells, spills or improper disposal of excess pesticide, spray mixture.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

This labeling must be in the possession of the user at the time of pesticide application.

Observe all cautions and limitations on this label and on the labels of products used in combination with PURSUIT. Do not use PURSUIT other than in accordance with the instructions set forth on this label. The use of PURSUIT not consistent with this label may result in injury to crops. Keep containers closed to avoid spills and contamination.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours. Exception: if the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls
- ~~waterproof~~ chemical-resistant gloves, such as butyl rubber ≥ 14 mils, or natural rubber ≥ 14 mils, or neoprene rubber ≥ 14 mils, or nitrile rubber ≥ 14 mils.
- shoes plus socks

8/62

STORAGE AND DISPOSAL

PROHIBITIONS:

KEEP FROM FREEZING

DO NOT store below 32°F.

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, by incineration or, if allowed by State and local authorities by burning. If burned, stay out of smoke.

CONTAINER DISPOSAL FOR BULK: Return empty container for reuse.

DISCLAIMER

The label instructions for the use of this product reflect the opinion of experts based on research and field use. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, herbicide resistant weed populations, or the use of, or application of the product contrary to label instructions, all of which are beyond the control of BASF Corporation (BASF). All such risks shall be assumed by the user.

BASF shall not be responsible for losses or damages resulting from use of this product in any manner not set forth on this label. User assumes all risks associated with the use of this product in any manner not specifically set forth on this label.

BASF warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use, subject to the risks referred to above. **BASF DOES NOT MAKE OR AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED AND EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.**

BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF PURSUIT. In no case shall BASF or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product.

Edible Bean User/Grower

THIS PRODUCT WHEN USED ON EDIBLE LEGUME CROPS MAY LEAD TO CROP INJURY, LOSS, OR DAMAGE. BASF RECOMMENDS THAT THE USER AND/OR GROWER TEST THE PRODUCT IN ORDER TO DETERMINE ITS SUITABILITY FOR SUCH INTENDED USE. BASF MAKES THIS PRODUCT AVAILABLE TO THE USER AND/OR GROWER SOLELY TO THE EXTENT THE BENEFIT AND UTILITY, IN THE SOLE OPINION OF THE USER AND/OR GROWER, OUTWEIGH THE EXTENT OF POTENTIAL INJURY ASSOCIATED WITH THE USE OF THIS PRODUCT. THE DECISION TO USE OR NOT TO USE THIS HERBICIDE MUST BE MADE BY EACH

9/62

INDIVIDUAL PURSUIT USER AND/OR GROWER ON THE BASIS OF POSSIBLE CROP INJURY FROM PURSUIT, THE SEVERITY OF WEED INFESTATION, THE COST OF ALTERNATIVE WEED CONTROLS, AND OTHER FACTORS. BASF INTENDS THAT BECAUSE OF THE RISK OF FAILURE TO PERFORM OR CROP DAMAGE THAT ALL SUCH USE IS AT THE USER'S AND/OR GROWER'S RISK.

BASF DISCLAIMS ANY LIABILITY FOR CLAIMS, CAUSES OF ACTION, FINES, PENALTIES, DAMAGES, INCLUDING CONSEQUENTIAL INCIDENTS AND DAMAGES, LOSSES, LIABILITIES, JUDGMENTS, AND EXPENSES ARISING OUT OF OR RELATING TO INJURY TO PERSONS, CROPS, OR PROPERTY RESULTING FROM THE USE OF PURSUIT HERBICIDE ON EDIBLE LEGUMES CONTRARY TO THE LABEL INSTRUCTIONS.

Uses With Other Products (Tank-mixes)

If this product is used in combination with any other product except as specifically recommended in writing by BASF, then BASF shall have no liability for any loss, damage or injury arising out of its use in any such combination not so specifically recommended. If used in combination recommended by BASF, the liability of BASF shall in no manner extend to any damage, loss or injury not directly caused by the inclusion of the BASF product in such combination use, and in any event shall be limited to return of the amount of the purchase price of the BASF product.

10/62

GENERAL INFORMATION

~~PURSUIT can be applied as an early preplant, preplant incorporated, preemergence, at ground cracking (peanuts) or early postemergence. The application method of choice will depend on the anticipated weed spectrum and the preference of the applicator.~~

PURSUIT kills weeds by root and/or foliage uptake and rapid translocation to the growing points. Adequate soil moisture is important for optimum PURSUIT activity. When adequate soil moisture is present, PURSUIT will provide residual control of susceptible germinating weeds; activity on established weeds will depend on the weed species and the location of its root system in the soil.

~~Apply PURSUIT herbicide only on selected field corn hybrids (CLEARFIELD corn) warranted by the seed company to possess resistance/tolerance to direct application of PURSUIT. DO NOT apply PURSUIT to corn hybrids which lack resistance/tolerance to PURSUIT herbicide. Contact your seed supplier, chemical dealer or BASF to obtain information regarding CLEARFIELD corn seed hybrids.~~

~~Crops growing under stressful environmental conditions can exhibit various injury symptoms which may be more pronounced if herbicides are used. Corn plants treated with PURSUIT may exhibit yellowing on new growth. Such effects occur infrequently and are temporary. Normal growth and appearance should resume within 1 to 2 weeks.~~

Occasionally, soybean internode shortening and/or temporary yellowing of crop plants may occur following PURSUIT applications. These effects occur infrequently and are temporary. Normal growth and appearance should resume within 1 to 2 weeks. These effects will not affect crop yields.

When organo-phosphate (such as Lorsban¹) or carbamate insecticides are tank-mixed with PURSUIT herbicide, temporary injury may result to the treated crops.

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible. Under some conditions (such as heavy texture soil, high organic matter, low pH or low rainfall) PURSUIT may cause injury to subsequent planted crops. Vegetable crops and particularly sugar beets are sensitive to PURSUIT residues in the soil.

Naturally occurring biotypes* of some of the weeds listed on this label may not be effectively controlled by this and/or other products with either the ALS/AHAS enzyme inhibiting mode of action. Other herbicides with the ALS/AHAS enzyme inhibiting mode of action include the sulfonylureas (e.g., Accent², Basise², Classic², ConcertHarmony GT²², Spirit-Exceed³³, Permit⁴, Pinnaele², etc.), the sulfonamides (e.g., Broadstrike¹FirstRate¹, etc.) and the pyrimidyl benzoates (e.g. Staple², etc.). If naturally occurring ALS/AHAS resistant biotypes are present in a field, Pursuit and/or any other ALS/AHAS enzyme inhibiting mode of action herbicide should be tank-mixed or applied sequentially with an appropriate registered herbicide having a different mode of action to ensure control.

* A weed biotype is a naturally occurring plant within a given species that has a slightly different, but distinct, genetic makeup from other plants.

Replanting: If replanting is necessary in a field previously treated with PURSUIT, the field may be replanted to soybeans, peanuts or CLEARFIELD corn (imidazolinone resistant/tolerant corn), lima beans or Southern peas. Rework the soil no deeper than the treated zone. Do not apply a second treatment of PURSUIT.

CLEARFIELD CORN

Apply PURSUIT herbicide only on selected field corn hybrids (CLEARFIELD corn) warranted by the seed company to possess resistance/tolerance to direct application of PURSUIT. DO NOT apply

PURSUIT to corn hybrids which lack resistance/tolerance to PURSUIT herbicide. Contact your seed supplier, chemical dealer or BASF to obtain information regarding CLEARFIELD corn hybrids.

Crops growing under stressful environmental conditions can exhibit various injury symptoms which may be more pronounced if herbicides are used. Corn plants treated with PURSUIT may exhibit yellowing on new growth. Such effects occur infrequently and are temporary. Normal growth and appearance should resume within 1 to 2 weeks.

EDIBLE LEGUME VEGETABLES

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. DO NOT apply PURSUIT if planting is delayed and chance of frost prior to maturity is likely.

USE PURSUIT ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans. Plant peas, lentils or lima beans at least 1/2 inch deep to reduce risk of crop injury.

DO NOT apply PURSUIT if cold and/or wet conditions are present or predicted to occur within one week of application. DO NOT apply PURSUIT postemergence after crop has begun to flower or crop injury may result. (Refer to specific legume vegetable crop for specific application timings recommended).

USE AREA RESTRICTIONS

In New York State - Not for Sale or Use on Long Island.

PURSUIT may be used in the following states: ~~Colorado, New Mexico, North Dakota, Oklahoma, South Dakota, Texas, Wyoming (following counties only: Sublette, Lincoln, Uinta, Sweetwater, Carbon, Natrona, Albany, Platte, Laramie, Goshen, Converse, and Niobrara)~~ and states east of these states.

MIXING INSTRUCTIONS

POSTEMERGENCE APPLICATIONS OF PURSUIT REQUIRE THE ADDITION OF AN ADJUVANT AND A FERTILIZER SOLUTION.

NOTE: Fertilizer solutions may not be used in the state of California.

I. ADJUVANTS

CROP OIL CONCENTRATE: A petroleum or vegetable seed based oil concentrate (such as ~~SUN-IT-II~~) may be used. Methylated seed oils, such as ~~SUN-IT-II~~ are recommended when weeds are under moisture or temperature stress. Use ~~SUN-IT-II~~ or other methylated seed oils at the rate of 1.0% v/v (1 gallon per 100 gallons of spray solution), or use a crop oil concentrate at 1.25% v/v (1.25 gallons per 100 gallons of spray solution). **DO NOT include a CROP OIL CONCENTRATE when applying PURSUIT to edible legume vegetable crops.**

OR

SURFACTANTS: Use a non-ionic surfactant containing at least 80% active ingredient. Apply the surfactant at the rate of 0.25% v/v (1 quart per 100 gallons of spray solution). An organo-silicone surfactant or dry surfactant may be used in place of a non-ionic surfactant.

AND

(All States Except California)

11/62

12/62

II. FERTILIZER SOLUTION

Recommended nitrogen based fertilizers include liquid fertilizers (such as 28%N, 32%N or 10-34-0) may be applied at the rate of 1.25 to 2.5 gallons per 100 gallons of spray solution. Use the higher rate when weeds are under moisture or temperature stress. Instead of a liquid fertilizer, spray grade ammonium sulfate may be used at the rate of 12-15 lbs. per 100 gallons of spray solution.

Note: ~~Liquid fertilizer solutions~~ is ~~are not~~ required in PURSUIT applications in use areas south of Interstate highway 40, except in the states of Texas, New Mexico and Oklahoma.

Fill the spray tank one-half full with clean water. Use a calibrated measuring device to measure the required amount of PURSUIT. Add PURSUIT to the spray tank while agitating. Add adjuvants and fill the remainder of the tank with water.

13/62

TANK MIX COMBINATIONS WITH OTHER HERBICIDES

If other herbicides are tank-mixed with PURSUIT, while agitating, add components in the following order:

- 1) Fill spray tank 1/2 full with clean water.
- 2) Add soluble packet products and thoroughly mix.
- 3) Add WP (wettable powder), DG (dispersible granule), DF (dry flowable) or liquid flowable formulations not in soluble packets.
- 4) Add PURSUIT and thoroughly mix.
- 5) Add other aqueous solution products.
- 6) Add EC (emulsifiable concentrate) products.
- 7) Add surfactant or crop oil to the spray tank.
- 8) Add liquid fertilizer.
- 9) While agitating, fill the remainder of the tank with water.

~~If dicamba sprayer contamination is possible, refer to the appropriate dicamba label (Banvel⁷, Clarity⁷, or Marksman⁷) for sprayer cleanup directions. Clean the sprayer prior to PURSUIT applications to avoid dicamba injury to soybeans.~~

To avoid injury to sensitive crops, spray equipment used for PURSUIT applications must be drained and thoroughly cleaned with water before being used to apply other products.

When PURSUIT is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

SPRAYING INSTRUCTIONS

DO NOT apply when wind velocity is greater than 10 mph, or when spray may be carried to sensitive crops. Sensitive crops include, but are not limited to, leafy vegetables and sugar beets.

GROUND APPLICATIONS

Uniformly apply with properly calibrated ground equipment in 10 or more gallons of water per acre. A spray pressure of 20 to 40 psi is recommended.

To ensure thorough coverage, use a minimum of 20 gallons of water per acre when applying PURSUIT to minimum or no-till crops. Use higher gallonage for fields with dense vegetation or heavy crop residues. Adjust the boom height to ensure proper coverage of weed foliage (according to the manufacturer's recommendation). Use only flat-fan nozzle tips for postemergence applications.

Avoid overlaps when spraying.

PURSUIT APPLICATIONS WITH A LOW VOLUME SPRAYER

PURSUIT may be applied to soybeans with a low volume (Spra-Coupe-type) sprayer. When applying PURSUIT with a low volume sprayer, spray the weeds before they reach the maximum size listed in this label. Adequate control of weeds is dependent upon good spray coverage of the weeds. The sprayer must be calibrated to deliver the recommended spray volume and pressure to ensure adequate spray coverage of the weeds.

When applying PURSUIT with a low volume sprayer, apply a minimum of 10 gallons per acre of spray solution with a nozzle pressure between 40-60 psi for optimum coverage. When spraying

combinations including Banvel or dicamba containing products on CLEARFIELD corn, do not exceed 40 psi sprayer pressure.

14/62

15/62

AERIAL APPLICATION

PURSUIT herbicide may be applied by air to crops listed in this label unless otherwise noted.

Uniformly apply with properly calibrated aerial equipment in 5 or more gallons of water per acre. When applied POSTEMERGENCE, the addition of a non-ionic surfactant AND fertilizer solution are required for optimum weed control. Apply a non-ionic surfactant at the rate of 1 quart per 100 gallons of spray solution **OR** a crop oil concentrate at the rate of 1.25 gallons per 100 gallons of spray solution **AND** a liquid fertilizer at the rate of 1.25 gallons per 100 gallons of spray solution. (See instructions under APPLICATION INFORMATION - POSTEMERGENCE).

To avoid injury to sensitive crops from drift, aerial applicators must adhere to the following SPECIAL AERIAL USE DIRECTIONS AND PRECAUTIONS.

- Always use a drift control agent to minimize spray drift.
- Do not spray within 100 feet of other crops (see directions for sensitive crops below).
- If field is less than 100 acres, application by helicopter is recommended.
- Nozzle height above ground must be a maximum of 10 feet.
- Nozzles must be pointed toward the rear of the aircraft. The downward angle of the nozzle should not be greater than 20 degrees:
- To minimize wing-tip vortex roll, nozzles or spray boom must not be located any closer to end of wing or rotor than three-fourths the distance from the center of the aircraft.
- Use a maximum spray pressure of 40 psi.
- DO NOT apply when inversion conditions exist.
- A buffer zone must be established between the area to be sprayed and sensitive crops.
- DO NOT spray when wind velocity is greater than 5 mph. Coarse sprays (larger droplets) are less likely to drift.

Sensitive Crop Precaution (Aerial Use Only)

PURSUIT may cause injury to non-target plants. FOLLOW THE PRECAUTION BELOW WHEN APPLYING PURSUIT BY AERIAL APPLICATION.

- Maintain a downwind buffer zone of at least a half mile from sensitive crops including: vegetables, fallow vegetable beds, sugarbeets, trees or vines which have broken dormancy or are in bloom, or other horticulturally sensitive vegetation.
- For upwind and side borders, maintain a minimum buffer zone of 150 feet from above sensitive crops.

Applicator is responsible for any loss or damage which results from spraying PURSUIT in a manner other than recommended in this label. In addition, applicator must follow all applicable state and local regulations and ordinances in regard to spraying.

APPLICATION INFORMATION

POSTEMERGENCE

PURSUIT is effective in controlling weeds in conservation tillage as well as in **conventional** production systems. **Apply PURSUIT herbicide as an early postemergence treatment when weeds are actively growing and before they exceed a height of 3 inches, unless otherwise indicated.** Delay application until the majority of the weeds are at the recommended growth stage. Application timing should be based on weed size and not crop growth stage. Apply PURSUIT to crops and weeds that are actively growing.

16/62
An adjuvant (either a crop oil concentrate or a surfactant) and a nitrogen-based fertilizer must be added to the spray solution for optimum weed control activity. See the ADJUVANTS section under MIXING INSTRUCTIONS for specific instructions.

When PURSUIT is applied postemergence, absorption will occur through both the roots and foliage. Susceptible weeds stop growing and either die or are not competitive with the crop. PURSUIT not only controls many existing broadleaf and grass weeds when applied postemergence, it also provides control of susceptible weeds that may emerge after application.

For maximum weed control, cultivate 7 - 10 days following a postemergence PURSUIT application. This timely cultivation will enhance residual weed control, especially under dry conditions.

PURSUIT should be applied a minimum of one hour before rainfall or overhead irrigation.

Unusually cool temperatures (50° F or less) reduce photosynthesis and transpiration and thus reduce uptake, translocation, and efficacy of PURSUIT herbicide in weeds. Delaying a PURSUIT application for 48 hours from the time the temperature increases above 50° F, if air temperature has been below 50° F for 10 hours or more hours, will improve weed control and reduce crop response.

NO-TILL/MINIMUM TILLAGE AND DOUBLE CROP SOYBEANS

PURSUIT controls existing weeds and provides residual control of most weeds when applied early postemergence to CLEARFIELD corn or soybeans in no-till or minimum tillage and double crop soybean production systems. The application may be applied either before or after emergence of the crop. (Refer to the WEEDS CONTROLLED POSTEMERGENCE chart for weeds controlled and recommended weed size).

If PURSUIT is applied prior to emergence of the crop, and weeds exceed the recommended size, a contact herbicide should be added to PURSUIT to enhance control. (See instructions for NO-TILL OR REDUCED TILLAGE under the PREEMERGENCE section of this label).

SOIL APPLICATIONS

PURSUIT provides effective weed control in conservation tillage systems designed to meet conservation compliance requirements. PURSUIT can be applied as an early preplant, preplant incorporated, or preemergence treatment in soybeans. It can also be applied in conventional, minimum tillage and no-till production systems. The application method of choice will depend on the anticipated weed spectrum and the preference of the applicator.

Adequate soil moisture is required for optimum activity. Rainfall or overhead irrigation is necessary to move PURSUIT into the weed germination zone. The amount of rainfall or irrigation required following application depends on existing soil moisture, soil texture and organic matter content. Sufficient water to moisten the soil to a depth of 2 inches is normally adequate. If adequate moisture is not received within 7 days after treatment, a cultivation is recommended to control escaped weeds. When adequate moisture is received after dry conditions, PURSUIT will provide residual control of susceptible germinating weeds; activity on established weeds will depend on the weed species and the location of its root system in the soil.

PURSUIT controls weeds by uptake by weed roots and translocation to the growing points where it stops weed growth. Susceptible weeds may emerge, growth will stop and the weeds will either die or are not competitive with the crop.

19/62

SOIL APPLICATIONS WITH LIQUID FERTILIZERS

PURSUIT can be applied to the soil in liquid fertilizers, alone or in combination with PROWL® 3.3 EC, or OUTLOOK® ~~Frontier⁷, Lasso⁴ or Dual³~~ to soybeans or CLEARFIELD corn. Mixtures including trifluralin may be applied to soybeans only. Follow all PURSUIT label recommendations regarding incorporation, timing of application, special instructions and precautions. Apply treatments in 20 or more gallons of liquid fertilizer per acre with ground equipment. Always test the compatibility of PURSUIT with the liquid fertilizer before mixing in the spray tank.

PREEMERGENCE (SURFACE APPLICATIONS)

PURSUIT offers flexibility in that it can be utilized in all production tillage systems. It can be applied prior to planting (up to 45 days prior to planting); at planting in conventional, reduced tillage or no-till production systems; or after planting and before crop emergence.

NO-TILL OR REDUCED TILLAGE

Apply PURSUIT treatments before, during or after planting. To ensure thorough coverage, use a minimum of 20 gallons of water per acre. Use higher gallonage for fields with dense vegetation or heavy crop residues.

For maximum grass control, tank-mix PURSUIT with PROWL 3.3 EC, or OUTLOOK® ~~Frontier, Lasso or Dual~~. To kill existing vegetation, ~~Gramoxone⁶ Gramoxone³ Extra, Starfire⁶ Starfire³, Roundup⁴ Ultra or 2,4-D~~ (early preplant - see 2,4-D label for limitations) may be tank-mixed with PURSUIT alone or in combination with PROWL 3.3 EC, or OUTLOOK ~~Frontier Lasso or Dual~~. ~~Gramoxone Extra, Starfire, Roundup Ultra or 2,4-D~~ should be deleted from the tank-mixture if vegetation is absent at the time of application.

NOTE: Adjust planters to ensure adequate soil ~~seed~~ coverage of seed.

PREPLANT INCORPORATED APPLICATIONS

PURSUIT may be applied following land preparation and should be **thoroughly incorporated** to a depth of 1 to 2 inches. If crops are planted on beds, apply and incorporate after bed formation using PTO-driven equipment or a rolling cultivator. Maintain PURSUIT in the surface 1 to 2 inches of the finished beds. Application may be made up to 45 days prior to planting soybeans.

When PURSUIT is soil applied for control of nutsedge in peanuts incorporate with two passes of the incorporation implement. Make the second pass at an offset angle to the first pass to minimize the potential for streaking.

18/62

FEDERAL CONSERVATION RESERVE PROGRAM AND AGRICULTURAL RESERVE PROGRAM LAND SEEDED TO FORAGE LEGUME SPECIES AND PERENNIAL FORAGE GRASSES.

DIRECTIONS FOR USE

PURSUIT is effective in controlling many annual broadleaf and grass weeds in **CONSERVATION RESERVE PROGRAMS** and **AGRICULTURAL RESERVE PROGRAMS (SET-ASIDE)** land seeded to forage legume or grass crops. A PURSUIT application may result in temporary reduction in growth of legumes and grasses. Plants overcome temporary effects and become well established due to reduced weed competition.

DO NOT feed or graze legumes or grasses following a PURSUIT application. DO NOT cut treated legumes or grasses for hay or forage. DO NOT harvest legume seed for livestock feed. DO NOT use seed from treated legumes for sprouting. Apply only one application of PURSUIT per year.

COVER CROPS:*

LEGUMES: Apply to forage legumes including alfalfa, clovers, crownvetch, birdsfoot trefoil and lespedeza.

GRASSES: PURSUIT may be applied to the following grasses: big bluestem, little bluestem, switchgrass, Russian wildrye, intermediate wheatgrass, crested wheatgrass, western wheatgrass, tall wheatgrass, smooth brome, canarygrass or orchardgrass.

***NOTE:** Cover crops may also be planted into fields previously treated with PURSUIT for weed control in soybeans. In this case, do not make a PURSUIT application to the cover crop until the following spring.

POSTEMERGENCE APPLICATIONS OF PURSUIT TO CRP COVER CROPS.

APPLICATION RATE: Apply PURSUIT at 4 fluid ounces per acre.

APPLICATION TIMING: PURSUIT may be applied postemergence to seedling legumes (with at least 3 fully expanded trifoliolate leaves) or to established legumes. On established legumes, PURSUIT may be applied in the fall or in the spring before weeds exceed the maximum recommended size for control.

DO NOT apply to seeded grasses until they have 4 leaves.

Refer to the WEEDS CONTROLLED under the SOYBEAN section of this label.

CROPS

ALFALFA - DIRECTIONS FOR USE

USE RATE (3 - 6 OUNCES PER ACRE)

Apply PURSUIT at a broadcast rate of 3 to 6 ounces per acre postemergence only.

A maximum of 6 ounces of PURSUIT per year may be applied to alfalfa.

DO NOT apply PURSUIT at more than 4 ounces per acre in North Dakota or Minnesota north of highway #210.

Do not apply more than 4 ounces of product to alfalfa during the last year of the stand.

SEEDLING ALFALFA

PURSUIT must be applied postemergence to seedling alfalfa. Apply PURSUIT when the seedling alfalfa is in the second (2nd) trifoliolate stage or larger and when the majority of the weeds are 1-3 inches. For low growing weeds (such as mustards) apply PURSUIT before the rosette exceeds 3 inches. When PURSUIT is applied to seedling alfalfa, there may be a temporary reduction in growth. Alfalfa soon outgrows any effects of the herbicide.

ESTABLISHED ALFALFA

PURSUIT can be applied to established alfalfa in the fall, in the spring to dormant, or semi-dormant alfalfa (less than 3 inches of re-growth), or between cuttings. Any application should be made before significant alfalfa growth or re-growth (3 inches) to allow PURSUIT to reach the target weeds.

Replanting: If replanting is necessary in a field previously treated with PURSUIT, do not plant alfalfa for 4 months following a PURSUIT application. Refer to the ROTATIONAL CROP GUIDELINE section on this label for plant-back interval of various crops.

WEEDS CONTROLLED

When applied as directed, PURSUIT will control or reduce competition from the weeds listed below. Refer to the MIXING INSTRUCTIONS section for recommendations when weeds are at the maximum recommended growth stage, or are under stress.

NOTE: R = Reduced Competition

Weeds noted with an "R" will be suppressed by PURSUIT. For best results, apply before the weeds exceed the size indicated below.

19/62

BROADLEAF WEEDS CONTROLLED

20/62

Weeds Controlled

PURSUIT APPLICATION RATE

3 Oz/A 4 Oz/A 6 Oz/A

Maximum Weed Size (Inches)

Weeds Controlled	3 Oz/A	4 Oz/A	6 Oz/A
Artichoke; Jerusalem	R	6	8
Beets, wild	4	5	6
Bedstraw, catchweed		3	4
Buckwheat, wild		3	4
Chickweed			
common	R	3	4
mouseear	R	3	3
Cocklebur, common	R	8	8
Cress, hoary		R	R
Dandelion		R	R(5)
Dock,			
broadleaf (seedling)			R(6)
curly (seedling)			R(6)
Dodder			R*
Fiddleneck			R(4)
Filaree			
redstem		R	3
whitstem		R	3
Fleabane, rough		3	3
Flixweed	R	3	4
Goosefoot, nettleleaf	R	3	4
Grounsel, common			R(3)
Henbit		R	3
Jimsonweed		3	4
Knotweed, prostrate		R	3
Kochia (non-ALS resistant)	R	3	3
Lambsquarter,			
common (1-2 leaves)		R	R(2)
Lettuce, miners		3	4
Mallow,			
common		3	3
little		3	3
Marshelder		4	6
Morningglory,			
entireleaf		R	3
ivyleaf		R	3
pitted		R	3
smallflower	R	3	4
tall		R	3
Mustards			
tumble	3	3	4
wild	3	3	4
black	3	3	4
Nettle, burning		3	4

BROADLEAF WEEDS CONTROLLED - Continued

21/62

Weeds Controlled	PURSUIT APPLICATION RATE		
	3 Oz/A	4 Oz/A	6 Oz/A
Maximum Weed Size (Inches)			
Nightshade			
black	3	3	4
Eastern black	3	3	4
hairy	3	3	4
Oxtongue, bristly			R(3)
Pennycress, field	3	3	4
Pepperweed,			
field	3	3	4
Virginia	R	3	3
Pigweed			
redroot	4	6	8
smooth	4	6	8
spiny		6	8
Radish, wild		R	4
Ragweed			
common		2	3
giant		3	3
Redmaids		3	4
Rocket			
London	3	4	6
yellow	R	3	4
Rockpurslane, desert			3
Shepherdspurse	3	3	4
Smartweed			
ladysthumb	R	3	4
Pennsylvania	R	3	4
swamp (seedling)		3	4
Spurge			
prostrate		R	3
spotted		R	3
petty		3	4
Spurry, corn		3	3
Sunflower, common	R	4	6
Swinecress		3	3
Tansymustard			
green	3	3	4
pinnate	3	3	4
Thistle, Russian	R	3	3
Velvetleaf	R	3	4
Wartcress, creeping		2	3
Watercress		3	3
Willowweed, panicle		3	3

*For best results in suppressing dodder (*cuscuta* spp), apply PURSUIT with crop oil concentrate or methylated seed oil after dodder has emerged but prior to or soon after attachment.

GRASSES AND SEDGES CONTROLLED

22/62

Weeds Controlled*

PURSUIT APPLICATION RATE

4 Oz/A 6 Oz/A

Maximum Weed Size (Inches)

Barnyardgrass	R	3
Bluegrass, annual		R(3)
Canarygrass, littleseed	R	R(3)
Cereals, volunteer		
barley	R	R(4)
oats	R	R(4)
wheat	R	R(4)
Crabgrass,		
large	R	3
smooth	R	3
Cupgrass, woolly**	3	3
Foxtail,		
giant	6	6
green	3	4 ²
yellow	3	3
Johnsongrass,		
seedling	8	8
rhizome	R	R(6-12)
Millet, wild proso	R	3
Nutsedge,		
yellow	R	R(6)
purple	R	R(6)
Oats, wild	R	R(4)
Rice, red	3	4
Shattercane	8	10
Signalgrass, broadleaf	R	8
Quackgrass ***		R(7)

* PURSUIT is active against many grass species. However, when heavy grass pressure is anticipated, PURSUIT should be used in a sequential application with a registered postemergence grass herbicide such as Poast Plus for optimum control.

** PURSUIT controls emerged woolly cupgrass only.

*** Quackgrass will be suppressed only when actively growing and before it exceeds 7 inches in height.

TANK MIX COMBINATIONS WITH OTHER HERBICIDES

To control weeds not listed on the PURSUIT label, herbicides such as ~~Bueth~~¹⁸Buctril⁵, 2,4-DB, Poast[®], or Poast Plus[®], Prism⁶ or ~~Select~~⁹Select⁶ may be tankmixed with PURSUIT. When PURSUIT is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded.

23/62

APPLICATION INFORMATION

PURSUIT is effective in controlling a broad spectrum of broadleaf and grass weeds. Alfalfa is tolerant to postemergence applications of PURSUIT after the second trifoliolate leaf has expanded. Minor height reduction or slight leaf yellowing may occur soon after application. The effects are temporary and the alfalfa will soon recover.

Apply PURSUIT as an early postemergence treatment when the weeds are actively growing. Weeds are generally easier to control before they exceed 3 inches in height. Weeds under stress are less susceptible to control in cold or drought stress conditions.

If applied to alfalfa under cool conditions (40° F or less), temporary stunting and yellowing of the alfalfa may occur.

STAND ESTABLISHMENT

Apply PURSUIT after the alfalfa has 2 fully expanded trifoliolate leaves. Weeds must not exceed the size listed in the WEEDS CONTROLLED tables. PURSUIT may be applied to summer, fall or spring seeded alfalfa.

Inter-seeded Oats

Oats inter-seeded with alfalfa will reduce soil erosion and allow the alfalfa to establish. Oats, however can compete with the alfalfa. An application of PURSUIT will kill or significantly reduce the growth of the oats and allow the alfalfa to establish with minimal erosion or competition from the oats. Apply PURSUIT to the oats when the oats have 3-4 leaves.

ESTABLISHED ALFALFA - DORMANT

PURSUIT may be applied to dormant alfalfa in the fall following the last cutting. PURSUIT may also be applied in the spring to dormant alfalfa, or as alfalfa breaks dormancy. Apply spring treatments prior to excessive alfalfa growth (less than 3 inches), to reduce spray interference.

ESTABLISHED ALFALFA - GROWING

For weed control during the season, apply PURSUIT following alfalfa cutting. Remove the hay from the field and apply PURSUIT prior to excessive alfalfa regrowth.

Perennial Grass Suppression

If perennial grasses (such as orchardgrass, fescues, bromes or timothy) are present in an alfalfa stand, PURSUIT will reduce the growth and competitive effect of the grass.

24/62

CLEARFIELD CORN - DIRECTIONS FOR USE

PURSUIT HERBICIDE USE AREA

~~(Not for use in New York)~~(Not for use in California)

**USE RATE
(4 OUNCES PER ACRE)**

Apply PURSUIT® at a broadcast rate of 4 ounces per acre (1/4 pint) for all methods of application: early preplant, preplant incorporated, preemergence, and postemergence (including minimum and no-till). At this broadcast rate, one gallon of PURSUIT will treat 32 acres of CLEARFIELD corn. (See instructions on page 24 for applications in North Dakota and Minnesota north of highway #210).

NOTE: Only one application of PURSUIT may be made during the season.

WEEDS CONTROLLED

When applied as directed, PURSUIT will control or reduce competition from the weeds listed below. Refer to the MIXING INSTRUCTIONS section for recommendations when weeds are at the maximum recommended growth stage, or are under stress.

NOTE: C = Control, R = Reduced Competition

(The number under Maximum Leaf Stage indicates the MAXIMUM number of leaves at which weeds should be sprayed postemergence).

BROADLEAF WEEDS

25/62

Weeds Controlled	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Alligator weed		4	1-3
Anoda, spurred	C	2	1-2
Artichoke, Jerusalem		8	6-10
Buffalobur	C*	R	1-3
Bristly starbur		2	1-2
Carpetweed	C		
Cocklebur, common	R	8	1-8
Galinsoga	C		
Jimsonweed	C*	4	1-3
Kochia (non-ALS resistant)	C	4	1-3
Lambsquarters, common	C*	R	1-2
Mallow, Venice	R		
Marshelder	C	4	1-3
Morningglory			
entireleaf	R	2	1-2
ivyleaf	R	2	1-2
pitted	R	2	1-2
smallflower	C	4	1-3
tall	R	2	1-2
Mustard sp.	C	4	1-3
Nightshade			
black	C	4	1-3
Eastern black	C	4	1-3
hairy	C	4	1-3
Pigweed			
redroot	C	8	1-8
smooth	C	8	1-8
spiny	C	8	1-8
Poinsettia, wild	C		
Puncturevine	C		
Purslane, common	C		
Pusley, Florida	C		
Sida, prickly	C*		
Ragweed,			
common	R	4	1-3
giant	R	4	1-3
Sage, barnyard		R	1-3
Smartweed			
ladysthumb	C	4	1-3
Pennsylvania	C	4	1-3
Spurge			
prostrate	C	4	1-3
spotted	C	4	1-3
Sunflower	C*	4	1-3
Velvetleaf	C*	4	1-3
Thistle, Canada		R	1-3

* When PURSUIT is soil applied, these weeds are more consistently controlled by preplant incorporated treatments.

DO NOT count cotyledon leaves when determining weed stage of growth.

26/62

GRASS WEEDS*

Weeds Controlled**	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Barnyardgrass	R	3	1-3
Crabgrass,			
large	R	3	1-3
smooth	R	3	1-3
Cupgrass, woolly		3	1-3
Foxtail,			
giant	C	6	1-6
green	C	3	1-3
yellow	C	3	1-3
Goosegrass	R		
Johnsongrass,			
seedling	C	6	1-8
rhizome		R	6-12
Millet, wild proso	R	R	1-3
Panicum,			
fall	R		
Texas	R		
Red rice		3	1-3
Sandbur, field	R	R	≤1
Shattercane	R	6	1-8
Signalgrass, broadleaf	R	4	1-8
Sorghum alnum	R	6	1-3

SEDGES

Nutsedge			
purple	R	R	1-3
yellow	R	R	1-3

* Preplant incorporated treatments of PURSUIT are more consistent for grass control.

** PURSUIT controls many grass species. However, when heavy grass pressure is anticipated, a soil applied grass herbicide underlay (such as PROWL, Dual, Eradicane⁶, or OUTLOOK Frontier, Harness⁴, Lasso, or Surpass⁶) is recommended for optimum control. DO NOT incorporate PROWL 3.3 EC herbicide; apply preemergence or early postemergence only. PURSUIT may also be used in sequential programs with registered burndown herbicides and/or soil applied atrazine-containing products.

TANK-MIXTURE HERBICIDE COMBINATIONS WITH PURSUIT (Postemergence)

27/62

Accent ^Δ	Dual ³
Atrazine ^{*,**}	FrontierOUTLOOK
Banvel ^{*,***}	Lasso EC, Lasso MT
Basagran ^{*,7,*}	Marksman ^{*,*}
Buctril ^{5*,**}	PROWL 3.3 EC
Clarity ^{*,***}	

DO NOT use crop oil concentrates as adjuvants in PURSUIT combinations with Buctril.

PURSUIT is active against many broadleaf and grass species. However, for long term weed management, alternate mode of action herbicides are recommended with PURSUIT. The application of a soil applied grass herbicide underlay will control grass weeds not on the PURSUIT herbicide label and enhance the control of certain broadleaf weeds such as common lambsquarters.

When PURSUIT is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT cannot be mixed with any product containing a label prohibiting such mixtures.

- ^Δ If Accent is used in combination with PURSUIT on Pioneer imidazolinone-resistant (IR) corn, any registered soil insecticide applications may be used.
If PURSUIT plus Accent tank-mixes are used on imidazolinone -tolerant (IT) hybrids, DO NOT use COUNTER[®] 15G. Other registered organophosphate insecticides such as COUNTER[®] CR[®] (banded applications only) or THIMET[®] or other registered carbamate or pyrethroid insecticides may be used when PURSUIT plus Accent tankmixes are applied to IT corn hybrids.
- ^{*} In some cases the grass activity of PURSUIT will be reduced when used in combination with atrazine, Buctril, Banvel, Basagran, Clarity, or Marksman.
- ^{**} Some corn leaf burn may result with Buctril or atrazine postemergence combinations with PURSUIT
- ^{***} Applications of Banvel or Clarity herbicide to corn during periods of rapid growth may result in temporary leaning.

25/02

**APPLICATIONS TO CLEARFIELD CORN IN NORTH DAKOTA AND MINNESOTA
(north of highway #210).**

APPLICATION RATE: Apply PURSUIT at 3 ounces per acre postemergence only.


Weeds Controlled	POSTEMERGENCE	
	Maximum Leaf Stage	Size (inches)
Kochia (non-ALS resistant)	4	1-3
Mustard, species	4	1-3
Nightshade, black	4	1-3
Eastern black	4	1-3
hairy	4	1-3
Pigweed, redroot	4	1-4
Wild oats*	3	1-4

*PURSUIT will reduce competition from wild oats.

NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE, CRANBERRY, PINTO, LIMA, AND SMALL WHITE TYPE DRY BEANS, LENTILS, WHITE LUPINS, CHICKPEAS (GARBANZO BEANS), DRY EDIBLE PEAS, ENGLISH AND SOUTHERN PEAS

29/62

DIRECTIONS FOR USE in the states east of and including: North Dakota, South Dakota, Wyoming, Colorado, and New Mexico (except the states east of and including: Vermont, Massachusetts, and Connecticut). Refer to map for geographical use area.



Use only nonionic surfactants as a spray additive for postemergence applications. **DO NOT** use crop oils, methylated seed oils, or petroleum oils, or ~~Dash~~.

DO NOT make more than one application of PURSUIT per year.

Allow at least 30 days between application and harvest of ^{postemergent} lima beans, English peas, and Southern peas. Allow at least 60 days between application and harvest of dry edible peas, lentils, chickpeas, and other dry bean or pea types listed on this label.

DO NOT APPLY PURSUIT POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRIFOLIATE LEAF OR PEAS ARE AT LEAST THREE INCHES IN HEIGHT OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT. DO NOT APPLY PURSUIT POSTEMERGENCE TO LIMA BEANS, LENTILS, WHITE LUPINS, OR CHICKPEAS.

DO NOT apply to Domino variety black turtle beans.

DO NOT apply this product through any type of irrigation system.

Pinto varieties UI-111 and Olathe are more sensitive to PURSUIT than other varieties.

APPLICATION INSTRUCTIONS

NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE, CRANBERRY, PINTO, AND SMALL WHITE DRY BEANS, DRY EDIBLE PEAS, ENGLISH AND SOUTHERN PEAS

In Michigan or the Delaware, Maryland, and Virginia (Delmarva) peninsula: DO NOT apply more than 2 oz of PURSUIT to sands or loamy sand soils.

In North Dakota or north of Highway 210 in Minnesota: DO NOT apply more than 2 oz of PURSUIT.

Preplant Incorporated Applications. Apply PURSUIT at the broadcast rate of up to 3 ounces per acre to dry beans (navy, great northern, red kidney, black turtle, cranberry, pinto and small white type dry beans), dry edible peas, and English peas, or up to 4 ounces per acre for southern peas only, within 1 week before planting. Applied preplant incorporated, PURSUIT may be tank-mixed with a registered grass herbicide.

Preemergence Applications. Apply PURSUIT at the broadcast rate of up to 3 ounces per acre to dry beans, dry edible peas and English peas, or up to 4 ounces per acre for southern peas only, immediately after, or up to 3 days after planting. PURSUIT may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

Early Postemergence Applications. Apply PURSUIT at the broadcast rate of up to 3 ounces per acre to dry beans, dry edible peas, and English peas, or up to 4 ounces per acre for southern peas only. Apply to dry beans with at least one fully expanded trifoliolate leaf. Apply to dry edible peas, English peas, and southern peas at least 3 inches in height but prior to 5 nodes and before flowering. The use of trifluralin prior to PURSUIT application may increase the likelihood and severity of crop injury. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

Basagran¹ may be tank-mixed with PURSUIT to control weeds not listed on the PURSUIT label. Addition of Basagran may also cause antagonism, thereby reducing control of grass weeds. Nitrogen-based fertilizer may be included as a spray additive **ONLY** when PURSUIT is tank-mixed with Basagran. Refer to the Basagran label for proper application rates and restrictions. Always use in accordance with the more restrictive label restrictions and precautions.

Success **LIMA BEANS, CHICKPEAS (GARBANZOS), LENTILS, ~~AND~~ WHITE LUPINS** *and*

DO NOT apply PURSUIT to white lupins grown on sand or loamy sand soils.

In Michigan or the Delaware, Maryland, and Virginia (Delmarva) peninsula: **DO NOT** apply more than 2 oz of PURSUIT to sands or loamy sand soils.

In North Dakota or north of Highway 210 in Minnesota: **DO NOT** apply more than 2 oz of PURSUIT.

Preplant Incorporated Applications. Apply PURSUIT at the broadcast rate of up to 3 ounces per acre within 1 week before planting. Applied preplant incorporated, PURSUIT may be tank-mixed with a registered grass herbicide.

Preemergence Applications. Apply PURSUIT at the broadcast rate of up to 3 ounces per acre immediately after or up to 3 days after planting. PURSUIT may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

WEEDS CONTROLLED

PURSUIT applied at the broadcast rate of 2 ounces per acre preplant incorporated, preemergence, or early postemergence will control:

- Mustard, wild
- Nightshade, black*
- Nightshade, eastern black*
- (*suppression only)

31/62

PURSUIT applied at the broadcast rate of 3 ounces per acre preplant incorporated, preemergence, or early postemergence will control:

Mustard, wild	Nightshade, hairy
Nightshade, black	Pigweed, redroot
Nightshade, Eastern black	

Postemergence applications of 3 ounces per acre must be made to weeds less than 2 inches tall for best results.

When applied as directed at the broadcast rate of 4 ounces per acre (for southern peas only), PURSUIT will control or reduce competition from the weeds listed below:

NOTE: C = Control, R = Reduced Competition

The number under Maximum Leaf Stage indicates the MAXIMUM number of leaves at which weeds should be sprayed postemergence.

BROADLEAF WEEDS

WEEDS CONTROLLED	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Anoda, spurred	C	2	1 - 2
Artichoke, Jerusalem		8	6 - 10
Buffalobur	C*		
Bristly starbur		2	1 - 2
Carpetweed	C		
Cocklebur, common	C*	8	1 - 8
Galinsoga	C		
Jimsonweed	C**	4	1 - 3
Kochia (non-ALS resistant)	C	4	1 - 3
Lambsquarters	C**	R	1 - 2
Mallow, Venice	R		
Morningglory			
entireleaf	R	2	1 - 2
ivyleaf	R	2	1 - 2
pitted	R	2	1 - 2
smallflower	C	4	1 - 3
tail	R	2	1 - 2
Mustard sp.	C	4	1 - 3
Nightshade,			
black	C	4	1 - 3
Eastern black	C	4	1 - 3
hairy	C	4	1 - 3
Pigweed,			
redroot	C	4	1 - 4
smooth	C	4	1 - 4
spiny	C	4	1 - 4
Poinsettia, wild	C		
Puncturevine	C		
Purslane, common	C		
Pusley, Florida	C		
Sida, prickly	C**		
Ragweed,			
common	R	4	1 - 3
giant	R	4	1 - 3
Sage, barnyard		R	1 - 3
Smartweed,			
ladysthumb	C	4	1 - 3
Pennsylvania	C	4	1 - 3
Spurge,			
prostrate	C	4	1 - 3
spotted	C	4	1 - 3
Sunflower, common	C**	4	1 - 3
Thistle, Canada		R	1 - 3
Velvetleaf	C**	4	1 - 3

*Use soil applications for light to moderate infestations only. Must be preplant incorporated for best results.

**When soil applied, common lambsquarters, jimsonweed, prickly sida, velvetleaf and common sunflower are more consistently controlled by preplant incorporated treatments.

GRASS WEEDS*

WEEDS CONTROLLED	SOIL APPLIED	POSTEMERGENCE	
		Maximum Leaf Stage	Size (inches)
Barnyardgrass	R	3	1 - 3
Crabgrass,			
large	R	3	1 - 3
smooth	R	3	1 - 3
Cupgrass, woolly		3**	1 - 3
Foxtail,			
giant	C	6	1 - 6
green	C	3	1 - 3
robust purple	C	3	1 - 3
robust white	C	3	1 - 3
yellow	C	3	1 - 3
Goosegrass	R		
Johnsongrass,			
seedling	C	6	1 - 8
rhizome		R	1 - 8
Panicum,			
fall	R		
Texas	R		
Red rice		3	1 - 3
Shattercane	R	6	1 - 8
Signalgrass, broadleaf	R	4	1 - 8

SEDGES

Nutsedge,			
purple	R	R	1 - 3
yellow	R	R	1 - 3

DO NOT count cotyledon leaves when determining weed stage of growth.

*When soil applied to grasses, more consistent control can be obtained from preplant incorporated treatments.

**PURSUIT controls emerged woolly cupgrass only.

Refer to the PRECAUTIONS section for additional instructions.

Dry

RED KIDNEY BEANS

DIRECTIONS FOR USE in the state of California.

Do Not Apply By Aerial Application.

APPLICATION RATE AND TIMING

Postemergence Applications:

Apply PURSUIT at the rate of 3 ounces per acre. A non-ionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

Apply PURSUIT herbicide when weeds are actively growing and red kidney beans have at least 1 fully expanded trifoliate leaf. DO NOT apply PURSUIT postemergence when the crop and weeds have been subjected to stress conditions such as temperature or moisture extremes.

For maximum weed control, cultivate 7-10 days following a postemergence PURSUIT application. This timely cultivation will enhance residual weed control, especially under dry conditions.

DO NOT APPLY PURSUIT POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

WEEDS CONTROLLED

When applied as directed, PURSUIT will control or reduce competition from the weeds listed below. Refer to the MIXING INSTRUCTIONS section for recommendations when weeds are at the maximum recommended growth stage, or are under stress.

(The number under Maximum Leaf Stage indicates the MAXIMUM number of leaves at which weeds should be sprayed postemergence).

WEEDS

Weeds Controlled	POSTEMERGENCE	
	Maximum Leaf Stage	Size (inches)
Kochia (non-ALS resistant)	4	1-3
Mustard, wild	4	1-3
Nightshade,		
black	4	1-3
Eastern black	4	1-3
hairy	4	1-2
Pigweed,		
redroot	4	1-3

Allow at least 60 days between application and harvest.
Do not make more than one application of Pursuit per year.

Refer to the PRECAUTIONS section for additional instructions.

35/62

SNAP BEANS

DIRECTIONS FOR USE in the states of Alabama, Florida, Georgia, Illinois, Minnesota, Michigan, New Jersey, and Wisconsin.

Do Not Apply By Aerial Application.

Do Not Apply PURSUIT After July 31st (June 20th in New Jersey).

APPLICATION INSTRUCTIONS

Preplant Incorporated Applications. Apply PURSUIT at 1.5 oz per acre within 1 week of planting. Applied preplant incorporated, PURSUIT may be tank-mixed with a registered grass herbicide.

Preemergence Applications. Apply PURSUIT at the broadcast rate of 1.5 oz per acre immediately after, or up to 1 day after planting. PURSUIT may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

WEEDS SUPPRESSED

PURSUIT applied at the broadcast rate of 1.5 oz/A preplant incorporated or preemergence will suppress or reduce competition of the following weeds:

- Common Purslane
- Eastern Black Nightshade
- Redroot Pigweed
- Wild Mustard

Allow at least 30 days between application and harvest.

Do not make more than one application of Pursuit per year.

Refer to the PRECAUTIONS section for additional instructions.

SNAP BEANS

36/62

DIRECTIONS FOR USE in the states of Arkansas, Missouri, Oklahoma, Texas (counties of Bailey, Castro, Lamb and Parmer only) and New Mexico (counties of Curry and Roosevelt only).

Do Not Apply By Aerial Application.

Do Not Apply PURSUIT After July 31.

APPLICATION INSTRUCTIONS

Postemergence Applications. Apply PURSUIT at 1.5 oz per acre in a tank-mix combination with Basagran. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

Refer to the Basagran label for proper application rates and restrictions.

DO NOT APPLY PURSUIT POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

WEEDS SUPPRESSED

PURSUIT applied at the broadcast rate of 1.5 oz/A postemergence will suppress or reduce competition of the following weeds:

Eastern Black Nightshade
Redroot Pigweed

✓ **Allow at least 30 days between application and harvest.**

Do not make more than one application of Pursuit per year.

Refer to the PRECAUTIONS section for additional instructions.

succulent

SUCCULENT PEAS, DRY EDIBLE PEAS, LENTILS, CHICKPEAS, AND LIMA BEANS

37/62

DIRECTIONS FOR USE in the states of Idaho, Montana, Nevada, Oregon, Utah, and Washington.

APPLICATION RATE AND TIMING

Preplant Applications for No-Till and Minimum Tillage Systems Only:

Apply PURSUIT at a broadcast rate of 3 ounces per acre within 30 days before planting. If incorporated, DO NOT incorporate deeper than 3 inches.

In no-till and minimum tillages systems, PURSUIT may be applied in the fall prior to spring planting. Rainfall is required for incorporation and activation. Unpredictable weed control can be expected since factors such as length of time between application and planting as well as uncontrollable weather factors will determine herbicide activity and longevity. Apply PURSUIT in the fall when soil temperature at the 4-inch depth is less than 55°F and before the ground is frozen.

Preplant Incorporated Applications:

Apply PURSUIT at the broadcast rate of 3 ounces per acre within 1 week before planting. DO NOT incorporate deeper than 3 inches.

Preemergence Applications:

Apply PURSUIT at the broadcast rate of 3 ounces per acre after planting, but prior to crop emergence.

PURSUIT may be tank-mixed with Sencor⁹-Sencor⁷ DF or Lexone² DF to assist in the control of lambsquarters or mayweed-chamomile (dogfennel). Refer to the Sencor or Lexone label for proper application rates and restrictions.

Postemergence Applications (Dry Edible Peas Only):

Apply PURSUIT at 2 ounces per acre. A nonionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and should be used at a rate of 2 pints per 100 gallons of spray mixture.

Basagran may be tank-mixed with PURSUIT to control weeds not listed on the PURSUIT label. Addition of Basagran may also cause antagonism, thereby reducing control of grass weeds. Nitrogen-based fertilizer may be included as a spray additive only when PURSUIT is tank mixed with Basagran. Use liquid fertilizer at 1.25 to 2.5 gal. per 100 gal. of spray solution or ammonium sulfate at the rate of 12-15 lbs/100 gal. of spray solution.

DO NOT APPLY PURSUIT POSTEMERGENCE BEFORE CROP HAS AT LEAST ONE TRIFOLIATE LEAF OR PEAS ARE AT LEAST THREE INCHES IN HEIGHT OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT. DO NOT APPLY PURSUIT POSTEMERGENCE TO LIMA BEANS, LENTILS, OR CHICKPEAS.

38/62

WEEDS CONTROLLED

PURSUIT applied PPI and/or Preemergence at 3 oz/A will control:

Weeds Controlled	Preplant Incorporated	Preemergence
Buckwheat, wild	C	C
Kochia (non-ALS resistant)	C	C
Lambsquarters, common	C	
Mustard, wild	C	C
Nightshade,		
Black	C	C
Eastern black	C	C
Hairy	C	C
Pigweed, redroot	C	C
Shepherdspurse	C	C
Thistle, Russian	C	C

NOTE: C = Control

PURSUIT applied postemergence at the broadcast rate of 2 ounces will control:

- Wild mustard
- Black nightshade*
- Eastern black nightshade*
- Hairy nightshade*

*Suppression only

Allow at least 30 days between application and harvest for succulent peas and lima beans.

Allow at least 60 days between application and harvest for dry edible peas, chickpeas and lentils.

Do not make more than one application of Pursuit per year.

Refer to the PRECAUTIONS section for additional instructions.

Dry or succulent - ? ask registrant

39/62

CHICKPEAS

DIRECTIONS FOR USE in the states of Arizona and California.

APPLICATION RATE AND TIMING

Preplant Incorporated Applications. Apply PURSUIT at the broadcast rate of up to 3 ounces per acre within 1 week before planting. Applied preplant incorporated, PURSUIT may be tank-mixed with a registered grass herbicide.

Preemergence Applications. Apply PURSUIT at the broadcast rate of up to 3 ounces per acre immediately after or up to 3 days after planting. PURSUIT may be applied in a tank-mix with a registered grass herbicide or applied preemergence following a preplant incorporated application of a registered grass herbicide.

WEEDS CONTROLLED

Weeds Controlled	Preplant Incorporated	Preemergence
Buckwheat, wild	C	C
Kochia (non-ALS resistant)	C	C
Lambsquarters, common	C	
Mustard, wild	C	C
Nightshade,		
Black	C	C
Eastern black	C	C
Hairy	C	C
Pigweed, redroot	C	C
Shepherdspurse	C	C
Thistle, Russian	C	C

NOTE: C = Control

Allow at least 30 days between application and harvest.
Do not make more than one application of Pursuit per year.
Refer to the PRECAUTIONS section for additional instructions.

*for succulent only
60 days for dry chickpeas
chick peas usually harvested at full maturity (dry)*

PEANUTS - DIRECTIONS FOR USE

**PURSUIT HERBICIDE USE RATE
(4 OUNCES PER ACRE)
(Not for use in California)**

Apply PURSUIT at a broadcast rate of 4 ounces per acre (1/4 pint) for all methods of application (except sequential - see below): preplant incorporated, preemergence, ground-cracking and postemergence. At this broadcast rate, one gallon of PURSUIT will treat 32 acres of peanuts.

PURSUIT may also be applied in a sequential application. Apply 2 ounces in a soil application (preplant incorporated or preemergence) followed by 2 ounces applied at ground-crack or postemergence.

NOTE: DO NOT apply more than 4 ounces of PURSUIT during the growing season.

NOTE: In Arizona for use only in Yuma and La Paz counties.

WEEDS CONTROLLED

When applied as directed, PURSUIT will control or reduce competition from the weeds listed below. Refer to the MIXING INSTRUCTIONS section for recommendations when weeds are at the maximum recommended growth stage, or are under stress.

NOTE: C = Control, R = Reduced Competition

(The number under Maximum Leaf Stage indicates the MAXIMUM number of leaves at which weeds should be sprayed postemergence).

BROADLEAF WEEDS

41/62

Weeds Controlled	SOIL APPLIED	AT-CRACK	POSTEMERGENCE	
			Maximum Leaf Stage	Size (inches)
Alligator weed		C	4	1-3
Anoda, spurred	C	C	2	1-2
Buffalobur	C*	C	R	1-3
Bristly starbur			2	1-2
Carpetweed	C	C		
Cocklebur, common	R	C	8	1-8
Devilsclaw	C	C		
Galinsoga	C	C		
Jimsonweed	C*	C	4	1-3
Lambsquarters, common	C*	C	R	1-2
Morningglory				
entireleaf	R	C	2	1-2
ivyleaf	R	C	2	1-2
pitted	R	C	2	1-2
smallflower	C	C	4	1-3
tall	R	C	2	1-2
Mustard sp.	C	C	4	1-3
Nightshade				
black	C	C	4	1-3
Eastern black	C	C	4	1-3
hairy	C	C	4	1-3
Pigweed				
redroot	C	C	8	1-8
smooth	C	C	8	1-8
spiny	C	C	8	1-8
Poinsettia, wild	C	C		
Puncturevine	C	C		
Purslane, common	C	C		
Pusley, Florida	C	C		
Ragweed,				
common	R	R	4	1-3
giant	R	R	4	1-3
Sida, prickly (teaweed)	C*	C		
Smartweed				
ladysthumb	C	C	4	1-3
Pennsylvania	C	C	4	1-3
Spurge				
prostrate	C	C	4	1-3
spotted	C	C	4	1-3
toothed	C	C		
Sunflower	C*	C	4	1-3
Velvetleaf	C*	C	4	1-3

* When PURSUIT is soil applied, these weeds are more consistently controlled by preplant incorporated treatments.

42/62

GRASS WEEDS

Weeds Controlled	SOIL APPLIED*	AT-CRACK	POSTEMERGENCE	
			Maximum Leaf Stage	Size (inches)
Barnyardgrass	R	R	3	1-3
Crabgrass,				
large	R	C	3	1-3
smooth	R	C	3	1-3
Cupgrass, woolly			3	1-3
Foxtail,				
giant	C	C	6	1-6
green	C	C	3	1-3
yellow	C	C	3	1-3
Goosegrass	R	R		
Johnsongrass,				
seedling	C	C	6	1-8
rhizome			R	6-12
Panicum,				
fall	R			
Texas	R			
Red rice			3	1-3
Shattercane	R	R	6	1-8
Signalgrass, broadleaf	R	C	4	1-6

SEDGES

Nutsedge				
purple	C	C	3	1-3
yellow	C	C	3	1-3

* When PURSUIT is soil applied to grasses, more consistent control can be obtained from preplant incorporated treatments.

Do not count cotyledon leaves when determining weed stage of growth.

AT CRACK APPLICATION refers to the time when the soil cracks due to the emerging peanut seedling. This generally occurs from 10 to 14 days following planting. At this time weeds have generally not germinated, or are in the seedling stage. If weeds have more than 2 true leaves, refer to the **POSTEMERGENCE** weed control column for weeds controlled.

In West Texas and New Mexico, wait until late cracking (most of the peanuts have emerged) before applying PURSUIT.

PURSUIT is active against many broadleaf and grass species. However, when heavy grass or common lambsquarters pressure is anticipated, PURSUIT should be used in combination with a registered soil-applied grass herbicide (See **HERBICIDE COMBINATIONS** section).

WEEDS CONTROLLED BY SEQUENTIAL APPLICATIONS OF PURSUIT

The sequential (split) application of PURSUIT consists of an application of 2 ounces of product soil applied (either preplant incorporated or preemergence) followed by 2 ounces applied either at ground-crack or postemergence.

When applied as a sequential treatment, PURSUIT will control the weeds listed under the "SOIL APPLIED" and "AT-CRACK" applications in the BROADLEAF WEEDS and GRASS WEEDS tables (in the peanut section of the label). It enhances the control of yellow and purple nutsedge. Apply the second application before the nutsedge exceeds 3 leaves.

HERBICIDE COMBINATIONS

44/62

GRASS WEEDS

When applied as directed, PURSUIT preplant incorporated or preemergence combination treatments with PROWL 3.3 EC[®] herbicide, trifluralin, Lasso¹, Dual, Balan¹ Sonalan¹ or Vernam³ will control the weeds listed in following table, in addition to those controlled by PURSUIT alone.

	PROWL 3.3 EC ^a	trifluralin ^b	Lasso	Dual	Balan ^b	Sonalan ^b	Vernam ^b
GRASSES							
Barnyardgrass	X	X	X	X	X	X	X
Crabgrass, smooth	X	X	X	X	X	X	X
Crabgrass, large	X	X	X	X	X	X	X
Crowfootgrass	X	X			X		
Goosegrass	X	X	X	X	X	X	X
Panicum, fall	X	X	X	X	X	X	X
Panicum, Texas	X	X			X	X	
Sandbur, field	X	X			X	X	
Signalgrass, broadleaf	X ^b	X	X	X	X	X	
Witchgrass	X	X	X	X		X	

^a Preplant incorporated tank-mixture applications of PURSUIT plus PROWL 3.3 EC will suppress the growth of itchgrass, and rhizome johnsongrass.

^b Preplant incorporated treatments only.

A selective postemergence grass herbicide such as Poast Plus, Bugle¹⁰-Bugle⁵ or Whip¹⁰-Whip⁵ may be mixed with PURSUIT to control grasses not controlled by PURSUIT. In some cases the activity of the grass herbicide may be reduced when mixed with PURSUIT. The reduction in activity may be overcome by delaying the application of the postemergence grass herbicide 7 days following the application of PURSUIT. If the postemergence grass herbicide is applied first, wait 3 days before applying PURSUIT. Refer to the respective grass herbicide label for recommended application rate, weed size and restrictions.

45/62

BROADLEAF WEEDS

Broadleaf herbicides that can be tank-mixed with PURSUIT include Basagran and Ultra Blazer, Starfire and 2,4-DB. Certain herbicides should not be applied with PURSUIT (see PRECAUTIONS section for restrictions).

For the control of sicklepod, morningglories, prickly sida and common ragweed, add 2,4-DB to the PURSUIT spray mixture. For the control of Florida beggarweed, add Starfire to the spray mixture. Refer to the 2,4-DB or Starfire label for specific directions for use, application rates and restrictions.

PURSUIT may also be applied postemergence in tank-mixture with Bravo^{1,2,3}, Bravo S^{1,2,3}, Orthene⁴-Orthene⁶ or Soluber^{1,4}-Solubor⁸.

SOYBEANS - DIRECTIONS FOR USE

46/62

USE RATE (4 OUNCES PER ACRE) (Not for use in California)

Apply PURSUIT at a broadcast rate of 4 ounces per acre (1/4 pint) for all methods of application: early preplant, preplant incorporated, preemergence, and postemergence (including minimum and no-till). At this broadcast rate, one gallon of PURSUIT will treat 32 acres of soybeans. (See instructions under section APPLICATIONS TO SOYBEANS IN NORTH DAKOTA AND MINNESOTA for applications in North Dakota and Minnesota north of highway #210).

NOTE: Only one application of PURSUIT may be made during the season.

WEEDS CONTROLLED

When applied as directed, PURSUIT will control or reduce competition from the weeds listed below. Refer to the MIXING INSTRUCTIONS section for recommendations when weeds are at the maximum recommended growth stage, or are under stress.

NOTE: C = Control, R = Reduced Competition

(The number under Maximum Leaf Stage indicates the MAXIMUM number of leaves at which weeds should be sprayed postemergence).

Do not count cotyledon leaves when determining weed stage of growth.

**BROADLEAF WEEDS
SOIL APPLIED**

47/62

Weeds Controlled (inches)		POSTEMERGENCE	
		Maximum Leaf Stage	Size
Alligator weed		4	1-3
Anoda, spurred	C	2	1-2
Artichoke, Jerusalem		8	6-
10			
Buffalobur	C*	R	1-3
Bristly starbur		2	1-2
Carpetweed	C		
Cocklebur, common	R	8	1-8
Galinsoga	C		
Jimsonweed	C*	4	1-3
Kochia (non-ALS resistant)	C	4	1-3
Lambsquarters, common	C*	R	1-2
Mallow, Venice	R		
Marshelder	C	4	1-3
Morningglory			
entireleaf	R	2	1-2
ivyleaf	R	2	1-2
pitted	R	2	1-2
smallflower	C	4	1-3
tall	R	2	1-2
Mustard sp.	C	4	1-3
Nightshade			
black	C	4	1-3
Eastern black	C	4	1-3
hairy	C	4	1-3
Pigweed			
redroot	C	8	1-8
smooth	C	8	1-8
spiny	C	8	1-8
Poinsettia, wild	C		
Puncturevine	C		
Purslane, common	C		
Pusley, Florida	C		
Sida, prickly	C*		
Ragweed,			
common	R	4R	1-3
giant	R	4R	1-3
Sage, barnyard		R	1-3
Smartweed			
ladysthumb	C	4	1-3
Pennsylvania	C	4	1-3
Spurge			
prostrate	C	4	1-3
spotted	C	4	1-3
Sunflower	C*	4	1-3
Velvetleaf	C*	4	1-3
Thistle, Canada		R	1-3

* When PURSUIT is soil applied, these weeds are more consistently controlled by preplant incorporated treatments.

48/62

GRASS WEEDS*

Weeds Controlled**	<u>SOIL APPLIED</u> (inches)	<u>POSTEMERGENCE</u>	
		Maximum Leaf Stage	Size
Barnyardgrass	R	3	1-3
Crabgrass,			
large	R	3	1-3
smooth	R	3	1-3
Cupgrass, woolly***		3	1-3
Foxtail,			
giant	C	6	1-6
green	C	3	1-3
yellow	C	3	1-3
Goosegrass	R		
Johnsongrass,			
seedling	C	6	1-8
rhizome		R	6-12
Millet, wild proso	R	R	1-3
Panicum,			
fall	R		
Texas	R		
Red rice		3	1-3
Shattercane	R	6	1-8
Signalgrass, broadleaf	R	4	1-8
Sorghum, alnum	R	6	1-3

SEDGES

Nutsedge			
purple	R	R	1-3
yellow	R	R	1-3

* Preplant incorporated treatments of PURSUIT are more consistent for grass control.

** PURSUIT is active against many broadleaf and grass species. However, when heavy grass or common lambsquarters pressure is anticipated, PURSUIT should be used in combination with a registered soil-applied grass herbicide (such as PROWL herbicide) for optimum control (See HERBICIDE COMBINATIONS section).

***PURSUIT only controls emerged woolly cupgrass

HERBICIDE COMBINATIONS

49/62

GRASS WEEDS

Use a soil applied grass herbicide (such as PROWL® 3.3 EC) to control grass weeds not on the PURSUIT label and to enhance the control of certain broadleaf weeds such as common lambsquarters and pigweeds. Refer to the PROWL 3.3 EC (or other grass herbicide) label for specific use recommendations, rates and precautions.

When applied as directed, PURSUIT preplant incorporated or preemergence combination treatments with PROWL 3.3 EC, TRI-4®, trifluralin, FrontierOutlook, Lasso or Dual will control the weeds listed in following table, in addition to those controlled by PURSUIT alone.

	PROWL 3.3 EC ^a	trifluralin ^b	Lasso	Dual	Outlook
GRASSES					
Barnyardgrass	X	X	X	X	X
Crabgrass, smooth	X	X	X	X	X
Crabgrass, large	X	X	X	X	X
Crowfootgrass	X	X			
Goosegrass	X	X	X	X	X
Millet, wild proso	X	X			
Panicum, fall	X	X	X	X	X
Panicum, Texas	X	X			
Sandbur, field	X	X			
Shattercane	X ^b	X			
Signalgrass, broadleaf	X ^b	X	X	X	X
Witchgrass	X	X	X	X	X

^a Preplant incorporated tank-mixture applications of PURSUIT plus PROWL 3.3 EC will suppress the growth of itchgrass, and rhizome Johnsongrass.

^b Preplant incorporated treatments only.

A selective postemergence grass herbicide such as ~~PRESTIGE⁷, Select⁹, Poast Plus~~, ~~Fusilade⁶, 2000~~, ~~Fusilade⁶, DX~~, ~~Fusion⁶, Assure² or Option¹⁰~~ may be mixed with PURSUIT to control volunteer corn or grasses not controlled by PURSUIT. For best results use ~~SUN-IT-II~~ or a crop oil concentrate AND liquid fertilizer with grass herbicide tank-mixtures.

PURSUIT + ~~PRESTIGE-POAST PLUS~~ For Enhanced Grass Control

Apply PURSUIT at the rate of 4 ounces per acre. Refer to the table below for the appropriate rate of ~~PRESTIGE-POAST PLUS~~ herbicide for enhanced grass control. The addition of ~~PRESTIGE-POAST PLUS~~ to PURSUIT at the recommended rates will control the grasses listed below. (Refer to the ~~PRESTIGE-POAST PLUS~~ label for additional weeds controlled).

PRESTIGE-POAST PLUS Rate * (ounces per acre)	Annual Grasses Controlled	Size (inches)
12 oz.	Wild Proso Millet Shattercane	4-10" 3-12"
16 oz.	Foxtail, Giant Junglerice Panicum, Fall Texas Signalgrass, Broadleaf	3-8" 3-8" 3-8" 3-8" 3-8"
20 oz.	Volunteer Corn	4-10"
24 oz.	Barnyardgrass Crabgrass, Large Smooth Cupgrass, woolly Foxtail, Green Yellow Goosegrass Johnsongrass, Seedling Sprangletop, Red Witchgrass	3-8" 3-6" 3-6" 3-8" 3-8" 3-8" 3-6" 3-8" 3-8" 3-8"

*If a mixture of grasses are present, use the highest rate indicated for the grasses present.

The addition of ~~PRESTIGE-POAST PLUS~~ herbicide to PURSUIT enhances the grass control, especially when heavy infestations of grass exist. It also provides control of grasses not controlled by PURSUIT. In some cases the activity of ~~PRESTIGE-POAST PLUS~~ may be reduced when mixed with PURSUIT. The reduction in activity may be overcome by delaying the application of ~~PRESTIGE-POAST PLUS~~ herbicide 7 days following the application of PURSUIT. If ~~PRESTIGE-POAST PLUS~~ is applied first, wait 3 days before applying PURSUIT.

For optimum control, apply the tank mixture to actively growing weeds at the sizes indicated in the table above, (for sequential applications refer to application rates and weeds sizes indicated in the PURSUIT and ~~PRESTIGE-POAST PLUS~~ labels). Refer to the ~~PRESTIGE-POAST PLUS~~ label for additional information regarding application rates, restrictions, precautions, weeds controlled, adjuvants recommended and other information.

BROADLEAF WEEDS

Broadleaf herbicides that can be tank-mixed with PURSUIT include ~~STATUSULTRA~~ BLAZER, Basagran, ~~Blazer⁷, Bronco⁴, Cobra⁹Cobra⁶, Galaxy⁷, FirstRate¹, Gramoxone Extra, Storm⁷, Flexstar⁶-Flexstar³ or Reflex⁶Reflex³~~. Roundup Ultra may be tank-mixed with PURSUIT to aid in control of certain weeds only in Roundup Ready Soybeans. See the Roundup Ultra label for rates

and weeds controlled and other restrictions. Certain herbicides should not be applied with PURSUIT
(see PRECAUTIONS section).

51/62

PURSUIT + STATUS-ULTRA BLAZER For Enhanced Control of Common Ragweed and Pigweeds (including tall and common waterhemp).

The addition of STATUS-ULTRA BLAZER to PURSUIT at the recommended rates will enhance the control of several broadleaf weeds, including common and giant ragweed, pigweed species and waterhemp. (Refer to the STATUS-ULTRA BLAZER label for additional weeds controlled.)

When tank-mixing STATUSULTRA BLAZER with PURSUIT, apply PURSUIT at the rate of 4 ounces per acre. Apply STATUSULTRA BLAZER at the following rates, depending on weed size:

	STATUSULTRA BLAZER Rate (ounces per acre):*		
	8 - 10 oz.	12 - 14 oz.	16 - 20 oz.
Weeds/Weed Size	1 - 4"	4 - 6"	6 - 8"
Common ragweed Pigweed species, Waterhemp, tall common			
Giant Ragweed		1 - 6"	6 - 8"***

* Use the higher rate if common ragweed is present or the weed population is high.

** Use the 20 ounce/acre rate if giant ragweed is 6 - 8 inches tall.

STATUSULTRA BLAZER Sequential Application Rates

When applying STATUSULTRA BLAZER following a PURSUIT application (sequential), apply STATUSULTRA BLAZER at the following rates:

	STATUSULTRA BLAZER Rate (ounces per acre):*		
	10 - 12 oz.	14 - 16 oz.	18 - 24 oz.
Weeds/Weed Size	1 - 4"	4 - 6"	6 - 8"
Common ragweed Pigweed species, Waterhemp, tall common			
Giant Ragweed		1 - 6"	6 - 8"***

* Use the higher rate if common ragweed is present or the weed population is high.

** Use the 24 ounce/acre rate if giant ragweed is 6 - 8 inches tall.

~~Blazer, Flexstar, Reflex, or Cobra may be used in tank mixture with PURSUIT instead of STATUSULTRA BLAZER for enhanced control of common ragweed or pigweeds. Refer to the Blazer, Flexstar, Reflex or Cobra labels for application rates and weed size recommendation.~~

PURSUIT + FirstRate for Enhanced Control of Ragweed Species.

FirstRate may be tank-mixed with PURSUIT to aid in the control of common and giant ragweed. See the FirstRate label for recommended rates and precautions.

53/62

PURSUIT + Sulfentrazone Containing Compounds.

PURSUIT provides control of many grasses and broadleaf weeds when applied to the soil or applied postemergence to weeds. It also provides season-long control of many weeds. Sulfentrazone-containing products (such as Authority¹ ~~Broadleaf~~ or Canopy² XL) may be tank-mixed with PURSUIT in soil applications for enhanced weed control in soybeans.

PURSUIT may be applied postemergence to soybeans previously treated with sulfentrazone-containing products.

NOTE: Sulfentrazone-containing products are only labeled for soil applications to soybean.

5/62

PURSUIT + ~~Concert or Pinnacle~~ Harmony GT For Enhanced Control of Common Lambsquarters.

For optimal weed control management, apply a soil applied grass herbicide such as PROWL[®], TRI-4[®], or ~~Treflan⁺~~-Trifluralin followed by PURSUIT postemergence. If common lambsquarters are not adequately controlled by the soil applied treatment, Harmony GT~~Concert, Concert SP, or Pinnacle~~ herbicide may be tank-mixed with PURSUIT for additional activity.

The addition of Harmony GT~~Concert, Concert SP, or Pinnacle~~ herbicides to PURSUIT may cause severe injury and/or stunting to soybeans, especially when applied under hot, humid conditions. The USER ASSUMES ALL RISKS AND CONSEQUENCES associated with applications of this tank-mixture to soybeans.

When tank-mixing Harmony GT~~Concert or Pinnacle~~ with PURSUIT, use the following rates:

PURSUIT - 4 ounces per acre

AND

Harmony GT - 1/24 ounce per acre~~Concert - 0.25 ounce per acre~~

OR

~~Concert SP - 0.25 ounce per acre (one 2 ounce soluble pack treats 8 acres)~~

OR

~~Pinnacle - 0.125 ounce per acre~~

Add to the spray mixture:

Non-ionic surfactant - 1 quart per 100 gallons (0.25% v/v)

AND

Liquid nitrogen based fertilizer (such as 28%N, 32%N, or 10-34-0) at the rate of 1.25 to 2.5 gallons per 100 gallons of spray solution. Instead of a liquid fertilizer, spray grade ammonium sulfate may be used at the rate of 12-15 lbs. per 100 gallons of spray solution.

Apply to 1-3 trifoliolate stage soybeans only.

Other Tankmixture Combinations

PURSUIT + SCEPTER DG for Volunteer Corn and Common Sunflower

The application of PURSUIT plus ~~SCEPTER or SCEPTER DG~~ may be applied to states or portions of states described as Region 2 or Region 3 on the ~~SCEPTER or SCEPTER DG~~ labels, and the following counties in South Dakota: Yankton, Bon Homme, Hutchinson, McCook, Hanson, Davison, Miner, Lake and Kingsbury. Refer to the respective labels for the recommended use area. Do not use this tank mixture in North Dakota or in Minnesota north of state highway #210.

Apply the products at the following rate:

PURSUIT 4 oz/A
AND
~~SCEPTER 2 oz/A (liquid)~~
 SCEPTER DG 0.53 oz/A*

* At the rate of 0.53 ounces per acre, one 14 ounce soluble bag of SCEPTER DG will treat 26.4 acres.

The tank mixture of PURSUIT plus ~~SCEPTER~~ or SCEPTER DG will suppress volunteer corn. Apply to volunteer corn up to 10 inches in height. The tank-mixture of PURSUIT and SCEPTER ~~or~~ SCEPTER-DG will enhance the control of common sunflowers. Apply to sunflowers up to 3 inches in size. Refer to the ~~SCEPTER~~ or SCEPTER DG label for additional weeds controlled.

A postemergence application of PURSUIT plus SCEPTER DG ~~products~~ will NOT suppress volunteer CLEARFIELD corn (field corn hybrids which possess tolerance or resistance to imidazolinone herbicides i.e., PURSUIT and SCEPTER DG).

~~PURSUIT + Metribuzin or Linuron Containing Products~~

~~PURSUIT may be applied either preplant incorporated or preemergence in tank mix combination with metribuzin (Sencor¹⁺ or Lexone²) or preemergence with linuron (Lorox² or Linex¹⁺²). PURSUIT may also be applied preplant incorporated, followed by a preemergence application of metribuzin or linuron. The tank mix or sequential application of PURSUIT and metribuzin will enhance the control of common cocklebur and common ragweed in addition to the weeds controlled by a soil application of PURSUIT. The addition of linuron to PURSUIT will enhance the control of common ragweed.~~

~~When tank-mixing herbicides, always follow the most restrictive label.~~

APPLICATIONS TO SOYBEANS IN NORTH DAKOTA AND MINNESOTA (north of highway #210).

Application Rate:

Apply PURSUIT at 3 ounces per acre postemergence only.

Weeds Controlled	POSTEMERGENCE	
	Maximum Leaf Stage	Size (inches)
Cocklebur, common*	4	1-4
Kochia (non-ALS resistant)	4	1-3
Mustard, species	4	1-3
Nightshade,		
black	4	1-3
Eastern black	4	1-3
hairy	4	1-3
Pigweed, redroot	4	1-4
Wild oats**	3	1-4

* For control of common cocklebur, add STATUSULTRA BLAZER herbicide at the rate of 12 ounces per acre to the spray solution.

**PURSUIT will reduce competition from wild oats.

Sugarbeet production can be reduced when grown in soil conditions with a pH less than 6.5. If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

57/62

58/62

Use of PURSUIT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

EXCEPTIONS TO ROTATIONAL CROP GUIDELINE

Barley: (States of Delaware, Indiana, Kentucky, Maryland, New Jersey, Ohio, Pennsylvania, and Virginia only). Barley may be planted 4 months following a PURSUIT application in these states.

CLEARFIELD canola: CLEARFIELD varieties of canola, such as Pioneer 45A71 and Pioneer 46A76, may be planted as a rotational crop the next season after an application of PURSUIT herbicide at label rates on registered crops.

Corn inbred lines: Corn inbred seed lines may be planted the year following an application of PURSUIT. Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, BASF has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of BASF, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT shall be assumed by the user.

Sweet corn and popcorn varieties: (States of Illinois, Indiana, Iowa, Minnesota, Ohio, Tennessee, and Wisconsin only). Sweet corn and popcorn varieties may be planted the year following an application of PURSUIT. Some sweet corn and popcorn varieties may be injured when planted at less than 18 months following an application of PURSUIT herbicide. Before planting sweet corn for processing, contact the processor company for information and recommendations regarding the tolerance of sweet corn varieties planned for fields treated with PURSUIT the previous year. **DO NOT** plant fresh market sweet corn varieties prior to 18 months after PURSUIT use. Before planting popcorn, contact the popcorn company for information and recommendations regarding the tolerance of popcorn varieties planned for fields treated with PURSUIT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of BASF, **ALL RISKS AND CONSEQUENCES ASSOCIATED WITH PLANTING SWEET CORN OR POPCORN VARIETIES INTO FIELDS TREATED PREVIOUSLY WITH PURSUIT SHALL BE ASSUMED BY THE USER.**

Stunting and maturity delay or other adverse effects may result when sweet corn or popcorn are planted following PURSUIT use.

Certain vegetable crops: (States of Alabama, Delaware, Florida, Georgia, Indiana, Kentucky, Maryland, New Jersey, North Carolina, Pennsylvania, South Carolina, and Virginia only). The following crops may be planted 18 months following the last application of PURSUIT: bahaigrass, cabbage, cantaloupe, cucumber, Irish potato, onion, sweet potato transplants, sweet pepper transplants, tomato transplants, and watermelon.

Cotton: (States of North Carolina, South Carolina and Virginia only) Cotton may be planted nine and one-half months after an application of PURSUIT if all of the following criteria are met:

- PURSUIT applied to peanuts only
- Soil texture is sandy loam or loamy sand only
- Greater than 16 inches of rainfall and/or irrigation is received following application of PURSUIT through October of the application year

59/62

Field Corn and Field Corn Grown for Seed: (Arizona, Hawaii, Idaho, Montana, Nevada, Oregon, Utah, Washington, and Wyoming). Nine and one-half months after PURSUIT application.

Snap Beans: When applied at no more than 1.5 ounces per acre to snap beans in the use areas defined on this label, snap beans may be replanted at anytime after application of PURSUIT.

Wheat: Wheat may be planted 3 months following a PURSUIT application in areas east of Interstate highway I-35.

When PURSUIT is applied at no more than 3 ounces per acre to edible legumes in the use areas described the following rotational restrictions apply:

- Chickpeas, lentils and peas may be planted anytime following a PURSUIT application.
- Snap beans may be planted 3 months and barley 4 months following an application of PURSUIT.

60/62

PRECAUTIONS

CLEARFIELD CORN

There should be an interval of at least 45 days between an application of PURSUIT and corn harvest (silage, fodder, or grain). **DO NOT** graze or feed treated corn forage, silage, fodder, or grain for at least 45 days after an application of PURSUIT.

All soil insecticides, including labeled banded or in-furrow applications, may be used in combination with Pioneer imidazolinone resistant (IR) corn hybrids.

Imidazolinone tolerant hybrids from other seed companies may occasionally exhibit injury symptoms when soil insecticides are used in combination with PURSUIT. **DO NOT USE COUNTER[®] 15G** systemic insecticide-nematicide in-furrow with imidazolinone tolerant corn hybrids. Other registered organophosphate insecticides such as banded applications of COUNTER 15G, COUNTER[®] CR[®] or THIMET[®] soil and systemic insecticide, or in-furrow applications of COUNTER CR or other registered carbamate or pyrethroid insecticides may be used in combination with PURSUIT applications. BASF has not tested all hybrids in which the imidazolinone tolerance trait is claimed and cannot be responsible for factors which are beyond its control, such as growing conditions, environmental conditions, grower practices and the specific genetics of each hybrid tolerance to PURSUIT and insecticide applications.

Alfalfa

Do not feed, graze or harvest alfalfa for 30 days following an application of PURSUIT to alfalfa.

Soybeans

If soybeans are furrow irrigated, till the soil prior to planting winter wheat or barley. The beds should be broken up and the soil mixed with tillage equipment set to cut 4-6 inches deep.

There should be an interval of at least 85 days between an application of PURSUIT and soybean harvest.

PURSUIT applications should be made before soybean bloom.

Do not graze or feed treated soybean forage, hay or straw to livestock.

DO NOT tank-mix PURSUIT with clomazone containing herbicides (~~Command⁹ or Commence¹⁵~~). PURSUIT may be applied postemergence following a soil application of ~~Command or Commence~~.

Peanuts

Do not graze or feed treated peanut forage, vines, hay or straw to livestock.

There should be an interval of at least 85 days between an application of PURSUIT and peanut harvest.

Classic may be applied postemergence to peanuts following a PURSUIT application. Refer to the Classic label for specific use recommendations.

DO NOT apply PURSUIT[®] PLUS EC ~~or PASSPORT[™]~~ to peanuts the same year as PURSUIT.

6/1/02

Edible Vegetable Legumes

succulent

Allow at least 30 days between application and harvest of snap beans, lima beans, chickpeas (Arizona and California), English peas, and Southern peas.

Allow at least 60 days between application and harvest of dry edible peas, lentils, chickpeas, red kidney beans, and other dry bean or pea types listed on this label.

General (all crops)

Full rate application of products containing chlorimuron ethyl (Classic, Canopy², XL, Synchrony², Reliance², Gemini², Lorox Plus², Preview², etc.) chloransulam-methyl (FirstRate), flumetsulam (Hornet¹, Python¹, Broadstrike + Dual, Broadstrike + Treflan, Broadstrike Plus Corn), imazaquin (SCEPTER³, SQUADRON[®], TRI SCEPT³, SCEPTER³ O.T.³, SCEPTER[®] 70DG or DETAIL³) or products containing imazethapyr (CONTOUR⁴, RESOLVE⁴, PURSUIT[®] DG or PURSUIT[®] PLUS EC) the same year as PURSUIT may increase the risk of injury to sensitive follow crops. Consult labels for recommended uses of these products in combinations.

Only rotational crops harvested at maturity may be used for feed or food.

In the event of a crop loss due to weather, soybeans, peanuts or CLEARFIELD corn can be replanted. DO NOT work the soil deeper than 2 inches.

For additional information regarding the use of PURSUIT herbicide, call telephone no. 800-942-0500.

¹ ~~Broadstrike~~, FirstRate, Treflan, Balan, Hornet, Python, Sonolan and Lorsban are trademarks of Dow Agro Sciences.

² Accent, Basise, Classic, Harmony GT, Concert, Pinnacle, Staple, Synchrony, Assure, Lexone, Lexone, Lorox, Lorox Plus, and Canopy XL, Gemini, Reliance and Preview are trademarks of E.I. duPont de Nemours & Co., Inc.

³ ~~Exceed and Bravo~~, Bravo S, Dual, are trademarks of Novartis Crop Protection, Inc. Flexstar, Reflex, Vernam, Gramoxone Extra, Spirit and Starfire are trademarks of Syngenta Crop Protection.

⁴ Permit, Lasso, and Roundup Ultra, and ~~Harness and Broneo~~ are trademarks of Monsanto Agricultural Company. ⁵ SUN-IT II is a trademark of Agseco, Inc.

⁸ ~~Buetril~~ ⁵ Buetril, Bugle and Whip are trademarks of Aventis Crop Science.

⁹ ~~Cobra~~ ⁶ Cobra, Prism, Select and Orthene are trademarks of Valent Chemical Co.

¹¹ ~~Sencor~~ ⁷ Sencor is a trademark of Bayer Corporation

¹³ ~~Bravo~~ ⁸ Bravo and Bravo S are trademarks of Fermenta ASC Corp.

⁸ ¹⁴ Solubor is a trademark of U.S. Borax & Chemical Corp.

¹⁵ ~~Command~~ ⁹⁰ Command and ~~Commence~~ are trademarks of FMC Corporation.

^{®/™} Trademarks of BASF.

BASF

62/62