

# RESTRICTED USE PESTICIDE

Due to Oncogenicity,

Formulation No. 2500-1, Restricted Applicators of pesticides are limited to those persons who are registered with the EPA under the provisions of 40 CFR 170.10. The use of this product may be restricted to those persons who are registered with the EPA under the provisions of 40 CFR 170.10. This product has been determined to cause tumors in laboratory animals.

# Ala-Scept

herbicide

## FOR WEED CONTROL IN SOYBEANS

### ACTIVE INGREDIENTS:

Aaachlor [2-chloro-2,6-diethyl-N-(methoxymethyl)acetanilide]

Imazaquin [2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-quinolinecarboxylic acid]

### INERT INGREDIENTS

TOTAL

Contains 3 pounds/gallon of aachlor and 0.1675 pounds/gallon of imazaquin.

EPA Reg. No. 241-311

EPA Est. No. 37420-GA-1

## KEEP OUT OF REACH OF CHILDREN: WARNING! ¡AVISO!

PRECAUCION AL USUARIO: Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada por un agente.

### FIRST AID

**IF IN EYES:** Immediately flush with plenty of water for at least 15 minutes. Call a physician.

**IF ON SKIN:** Immediately flush with plenty of water while removing contaminated clothing. If a hot shower is available, wash thoroughly with soap and water. Wash clothing before reuse. Sensitized persons should avoid further contact and reuse of contaminated clothing.

**IF INHALED:** Remove victim to fresh air. If breathing is difficult, give oxygen. If breathing has stopped, give artificial respiration. Call a physician.

**IF SWALLOWED:** Do not induce vomiting. Give water to drink. Call a physician. **NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.**

In case of emergency endangering life or property involving this product, call Collect, day or night, Area Code 291-635-3100.

See Next Page For Additional Precautionary Statements.  
See Directions For Use Inside.



American Cyanamid Company  
Agricultural Division  
Crop Protection Chemicals Department  
Wayne, NJ 07470 1989

Net Contents: 2.5 gallons

D40  
1/89

P. 201

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

# WARNING

- CAUSES EYE IRRITATION**
- HARMFUL IF SWALLOWED**
- HARMFUL IF INHALED**
- MAY CAUSE ALLERGIC SKIN REACTION**
- AVOID CONTACT WITH EYES, SKIN, OR CLOTHING**
- AVOID BREATHING SPRAY MIST**
- WASH THOROUGHLY WITH SOAP AND WATER AFTER HANDLING**

Wear goggles or face shield, rubber gloves, long trousers, long sleeves, shirt or jacket of tightly woven material, along with boots high enough to cover ankles when transferring and mixing, and when adjusting, repairing or cleaning equipment. Wear rubber boots when pouring from open containers. Clothing which comes in contact with this product must be washed before reuse. Clothing or other materials which become drenched with the concentrated pesticide must be disposed of in a sanitary landfill by incineration or, if allowed by state and local authorities, by burning. Wash thoroughly with soap and water after handling.

### ENVIRONMENTAL HAZARDS

This product contains a herbicide which has been identified as a potential ground water contaminant and there is the possibility that it may leach through the soil to ground water, especially where there is no clay and ground water is near the surface. For a more detailed description of this hazard, consult the chemical manufacturer's data sheet. Do not apply to streams, rivers, and reservoirs.

Care must be taken when using this product to prevent leaks or spills from entering wells, springs, or other water sources. Do not apply to the soil surface or in ditches.

Check valves or antisiphoning devices must be used on all mixing equipment. If excess pesticide, spray mixtures, or rinsate should be according to label use instructions or according to the state pesticide or environmental control agency or the hazardous waste representative at the nearest EPA regional office.

Do not contaminate water when disposing of equipment washwaters.

### DIRECTION OF USE

If a violation of Federal law to use this product in any manner inconsistent with the labeling, DO NOT apply this product through any type of mechanical system. Do not apply this product to soil that is wet or has a high water table. **CHARL WELL BEFORE USE**

### STORAGE AND DISPOSAL

**STORAGE:** STORE AWAY FROM CHILDREN. KEEP FROM THE SUN. Do not store in a warm room and do not store in a room where people live. Do not store in a room and do not store in a room where people live. Do not store in a room and do not store in a room where people live. Do not store in a room and do not store in a room where people live.


**DISPOSAL:** Wastes of this pesticide may cause eye irritation and may be dangerous. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law. If these wastes cannot be disposed of according to label use instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

mental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Empty containers must be cleaned and product residue removed. Containers must be triple rinsed, cleaned, and recycled, or disposed of in a sanitary landfill, or by incineration, or by other methods approved by the local authority. Burned or burned stay out of view.

### USE AREA

ALA SCEPT herbicide can be applied only in the states or parts of states shaded in the following map.



**Northern Use Area:** Includes Illinois, Indiana, Michigan, Ohio, Pennsylvania, West Virginia, Kentucky, Tennessee, Mississippi, Alabama, Georgia, Florida, and parts of the Carolinas and Virginia.

**Southern Use Area:** Includes Alabama, Georgia, Florida, and parts of the Carolinas and Virginia.

**NOTE:** See ROTATIONAL CROP RESTRICTIONS section for recommendations applying to the Northern Use Area.

### GENERAL INFORMATION

Read and carefully observe precautionary statements and all other information appearing on the labeling of all products used in mixtures and sequential treatments.

Some softening of soybean plants may occur with application of this product. This is the result of the herbicide action on the plant and is not an effect of the herbicide.

Do not apply this product to soybean plants that are in the early stages of growth.

Do not apply this product to soybean plants that are in the late stages of growth.

Do not apply this product to soybean plants that are in the late stages of growth.

Do not apply this product to soybean plants that are in the late stages of growth.

Do not graze treated area or feed treated soybean forage, hay or straw to livestock following application.

### ROTATIONAL CROP RESTRICTIONS

The following rotational crops may be planted after applying ALA SCEPT in soybeans:

1. Wheat may be planted in the fall, four months after ALA SCEPT application in soybeans. Rice may be planted in the spring of the year following ALA SCEPT application. (See Northern Use Area Restrictions.)
2. Eleven months after ALA SCEPT application:
  - Barley (See Northern Use Area Restrictions)
  - Corn (See Northern Use Area Restrictions)
  - Field Beans (See Northern Use Area Restrictions)
  - Grain Sorghum (See Northern Use Area Restrictions)
  - Grain Sorghum
  - Grain Sorghum
  - Peas
  - Tobacco

### NORTHERN USE AREA RESTRICTIONS

For the Northern Use Area as defined in the USE AREA section of this label, DO NOT plant the following crops as a rotational crop in the fall or in the spring of the year following ALA SCEPT application:

### SOUTHERN USE AREA RESTRICTIONS

For the Southern Use Area as defined in the USE AREA section of this label, field corn may be planted in the spring of the year following ALA SCEPT application, unless extreme drought conditions develop and less than 15 inches of rainfall or irrigation is received from the date of ALA SCEPT application through October of the year of application. For example, in Alabama, Arkansas, the District of Columbia, Tennessee, Texas, Virginia, and West Virginia, field corn may be planted in the spring of the year following ALA SCEPT application, unless extreme drought conditions develop and less than 15 inches of rainfall or irrigation is received from the date of ALA SCEPT application.

Do not graze or feed treated soybean forage hay or straw to livestock until 90 days after application. Do not graze or feed treated soybean forage hay or straw to livestock until 90 days after last ALA SCEPT application. Do not graze or feed treated soybean forage hay or straw to livestock until 90 days after last ALA SCEPT application.

Do not graze or feed treated soybean forage hay or straw to livestock until 90 days after application. Do not graze or feed treated soybean forage hay or straw to livestock until 90 days after last ALA SCEPT application.

Do not graze or feed treated soybean forage hay or straw to livestock until 90 days after application. Do not graze or feed treated soybean forage hay or straw to livestock until 90 days after last ALA SCEPT application.

Do not graze or feed treated soybean forage hay or straw to livestock until 90 days after application. Do not graze or feed treated soybean forage hay or straw to livestock until 90 days after last ALA SCEPT application.

### MIXING, SPRAYING AND HANDLING INSTRUCTIONS

The use of a mechanical transfer system is required to be used by a mixer loader and/or applicators who treat 300 acres or more annually with pesticides containing alachlor as an active ingredient.

Direct contact or exposure to this product should be minimized. The following instructions for transfer, mixing, and application equipment should be followed in order to minimize exposure. However, no protective clothing requirements are listed in the PRECAUTIONARY STATEMENTS section of this label and therefore, no special clothing or washing the face, eyes, protection of hands.

### Equipment Cleaning and Repair

Cleaning and repair of transfer systems and application equipment is a source of exposure to this product. Care should be taken to minimize exposure during cleaning and repair of transfer systems and application equipment. Whenever possible, these systems or equipment should be rinsed before being cleaned or repaired. When repairs must be made during transfer or application, the equipment should be shutdown, and special care taken to avoid contact with the pesticide.

Always predetermine the compatibility of this product in a tank with the other materials to be mixed. Add this product to the tank last, and by using small proportional quantities in advance, check the compatibility. AVAILABLE FOR FERTILIZER TANK MIXTURES: This product is compatible with water carrier, spray carrier, and fertilizer carrier.

Always predetermine the compatibility of this product in a tank with the other materials to be mixed. Add this product to the tank last, and by using small proportional quantities in advance, check the compatibility.

Make this product or a dilute end use mixture of this product with the appropriate carrier.

1. Place a 20 to 35 mesh screen or wetting pack over the sprayer tank.
2. Through the screen, fill the sprayer tank with water or the appropriate carrier.
3. If a compatibility agent is necessary to improve compatibility, prevent the formation of undesirable and unusable spray droplets, and to prevent wheeling, add it to the carrier already in the tank. Use only compatibility agents created by EPA for this use. Read and follow the label directions for use on the compatibility agent label. Do not use any other compatibility agents.
4. Dump the contents of the sprayer tank into the carrier source. Mix the contents of the sprayer tank with water or the appropriate carrier. Do not spray the contents of the sprayer tank through the screen of the tank. Continue agitation.
5. Complete filling the sprayer tank with carrier. If the sprayer tank is empty, Super is used, add the required Super to the tank. If the sprayer tank is not empty, add the required Super to the tank. After this product and prior to application, Super is used, Super is used. Remove hose from tank immediately after use, and keep it hanging back onto the carrier source.

Through agitation is required until the contents of the tank are sprayed. If the spray mixture is allowed to settle, the contents of the tank must be resuspended before spraying is resumed. This may involve agitation and prolonged agitation.

Keep by pass line on or near bottom of tank to minimize foaming. Nozzle or line strainer screen size should be no finer than 40 mesh. Carefully select correct nozzles to avoid spraying a fine mist. Check for uneven distribution of spray droplets. To reduce loss of the tank, always spray at a minimum application nozzle pressures below 40 psi.

Do not use this product in a tank with other materials. Do not use this product in a tank with other materials.

### STANDARD SPRAYABLE FLUID FERTILIZER COMPATIBILITY TEST

Make this test in a tank always mix evenly throughout a spray mixture. Do not use this product in a tank with other materials. Do not use this product in a tank with other materials.

**BEST AVAILABLE COPY**

- A. Materials Required:**
- Two one quart jars (one jar marked "With" and "Without")
  - TE Aspoon (one level TE Aspoon is equal to one level teaspoon)
  - Sprayable fluid fertilizer
  - The herbicide(s) to be mixed
  - A compatibility agent (the purpose of the adjuvant is to help keep the fertilizer and crop protection chemical in suspension, if this as substance is needed)

- B. Procedure:**
- Add one pint of the sprayable fluid fertilizer that will be used or other herbicide carrier to each jar marked "With" and "Without"
 

<b>WITH</b>	<b>Add One Pint Liquid Fertilizer To Two Quart Jars</b>	<b>WITHOUT</b>
-------------	---	----------------
  - To the jar marked "With" add 1/4 TE Aspoon or 1/2 milliliters of a suitable compatibility agent (shake gently for five to ten seconds to mix) 1/4 TE Aspoon in one pint is the equivalent of two pints per 100 gallons of liquid fertilizer
 

<b>WITH</b>	<b>To Jar Marked "With" Add Compatibility Agent and Shake 5 Min.</b>	<b>WITHOUT</b>
-------------	--	----------------
  - To each jar add the appropriate amount of the herbicide(s). If more than one is used, add them together with the compatibility agent added first. Shake the jar marked "With" 10-15 seconds for five to ten minutes after each addition
 

<b>With</b>	<b>Add Herbicide(s) to Jar Marked "With" and Shake 10-15 Min.</b>	<b>WITHOUT</b>
-------------	---	----------------

**Amount to Be Added Per Pint of Sprayable Fluid Fertilizer (Assuming Volume is 2 1/2 gal/A)**

HERBICIDE	RATE A	Level TE Aspoons	Milliliters
Wettable Powders	1 pt.	1 1/2	15
or		1	10
Dry Fertilizer	1 pt.	4 1/2	45
or		3	30
or		2 1/2	25

HERBICIDE	RATE A	Level TE Aspoons	Milliliters
Emulsifiable Concentrate	1 qt.	1	10
Emulsifiable Concentrate	1 qt.	1	10
Emulsifiable Concentrate	2 qt.	2	20
Emulsifiable Concentrate	3 qt.	3	30
Soluble Concentrate	1 qt.	1	10
Soluble Concentrate	2 qt.	2	20

This compatibility agent is designed to be used only with the maximum labeled rate of herbicide(s) and fertilizer. Use of herbicide rates in excess of the label rate will result in a herbicide rate in excess of the label rate. Regardless of spray volume, the herbicide rate should not exceed the label rate. If the herbicide rate is in excess of the label rate, the herbicide rate should be reduced to the label rate.

- C. OR Application and Precautions:**
- Apply herbicide(s) immediately after mixing with the liquid fertilizer.
  - If a compatibility agent is used, mix for five minutes after the final addition of herbicide(s) to allow for the formation of large flocs. Suspend the mixture for 15 minutes. Observe if the herbicide(s) cannot be physically mixed with the liquid fertilizer but remains as small oily particles in the solution. If incompatibility in any form develops (either above or occurs in the jar "with" the compatibility agent added), the liquid fertilizer and the herbicide(s) should not be together in the same spray tank. If incom-

patibility develops, the herbicide(s) should be mixed with the adjuvant but not in the same tank as the liquid fertilizer. An adjuvant is recommended.

Both jars should be shaken thoroughly and the herbicide(s) applied for one-half hour after the application of liquid fertilizer and adjuvants can be suspended. If the herbicide(s) normally will settle to the top after standing, wettable powders will either settle to the bottom of the tank or jar or float to the top, depending upon the density of the fertilizers.

If the herbicide(s) is compatible with fluid fertilizer in the foregoing test without having to use a compatibility agent, fluid fertilizer may be used for the pre-mixing. If this is not compatible without the compatibility agent, the herbicide(s) should be pre-mixed with water before adding to the spray tank.

**APPROVED APPLICATION TIMING AND METHODS**

**Preplant Incorporation:** This product may be mixed into the soil using shallow incorporation equipment any time within 7 days prior to planting. Apply the recommended treatment rate to the soil surface as a broadcast application. If weeds emerge after treatment, rotary hoe or shallowly cultivate immediately to improve performance. Refer to the label for additional information.

If soybeans are planted to best apply and incorporate after bed formation.

**NOTE:** The incorporation of this product on coarse soil types may result in weed control that is not uniform due to rainfall or irrigation.

**Preplant Surface Application:** Sprayable fluid fertilizer should be applied to the soil surface of the soil. Equipment should be operated at a minimum design speed for incorporation to ensure adequate mixing and distribution of the herbicide treatment in the soil. Equipment design should be such that the herbicide is applied to the soil surface and will not be lost to the atmosphere. Seed conditions, including moisture content and crop residue, may not be suitable to allow thorough and uniform mixing with the equipment used for 1 pass incorporation.

Contact American Cyanamid Company for additional information regarding recommended equipment and methods for surface broadcast application.

**2 Pass Incorporation:** When 2 pass incorporation is used, spray and incorporate the herbicide treatment into the upper 2 to 4 inches of the soil with equipment that will be capable of **DEEPER THAN 4 INCHES**. The second pass must be made at an angle and deeper than the first pass to ensure uniform distribution of the herbicide treatment in the soil.

**Preemergence Surface Application:** This product may be applied to the soil surface before or after the soil has been prepared. Apply with the appropriate equipment and follow the label directions for application. If weeds emerge after the herbicide treatment, rotary hoe or shallowly cultivate immediately to improve performance. Refer to the label for additional information.

**Postemergence Surface Application:** Apply before soybeans exceed the **unifoliate stage of growth** (first two true leaves). Weeds which are emerged at the time of application may not be controlled by this product. Rainfall or overhead sprinkler irrigation is required after application to move the herbicide treatment into the weed germination zone to control unemerged weeds. If weeds are emerged at application or after treatment, shallowly





Apply the product using the sprayer. Apply the recommended rates of the product to approved tank mixtures in 3-10 gallons of water per acre. **AVOID DRIFT. DO NOT APPLY DURING INVERSION CONDITIONS WHEN WINDS ARE LIGHTLY OR UNDER ANY OTHER CONDITION WHICH WOULD CAUSE DRIFT.** Coarser sprays are less likely to drift, therefore do not use nozzles or nozzle configurations which dispense spray or fine spray droplets. Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure. Ensure uniform application. To avoid streaked, uneven or overlapped application use mechanical mixing systems only. **HUMAN FLAGGERS PROHIBITED!**

**ALA-SCEPT APPLICATION RATE**

Apply the product at a broadcast rate of 2-2.3 quarts per acre. Refer to the "WEEDS CONTROLLED" section of this label to determine specifics, such as method of application for corresponding weed species.

**TANK MIXTURES  
ALA-SCEPT PLUS LASSO MICRO-TECH**

Use in areas of heavy weed infestations or for improved control of hard to control weeds, such as seedling Johnsongrass, groundcherry, nightshade, smartweed, Texas panicum. Lasso Micro-Tech herbicide can be added to the recommended rate of ALA-SCEPT. A maximum of 2 quarts of Lasso Micro-Tech herbicide can be added to the 2.2 to 2.3 quart per acre rate of ALA-SCEPT.

**ALA-SCEPT plus ROUNDUP or ALA-SCEPT plus  
GRAMOXONE SUPER**

To apply, mix the Roundup or Gramoxone Super with the tank mixture of 2.2 quarts of water per acre immediately before, during or after planting. **DO NOT APPLY DURING INVERSION CONDITIONS WHEN WINDS ARE LIGHTLY OR UNDER ANY OTHER CONDITION WHICH WOULD CAUSE DRIFT.** At a density of stubble weeds, control of weeds may require a higher application rate. The application rate should be increased within the recommended range to insure complete coverage.

**CONTROL OR SUPPRESSION OF EMERGED WEEDS**

**Annual Weeds:** Apply 2 quarts of Roundup herbicide per acre in this tank mixture if weeds are less than 6 inches tall and use 1.5 quarts of Roundup if weeds are more than 6 inches tall for control of many emerged annual grasses and broadleaf weeds. For emergent annual weeds controlled see the "WEEDS CONTROLLED" section of the label booklet for Roundup.  
**Perennial Weeds:** At the time application of the tank mixture, apply 1.5 quarts of Roundup per acre to control many emerged perennial grasses and broadleaf weeds. For perennial weeds controlled see the "WEEDS CONTROLLED" section of the label booklet for Roundup.  
To obtain control of perennial weeds, follow recommendations on the label booklet for Roundup for stage of growth and rate of application.

**USE OF THE TANK MIXTURE FOR BERMUDAGRASS OR  
JOHNSONGRASS CONTROL IS NOT RECOMMENDED**

**NOTE:** When using the tank mixture of ALA-SCEPT plus Roundup herbicide per acre, **DO NOT** use the tank mixture with Lasso Micro-Tech herbicide included in the tank mixture.

**GRAMOXONE SUPER**

**NOTE THAT GRAMOXONE SUPER HERBICIDE IS A POISON WHICH MAY BE FATAL OR HARMFUL TO HUMANS AND DOMESTIC ANIMALS.**

**Broadcast Treatment:** Apply 1-1.2 to 2-1.2 pints of Gramoxone Super to this tank mixture immediately before, during or after planting of the CROP EMERGENCE. Use the 2-1.2 pint rate if any weeds are 4 inches tall. This mixture may not control weeds taller than 6 inches. An increase in stubble, crop residue or weeds increases spray volume and should be increased within the recommended range for complete coverage. Add a nonionic spreader/surfactant approved for use on growing crops containing at least 75 percent surfactant active agent at 8 fluid ounces per 100 gallons of diluted spray. Refer to the Gramoxone Super label for additional information.

**SEQUENTIAL TREATMENT**

ALA-SCEPT and Roundup are registered for use on a wide range of weeds and grasses in Florida, Georgia, Louisiana, Mississippi, Alabama, Arkansas, California, Texas, and Tennessee. Gramoxone Super is registered for use on a wide range of weeds and grasses in Florida, Georgia, Louisiana, Mississippi, Alabama, Arkansas, California, Texas, and Tennessee.

**DISCLAIMER**

The label instructions for the use of this product, their interpretation, and their performance are based on research and field use. The product is not a guarantee of results and should be used carefully. However, the use of this product may eliminate all risks inherent and associated with use of this product, including injury, ineffectiveness, or other unintended consequences, may result. Cause of such factors as weather conditions, presence of other plants, or the use of, or application of, the product in a manner not specified on the label are beyond the control of American Cyanamid Company. American Cyanamid Company makes no other express or implied warranty including other express or implied warranty of FITNESS or MERCHANTABILITY.

INBOOK™, INTEG™ is protected by the following Patents: U.S.A. 4680083, 4592572, 4675062, Canada 422793, and others.



# RESTRICTED USE PESTICIDE

Due to Oncogenicity

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification. The use of this product may be hazardous to your health. This product contains alachlor, which has been determined to cause tumors in laboratory animals.



FOR WEED CONTROL IN SOYBEANS

**ACTIVE INGREDIENTS** Alachlor (2-chloro-2,6-diethyl-  
N-(methoxymethyl)acetamide) 34.5%  
Imazaquin, 2-(4,5-dihydro-4-methyl-4,1-methyl-5-oxo-  
1H-imidazol-2-yl)-3-quinolinecarboxylic acid 2.0%  
**INERT INGREDIENTS** 63.5%  
**TOTAL** 100.0%

Contains 3 pounds/gallon of alachlor and 0.1875 pounds/gallon of imazaquin.

EPA Reg. No. 241-311

EPA Est. No. 37429-GA-1

**KEEP OUT OF REACH  
OF CHILDREN  
WARNING!  
¡AVISO!**

**PRECAUCION AL USUARIO:** Si usted no lee, entiende, no sabe interpretar o no está seguro de lo que dice la etiqueta le haya sido explicada ampliamente.

**FIRST AID**

**IF IN EYES:** Immediately flush with plenty of water for at least 15 minutes. Call a physician.

**IF INHALED:** remove victim to fresh air. If not breathing give artificial respiration preferably mouth to mouth. Get medical attention.

**IF ON SKIN:** immediately flush with plenty of water while removing contaminated clothing. As soon as soap is available wash thoroughly with soap and water. Wash clothing before reuse. Sensitized persons should avoid further contact and reuse of contaminated clothing.

**IF SWALLOWED: DO NOT** induce vomiting. Give large quantities of water. Call a physician. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

In case of emergency endangering life or property involving this product, Call Collect, day or night, Area Code 201-835-3100.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC  
ANIMALS**

CAUSE SEVERE IRRITATION. HARMFUL IF SWALLOWED. HARMFUL IF INHALED. MAY CAUSE ALLERGIC SKIN REACTION. AVOID CONTACT WITH EYES, HAIR OR CLOTHING. AVOID BREATHING VAPORS. WASH THOROUGHLY WITH SOAP AND WATER AFTER HANDLING.

Wearing eye protection and rubber gloves is preferable to wearing clothing that gets dirty and requires frequent washing. Wash thoroughly after spraying, stirring and mixing, and when adjusting, repairing or cleaning equipment. Wear rubber boots when pouring from open containers. Clothing which comes in contact with this product must be washed before reuse. Clothing or other materials which become saturated with the concentrated pesticide must be disposed of in a sanitary and safe manner or if allowed by state and local authorities by burning. Wash thoroughly with soap and water after handling.

**ENVIRONMENTAL HAZARDS**

This product contains alachlor, which has been identified in limited ground water sampling and there is the possibility that it can leach through the soil to ground water, especially where soils are coarse and ground water is near the surface. Following application, and during rainfall periods that cause runoff, this chemical may reach surface water bodies such as streams, rivers, and reservoirs. Do not discharge into any water body. Do not use in areas where runoff may enter a stream or lake. Do not use in areas where runoff may enter a stream or lake. Do not use in areas where runoff may enter a stream or lake. Do not use in areas where runoff may enter a stream or lake.

**DIRECTIONS FOR USE**

For use only by persons certified with this product. See label for complete directions for use. Do not use this product in any way that is prohibited by the label. Do not use this product in any way that is prohibited by the label. Do not use this product in any way that is prohibited by the label.

**STORAGE AND DISPOSAL**

**STORAGE:** STORE ABOVE 32°F TO KEEP PRODUCT IN SOLUTION. Below 32°F crystals may form and settle to the bottom. If crystals form, place in a warm room and roll and/or shake the container frequently to redissolve before using.

**DISPOSAL:** Wastes of this pesticide may cause eye irritation and may be dangerous. Improper disposal of excess pesticide spray mixtures or residues is a violation of Federal law. If these wastes cannot be disposed of according to label use instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Emptied container retains vapor and product residue. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary and safe manner or if allowed by state and local authorities by burning. If burned, stay out of smoke.

See Directions for Use on Label.



Net Contents: 2.5 gallons

D40 1 89

**BEST AVAILABLE COPY**