


Reg no: 241-317

PM-25

175



EVENT™

Grass Growth Regulator

For Use on Tall Fescues, Perennial Ryegrass, Bluegrasses, and Bahiagrasses

ACTIVE INGREDIENTS:

Ammonium salt of imazethapyr [(+)-2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1-H-imidazol-2-yl]-5-ethyl-3-pyridinecarboxylic acid]*..... 17.26%

Ammonium salt of imazapyr [2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1-H-imidazol-2-yl]-3-pyridinecarboxylic acid]]*..... 0.64%

INERT INGREDIENTS..... 82.10%

TOTAL.....100.00%

*Equivalent to 16.3% imazethapyr and 0.6% imazapyr or 1.46 pounds acid per gallon.

EPA Reg. No. 241-

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN

CAUTION!

PRECAUCION!

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

STATEMENT OF PRACTICAL TREATMENT

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF IN EYES: Flush with plenty of water.


In case of an emergency endangering life or property involving this product, call collect, day or night, Area Code 201-835-3100.

See Next Page for Additional Precautionary Statements

Net Contents: 1 gallon (3.78 liters)

D40 7/13/88

™Trademark of American Cyanamid Company



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

CAUTION!

Harmful if inhaled or absorbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water or wetlands.
DO NOT contaminate water when disposing of equipment washwaters.

GENERAL INFORMATION

EVENT reduces the foliar growth of established tall fescue, perennial ryegrass, bluegrass, and bahiagrass. EVENT controls seedhead production by tall fescue and controls or suppresses seedhead production by perennial ryegrass, bluegrass, and bahiagrass.

EVENT may be used on limited care-low maintenance areas, such as roadsides, airports, fairgrounds, and golf course roughs, and limited wear areas such as industrial, institutional, and cemetery grounds.


DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Observe all cautions and limitations on this label and on the labels of products used in combination with EVENT. When EVENT is used in tank mixture with another product, refer to each label for rates, methods of application, proper timing, limitations and precautions. ALWAYS USE in accordance with the most restrictive label limitations. EVENT cannot be mixed with any product having a label prohibiting against such mixing. Keep containers closed to avoid spills and contamination. DO NOT apply EVENT through any type of irrigation system.

STORAGE AND DISPOSAL

PROHIBITIONS: KEEP FROM FREEZING
DO NOT store below 32°F.
DO NOT contaminate water, food, or feed by storage or disposal.



PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, by incineration or, if allowed by State and Local authorities by burning. If burned, stay out of smoke.

DOSAGE RATES AND TIMING

Apply EVENT at a rate of 8 to 10 fluid ounces per acre. Use the higher rate for more extended periods of foliar growth suppression. Apply after grasses are at 100 percent greenup and have at least two inches of vertical growth. On finer turfgrass areas where more uniform quality is desired, apply after two mowings. For seedhead suppression, apply prior to the time of seed stalk elongation. EVENT may be applied to mixed grass-legume (e.g., crown vetch) stands for legume release.

ADDITION OF SURFACTANTS

THE ADDITION OF A SURFACTANT TO EVENT SPRAYS IS REQUIRED. Add a nonionic surfactant containing at least 80 percent active ingredient at 0.25 percent v/v of the spray (2 pints per 100 gallons of spray mixture). An antifoam agent may be added to the tank, if needed.

SPRAYING INSTRUCTIONS

Uniformly apply EVENT sprays with a properly calibrated sprayer in sufficient volume to insure adequate coverage. AVOID DRIFT onto the foliage of desirable nontarget plants or injury may result. AVOID overlaps when spraying. A spray indicator dye is useful for preventing overlaps. DO NOT apply when wind velocity is greater than 10 mph.

HERBICIDE COMBINATIONS

EVENT may be used in combination with EMBARK* or TELAR** for broad-spectrum grass seedhead suppression. To use, mix 2 to 4 ounces of EVENT with either 8 ounces of EMBARK or 1/8 to 1/4 ounce of TELAR per acre.

Where control of broadleaf weeds is desirable, EVENT may be tank-mixed with most broadleaf herbicides currently registered for use in turf. EXCEPT when in combination with TELAR, DO NOT tank-mix with sulfonylurea or imidazolinone herbicides. EVENT may be tank-mixed with methylarsoate herbicides such as MSMA or DSMA. DO NOT tank-mix with other postemergence grass herbicides labeled for use in turfgrasses. EVENT may be used prior to or sequential to all currently registered herbicides EXCEPT there must be a three month interval of grass growth between applications of EVENT and sulfonylurea and imidazolinone herbicide applications.

*Registered Trademark of P.B.I. Gordon

**Registered Trademark of DuPont

MIXING INSTRUCTIONS

Fill the spray tank one-half to three-quarters full with clean water. While agitating, add the required measured amount of EVENT. When thoroughly mixed, add the required measured amount of nonionic surfactant. Fill the tank to required volume and add an antifoam agent, if needed.

When tank mixing EVENT with recommended herbicides, add the herbicide first and make sure it is thoroughly mixed. Then add EVENT and proceed as above.

SPECIAL PRECAUTIONS

USE ONLY on well-established non-stressed grass stands. DO NOT use on newly established stands less than 1-year old.

DO NOT use on highly managed turfgrasses.

DO NOT reseed before three months following an application of EVENT.

DO NOT use on intense wear areas such as athletic fields, playgrounds, or footpaths.

DO NOT use on ornamental plants as injury may result.

DO NOT graze or feed clippings of treated grasses to livestock.


DO NOT apply aerially.

IMPORTANT: Thoroughly clean application equipment after use. Flush tank, pump, hoses, and boom with several changes of water after removing nozzle tips and screens (clean these parts separately).

DISCLAIMER

The label instructions for the use of this product reflect the opinion of experts based on research and field use. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Turf injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the use of, or application of the product contrary to label instructions, all of which are beyond the control of American Cyanamid Company. All such risks shall be assumed by the user.

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use, subject to the risks referred to above.



Any damages arising from breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or values or any other special or indirect damages.

American Cyanamid Company makes no other express or implied warranty, including other express or implied warranty of FITNESS or of MERCHANTABILITY.

AMERICAN CYANAMID COMPANY
AGRICULTURAL DIVISION
VPC VENTURES
WAYNE, NJ 07470

©1988

