

TANK MIXTURE WITH SCEPTER**FOR USE IN SOYBEANS IN LOUISIANA, EASTERN TEXAS,
AND MISSISSIPPI SOUTH OF INTERSTATE 20**

OBSERVE ALL PRECAUTIONARY STATEMENTS AND LIMITATIONS ON THE SCEPTER O.T. AND SCEPTER® LABELS. ALWAYS FOLLOW THE MOST RESTRICTIVE LABEL. SEE THE SCEPTER O.T. LABEL FOR MIXING INSTRUCTIONS, SPRAYING INSTRUCTIONS, APPLICATION INFORMATION, AND FOLLOW CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label should be in the possession of the user at the time of pesticide application. DO NOT apply this product through any type of irrigation system.

TANK MIXTURES

SCEPTER® at 1/3 pint per acre may be applied as a tank mixture with SCEPTER O.T. at 1 pint per acre.

If this tank mixture is used following a soil application of SCEPTER, SQUADRON®, or TRI-SCEPT®, no additional postemergence applications of SCEPTER O.T. or SCEPTER, may be used during the same year. DO NOT apply more than 1-1/3 pints of SCEPTER in any form (SCEPTER, SCEPTER® 70DG, SCEPTER O.T., SQUADRON, or TRI-SCEPT) in a single growing season.

The above tank mixture should be applied early postemergence to soybeans after crop emergence, but before weeds exceed the 2 leaf stage of growth.

The addition of a non-ionic surfactant is required. The non-ionic surfactant should contain at least 80% active ingredient and should be mixed at the rate of 2 pints per 100 gallons of spray mixture.

There must be an interval of at least 90 days before the application of this tank mixture and soybean harvest.

WEEDS CONTROLLED

The above tank mixture will control the following weeds in addition to those listed as being controlled by postemergence applications on the SCEPTER O. T. and SCEPTER labels.

Alligatorweed	(<i>Alternanthera philoxeroides</i>)	Redweed	(<i>Melochia corchorifolia</i>)
Dayflower	(<i>Commelina communis</i>)	Rice Fl. edge	(<i>Cyperus iria</i>)
Eclipta	(<i>Eclipta alba</i>)	Texasweed	(<i>Caperonia palustris</i>)
Red Rice*	(<i>Oryza sativa</i>)		

*For control of red rice, the above tank mixture must follow a soil application of PROWL®, trifluralin, Lasso¹, or Dual². The above tank mixture should be applied before the height of the red rice exceeds 3 inches.

¹Lasso is a trademark of Monsanto Agricultural Products Company.
²Dual is a trademark of Ciba-Geigy Corporation.
 *Registered Trademark of American Cyanamid Company
 PE-12222 2/7/92 SO

