

PM

23

241-321

SCEPTER® O.T.®

herbicide

FOR USE IN SOYBEANS

Page 1 of 12

ACTIVE INGREDIENTS:

Sodium salt of imazaquin (2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-quinolinecarboxylic acid)*..... 5.61%

Sodium salt of acifluorfen: Sodium 5-(2-chloro-4-(trifluoromethyl)phenoxy-2-nitrobenzoate)..... 20.93%

INERT INGREDIENTS 73.46%

TOTAL100.00%

*Equivalent to 5.24%, 2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-quinolinecarboxylic acid.

SCEPTER O.T. contains 2.5 pounds of active ingredients per gallon (0.5 pounds active ingredient of imazaquin and 2.0 pounds active ingredient of acifluorfen per gallon).

EPA Reg. No. 241-321

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN

DANGER!/PELIGRO!

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention.

IF SWALLOWED: Drink promptly a large quantity of milk, egg white, gelatin solution, or if these are not available, large quantities of water. Avoid alcohol. Get medical attention.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF INHALED: Remove victim to fresh air.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

In case of emergency endangering life or property involving this product, call collect, day or night, Area Code 201-835-3100.

See Next Page For Additional Precautionary Statements.

ACCEPTED

29 SEP 1993

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 241-321

AMERICAN CYANAMID COMPANY
AGRICULTURAL PRODUCTS DIVISION
CROP PROTECTION CHEMICALS DEPARTMENT
WAYNE, NJ 07470 ©1993

Net Contents:

®Registered Trademark of American Cyanamid Company

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
DANGER!**

**CAUSES IRREVERSIBLE EYE DAMAGE
HARMFUL IF SWALLOWED,
ABSORBED THROUGH
THE SKIN, OR INHALED**

Avoid contact with eyes, skin, or clothing. Wear goggles or face shield. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. DO NOT apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. DO NOT clean equipment or dispose of equipment washwaters in a manner that will contaminate water resources.


DISCLAIMER

The label instructions for the use of this product reflect the opinion of experts based on research and field use. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, herbicide resistant weed populations, or the use of, or application of the product contrary to label instructions, all of which are beyond the control of American Cyanamid Company. All such risks shall be assumed by the user.

American Cyanamid Company shall not be responsible for losses or damages resulting from use of this product in any manner not set forth on this label. User assumes all risks associated with the use of this product in any manner not specifically set forth on this label.

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use, subject to the risks referred to above. CYANAMID DOES NOT MAKE OR AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED AND EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

BUYER'S EXCLUSIVE REMEDY AND AMERICAN CYANAMID'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF SCEPTER O.T. In no case shall, Cyanamid or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product.



USES WITH OTHER PRODUCTS (TANK MIXES)

If this product is used in combination with any other product except as specifically recommended in writing by American Cyanamid Company, then American Cyanamid Company shall have no liability for any loss, damage, or injury arising out of its use in any such combination not so specifically recommended. If used in a combination recommended by American Cyanamid Company, the liability of American Cyanamid Company shall in no manner extend to any damage, loss or injury not directly caused by the inclusion of the American Cyanamid Company product in such combination use, and in any event shall be limited to return of the amount of the purchase price of the product.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of herbicide application.

Observe all cautions and limitations in this leaflet. The use of SCEPTER O.T. not consistent with this label can result in injury to crops, animals, or persons. Keep container closed to avoid spills and contamination.

DO NOT apply this product through any type of irrigation system.

STORAGE AND DISPOSAL


STORAGE: KEEP FROM FREEZING. DO NOT STORE BELOW 32°F. SHAKE WELL BEFORE USING.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. DO NOT contaminate water, food, or feed by storage or disposal. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. Dispose of excess or waste pesticide by use according to label directions, or contact your State Pesticides or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

SCEPTER O.T. is a selective postemergence herbicide for the control of important broadleaf weeds in soybeans. For best results, apply SCEPTER O.T. to small actively growing weeds.



Within any weed population, there may be the potential for the occurrence of weeds resistant to herbicides when compounds that affect the same site of action are used continuously on the same acre. Therefore, any multiple year weed management program should include, when local agronomic practices allow, as many of the following practices as possible:

- use scouting and weed thresholds to determine the need for herbicide treatment.
- utilize tank-mixes or sequential applications of different mode-of-action herbicides to control the same target weeds
- combine cultivation with herbicide treatments

See your Cyanamid representative for additional information.

PRECAUTIONS:

DO NOT use on crops other than soybeans. Crops other than soybeans, such as cotton, corn and vegetables, may be injured by spray drift or other indirect contact with SCEPTER O.T.

To avoid injury to sensitive crops from spray drift, follow all use directions and precautions in SPRAYING INSTRUCTIONS on Page 8.


To avoid injury to sensitive crops, spray equipment used for SCEPTER O.T. applications must be drained and thoroughly cleaned with water before being used to apply other products to these crops. See Page 10 for ROTATIONAL CROP RESTRICTIONS.

DO NOT use SCEPTER O.T. other than in accordance with the instructions set forth on this label.

Use of SCEPTER O.T. herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

CULTIVATION

A timely cultivation may aid in the control of certain weeds or improve general weed control. After postemergence treatments, wait at least 10 days before cultivating.



BIOLOGICAL ACTIVITY

When SCEPTER O.T. is applied postemergence, it is absorbed through the foliage and roots (root uptake occurs if adequate moisture is received from rainfall or overhead irrigation). After application, SCEPTER O.T. acts in three ways:


1. As a contact herbicide producing rapid necrosis of the plant tissue.
2. As a foliar systemic herbicide with movement to the growing points (roots and leaves) of the weed.
3. And as a soil residual herbicide with activity on some newly germinating susceptible weeds.

Susceptible weeds treated with SCEPTER O.T. stop growing and rapidly become chlorotic and necrotic; especially at the growing points, and either die or become non-competitive with the crop.

CROP RESPONSE

The aciflourfen in SCEPTER O.T. may cause some necrosis, cupping, speckling, and bronzing of soybean leaves. Plants quickly recover without any effect on soybean yields.


There must be an interval of 90 days between the last SCEPTER O.T. application and soybean harvest. DO NOT graze or feed treated soybean forage hay or straw to livestock.



USE AREA

6 7 12

SCEPTER O.T. can be applied only in the states or parts shaded in the following map:



Use Region 1 includes eastern Oklahoma, Arkansas, the Missouri bootheel, Tennessee, North Carolina, South Carolina, Georgia, Florida, Alabama, Mississippi, Louisiana, and eastern Texas.

Use Region 2 includes eastern Kansas (east of U.S. 81; the counties of Cloud, Ellsworth, Harvey, Jewell, Lincoln, Mitchell, McPherson, Ottawa, Republic, Saline, Sedgewick, and Sumner), southeastern Nebraska (east of U.S. 81, south of U.S. 34), Missouri, Illinois (south of S.R. 116 west of Peoria; south of U.S. 24 east of Peoria), Indiana, Ohio, Michigan, Kentucky, Virginia, West Virginia, Pennsylvania, Maryland, Delaware, and New Jersey.

Use Region 3 includes Nebraska (east of U.S. 81, north of U.S. 34 and also that area east of U.S. 283, south of U.S. 30, and west of U.S. 81), South Dakota (east of U.S. 81), Illinois (north of S.R. 116 west of Peoria; north of U.S. 24 east of Peoria), Wisconsin, Iowa, and southern Minnesota.

Note: See USE RATES, SEQUENTIAL TREATMENTS and ROTATIONAL CROP RESTRICTIONS sections for recommendations applying to each Use Region.

WEEDS CONTROLLED

WEEDS CONTROLLED		WEED HEIGHT (inches)	MAXIMUM LEAF STAGE
Cocklebur	(<u>Xanthium strumarium</u>)	up to 6	6
Copperleaf, hophornbeam	(<u>Acalypha ostryifolia</u>)	up to 2	2
Corn ¹ , volunteer	(<u>Zea mays</u>)	up to 8	4
Morningglory			
Bigroot ²	(<u>Ipomoea pandurata</u>)	--	-
Entireleaf	(<u>Ipomoea hederacea</u> var. <u>intergriuscula</u>)	up to 2	3
Ivyleaf	(<u>Ipomoea hederacea</u>)	up to 2	3
Palmleaf	(<u>Ipomoea wrightii</u>)	up to 2	3
Pitted	(<u>Ipomoea lacunosa</u>)	up to 2	3
Smallflower	(<u>Jacquemontia tannifolia</u>)	up to 2	3
Tall	(<u>Ipomoea purpurea</u>)	up to 2	3
Pigweed			
Redroot	(<u>Amaranthus retroflexus</u>)	up to 6	6
Smooth pigweed	(<u>Amaranthus hybridus</u>)	up to 4	4
Waterhemp, Tall	(<u>Amaranthus tuberculatus</u>)	up to 4	4
Sesbania, hemp	(<u>Sesbania exaltata</u>)	up to 12	12
Sida ³ , prickly	(<u>Sida spinosa</u>)	up to 3	3
Smartweed, Pennsylvania	(<u>Polygonum pennsylvanicum</u>)	up to 2	2
Sunflower, common	(<u>Helianthus annuus</u>)	up to 4	4

¹SCEPTER O.T. will control only those field corn hybrids which DO NOT possess tolerance or resistance to SCEPTER O.T. or other imidazolinone herbicides (i.e. PURSUIT®).

²SCEPTER O.T. will provide suppression of bigroot morningglory. Applications should be made when vines are 24 to 36 inches in length.

³SCEPTER O.T. will provide suppression of prickly sida. A nonphytotoxic oil concentrate should be used in place of nonionic surfactant. Use a jar test to ensure compatibility with oil concentrates. DO NOT use oil concentrates when tank mixing SCEPTER O.T. with 2,4-DB.

APPLICATION RATE

Apply SCEPTER O.T. at a rate of 1.0 pint per acre with a nonionic surfactant.

APPLICATION INSTRUCTIONS

Apply SCEPTER O.T. as a postemergence treatment after crop emergence and in accordance with the weeds controlled, weed height, and rates of application information listed in this label.

SCEPTER O.T. can provide residual control of cocklebur and pigweed that may emerge after application. Application should be made to small soybeans so that more herbicide reaches the soil surface. The level of residual control will depend on weed density and canopy closure which affect the amount of SCEPTER O.T. reaching the soil surface and environmental conditions, such as rainfall, at the time of application.

•Registered Trademark of American Cyanamid Company

To maximize weed control following a postemergence SCEPTER O.T. application, wait at least 10 days before cultivating. This timely cultivation will enhance residual weed control, especially under dry conditions.

Apply when weeds are actively growing. DO NOT apply SCEPTER O.T. when weeds have been subjected to stress conditions such as temperature and moisture extremes.

There must be an interval of at least 90 days between the last SCEPTER O.T. application and soybean harvest.

In Use Region 3 as defined by the Use Area section of the label, apply SCEPTER O.T. prior to July 1.

HERBICIDE COMBINATIONS

SCEPTER O.T. PLUS 2,4-DB TANK MIXTURE

SCEPTER O.T. may be tank mixed with 2,4-DB at .015 to .030 lb. a.i./A for improved control of labeled morningglory species. Apply by ground to young actively growing weeds, and before annual morningglories reach 4 inches in height. Observe all use restrictions on 2,4-DB label.

MIXING INSTRUCTIONS

Fill the spray tank one-fourth to one-half full with clean water. While agitating add the required amount of SCEPTER O.T. and as the last ingredient in the tank add the nonionic surfactant. An antifoaming agent may be added to the tank if needed. Maintain agitation while spraying to ensure a uniform spray mixture.

SPRAYING INSTRUCTIONS


GROUND APPLICATIONS:

Uniformly apply with properly calibrated ground equipment in 10 to 30 gallons of water per acre and maintain a sprayer pressure of 30-40 PSI. Use higher gallonage for fields with dense vegetation. Use sprayers equipped for postemergence applications. Use nozzles that provide accurate and uniform application (tapered flat fan spray tips with 80-110 degree spray angles, and tip sizes should not exceed -06 to insure small droplet size of the spray solution; hollow cone nozzle tips may also be used).

A second application may be necessary if additional weeds emerge after treatment (total SCEPTER O.T. applied per season can not exceed 2 pints per acre).

DO NOT apply with ground equipment when wind velocity is greater than 10 mph or when spray may be carried by wind to sensitive crops.

Avoid overlaps when spraying.



AERIAL APPLICATIONS:

Uniformly apply with properly calibrated aerial equipment in 5 or more gallons of water per acre.

To avoid injury to sensitive crops from drift, aerial applicators must adhere to the following SPECIAL USE DIRECTIONS AND PRECAUTIONS:

- Use diaphragm-type nozzles which produce cone or flat spray patterns.
- Use a maximum spray pressure of 40 PSI.
- Nozzle height above ground must be a maximum of 10 feet.
- Nozzles must be pointed toward the rear of the aircraft. The downward angle of the nozzle should not be greater than 20 degrees.
- To minimize wing-tip vortex roll, nozzles or spray boom must not be located any closer to end of wing or rotor than three-fourths the distance from the center of the aircraft.
- A buffer zone must be established between the area to be sprayed and sensitive crops.
- DO NOT spray when wind velocity is greater than 5 mph.

Applicator is responsible for any loss or damage which results from spraying SCEPTER O.T. in manner other than recommended in this label. In addition, applicator must follow all applicable state and local regulations and ordinances in regard to spraying.

ADJUVANT USE

Use a nonionic surfactant at least 80% active at 1 quart per 100 gallons or 0.25% volume per volume (v:v) to enhance the wetting, contact, and uptake activity of SCEPTER O.T. on the plant foliage.

SEQUENTIAL TREATMENTS

For Use Region 1 and Use Region 2 as defined in the USE AREA section of this label; a postemergence application of SCEPTER O.T. may be applied at 1 pint per acre following a soil application of SCEPTER[®], SCEPTER[®] 70DG, SQUADRON[®], or TRI-SCEPT[®] to control weeds listed in the WEEDS (CONTROLLED) section of this label. (See labels of above herbicides for additional weeds controlled by this sequential program.) The total amount of SCEPTER O.T. applied can not exceed 1 pint per acre in USE REGION 2 or 2 pints per acre in USE REGION 1 in one season following a soil application of the herbicides listed above..

In USE REGION 2, refer to the ROTATIONAL CROP RESTRICTIONS section of this label for instructions on planting follow crops when using this program.

[®]Registered Trademarks of American Cyanamid Company

ROTATIONAL CROP RESTRICTIONS

The following rotational crops may be planted after applying SCEPTER O.T. in soybeans:

<u>CROP</u>	<u>USE REGION 1</u>	<u>USE REGION 2</u>	<u>USE REGION 3</u>
Soybeans	No restrictions	No restrictions	No restrictions
Wheat	4 months	4 months ³	See FOOTNOTE 4
Rice	Spring following SCEPTER O.T. application	Spring following SCEPTER O.T. application ³	---
Barley	11 months	11 months ³	See FOOTNOTE 4
Field Corn	11 months ¹	11 months ^{2, 3}	See FOOTNOTE 4
Edible Beans	11 months	11 months ³	11 months
Grain Sorghum	11 months	11 months ³	11 months
Oats	11 months	11 months ³	See FOOTNOTE 4
Peanuts	11 months	11 months ³	11 months
Tobacco	11 months	11 months ³	11 months
Sugar Beets	26 months	26 months	26 months
Other Crops	18 months	18 months	See FOOTNOTE 5

1 For USE REGION 1 as defined in the USE AREA section of this label; field corn may be planted in the spring of the year following a single SCEPTER O.T. application unless extreme drought conditions develop (less than 10 inches of rainfall or irrigation is received within 6 months following the date of application).

2 For USE REGION 2 as defined in the USE AREA section of this label; field corn may be planted in the spring of the year following a single SCEPTER O.T. application unless extreme drought conditions develop (less than 10 inches of rainfall or irrigation is received from two weeks prior to the date of SCEPTER O.T. application through November 15 of the same year).

If the minimum rainfall requirement is not met, only field corn hybrids which possess tolerance or resistance to SCEPTER O.T. and other imidazolinone herbicides may be planted the spring of the year following a SCEPTER O.T. application (SEE NOTE).

3 ROTATIONAL CROP RESTRICTIONS for sequential SCEPTER O.T. application following a soil application of SCEPTER, SCEPTER 70DG, SQUADRON, or TRI-SCEPT (USE REGION 2 only):

For Use Region 2 as defined in the USE AREA section of this label; soybeans may be planted anytime. Barley, edible beans, grain sorghum, oats, peanuts, rice, tobacco, and wheat may be planted 15 months after the last herbicide application. Cotton may be planted 18 months after the last herbicide application.

Only field corn hybrids which possess tolerance or resistance to SCEPTER O.T. and other imidazolinone herbicides may be planted the spring of the year following a sequential SCEPTER O.T. application (SEE NOTE). Other field corn varieties may be planted 15 months after the last herbicide application.


4 For USE REGION 3 as defined in the USE AREA section of this label; field corn, wheat, barley, or oats may be planted as a rotational crop in the fall or in the spring of the year following SCEPTER O.T. application ONLY if SCEPTER O.T. is applied as a postemergence application at 1 pint per acre AND 10 inches of rainfall or irrigation has been received from the date of application through October of the same year.

5 For USE REGION 3 as defined in the USE AREA section of this label; canola, table beets, strawberries, cabbage, tomatoes, potatoes, carrots, celery, cole crops, garlic, onions, spinach, asparagus, cauliflower, and brocoli may be planted 26 months after a SCEPTER O.T. application. Other crops may be planted 18 months after a SCEPTER O.T. application.

NOTE: Contact your chemical dealer, seed supplier, or American Cyanamid to obtain information regarding the availability of imidazolinone tolerant field corn hybrids which are adapted to your area.

ONLY rotational crops harvested at maturity may be used for feed or food.

Use of SCEPTER O.T. herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.



Except where noted on a Supplemental Label or in the SEQUENTIAL TREATMENTS section of this label DO NOT apply products containing chlorimuron ethyl (e.g. Classic¹, Canopy¹, Gemini¹, Lorox Plus¹, Preview¹, etc.), or imazaquin (e.g. SCEPTER, SCEPTER 70DG, SQUADRON, TRI-SCEPT, etc.) the same year as SCEPTER O.T. DO NOT apply products containing imazethapyr (e.g. PURSUIT, PURSUIT[•] PLUS, PASSPORT[•], etc.) the same year as SCEPTER O.T. or injury to follow crops may occur.

DO NOT graze or feed treated soybean forage, hay, or straw to livestock.

¹Trademarks of E. I. duPont de Nemours and Company

[•]Registered Trademarks of American Cyanamid Company

