+1122 24/2765

PASSPORT® herbicide

FOR USE IN SOYBEANS

Ammonium salt of imazethapyr: (\pm) -2-[4,5-dihydro-4-methyl-4-(1-methyl-ethyl)-5-oxo-1H-imidazol-2-yl]-5-ethyl-3-pyridinecarboxylic acid* . . . 2.2%

trifluralin: <, < < trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine . . . 27.5%

*Equivalent to 2.1% (\pm) -2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-lH-imidazol-2-yl]-5-ethyl-3-pyridinecarboxylic acid. **Contains petroleum distillates

PASSPORT contains 2.6 pounds of active ingredients per gallon (2.4 pounds ai of trifluralin and 0.2 pounds acid equivalent of imazethapyr).

EPA Reg. No. 241-325

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN

DANGER/PELIGRO!

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Hold eyelids open and flush with a steady, gentle steam of water for 15 minutes. Call a physician.

IF SWALLOWED: DO NOT induce vomiting. Call a physician or Poison Control Center immediately.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

NOTE TO PHYSICIAN: Because of increased risk of chemical pneumonia or rulmonary edema caused by aspiration of the hydrocarbon solvent, vomiting should be induced only under professional supervision.

In case of an emergency endangering life or property involving this product, call collect, day or night, Area Code 201-835-3100.

See Next Page for Additional Precautionary Statements

American Cyanamid Company
Agricultural Division
Crop Protection Chemicals Department
Wayne, NJ 07470 •1992

Net Contents: 2.5 gallons (9.45 liters)

BEST AVAILABLE COPY

Registered Trademark of American Gyanamid Company

ACCEPTED 24 AUG 1993

Under the Federal Insecticity.
Funging the and Potential to each
as about the pessicide
repotent doubter 141-325
EPA Reg. No. 241-325

20415

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)

DANGER

Causes irreversible eye damage. Harmful if swallowed, inhaled or absorbed through skin. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. DO NOT apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to fish in neighboring aquatic sites. DO NOT contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This label must be in the possession of the user at the time of pesticide application.

Observe all cautions and limitations in this leaflet. Do not use PASSPORT other than in accordance with the instructions set forth on this label. The use of PASSPORT not consistent with this label can result in injury to crops, animals, or persons. Keep container closed to avoid spills and contamination.

DO NOT apply this product through any type of irrigation system.

STORAGE AND DISPOSAL

STORAGE:

)

KEEP FROM FREEZING. DO NOT STORE BELOW 40°F. SHAKE WELL BEFORE USING.

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL:

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL:

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

2

DISCLAIMER

The label instructions for the use of this product reflect the opinion of experts based on research and field use. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the use of, or application of the product contrary to label instructions, all of which are beyond the control of American Cyanamid Company. All such risks shall be assumed by the user.

American Cyanamid Company shall not be responsible for losses or damages resulting from use of this product in any manner not set forth on this label. User assumes all risks associated with the use of this product in any manner not specifically set forth on this label.

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use, subject to the risks referred to above. CYANAMID DOES NOT MAKE OR AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED AND EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

BUYER'S EXCLUSIVE REMEDY AND AMERICAN CYANAMID'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF PASSPORT. In no case shall Cyanamid or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product.

USES WITH OTHER PRODUCTS (TANK MIXES)

If this product is used in combination with any other product except as specifically recommended in writing by American Cyanamid Company then American Cyanamid Company shall have no liability for any loss, damage or injury arising out of its use in any such combination not so specifically recommended. If used in combination recommended by American Cyanamid Company, the liability of American Cyanamid Company shall in no manner extend to any damage, loss or injury not directly caused by the inclusion of the American Cyanamid Company product in such combination use, and in any event shall be limited to return of the account of the purchase price of the American Cyanamid Company product.

BEST AVAILABLE COPY

GENERAL INFORMATION

Entry into treated fields is not permitted without protective clothing until sprays have dried.

PASSPORT is effective in providing weed control in reduced tillage (fields not tilled the previous fall) as well as conventional tillage production systems. Apply PASSPORT preplant incorporated from 45 days prior to planting up to planting. Incorporate PASSPORT within 24 hours of application. After PASSPORT is applied, some susceptible weeds emerge, growth stops, and the weeds either die or are not competitive with the crop.

The weed killing activity of PASSPORT involves herbicide uptake by weed roots and rapid translocation to the growing points. Therefore, adequate soil moisture is important for optimum PASSPORT activity. When adequate soil moisture is present, PASSPORT will provide residual control of susceptible germinating weeds.

Occasional internode shortening and/or temporary yellowing of soybean plants may occur following PASSPORT applications. This will not affect soybean yields.


Replanting: If replanting is necessary in a field previously treated with PASSPORT, the field may be replanted to soybeans. Rework the soil no deeper than the treated zone. Do not apply a second treatment of PASSPORT.

MIXING INSTRUCTIONS

Fill the spray tank one-half to three-quarters full with clean water. While agitating add the required amount of product and then fill the remainder of the tank with water. Maintain agitation while spraying to ensure a uniform spray mixture.

To avoid injury to sensitive crops, spray equipment used for PASSPORT applications must be drained and thoroughly cleaned with water before being used to apply other products.

To avoid injury to sensitive crops, spray equipment used for PASSPORT applications must be drained and thoroughly cleaned with water before being used to apply other products.



HERBICIDE COMBINATIONS

このかし

PASSPORT may be applied preplant incorporated in tank mix combination with metribuzin (Sencor¹ or Lexone²) PASSPORT may also be applied preplant incorporated, followed by a preemergence application of metribuzin or linuron (Lorox² or Linex³). The tank mix or sequential application of PASSPORT and metribuzin will enhance the control of common cocklebur and common ragweed in addition to the weeds controlled by a soil application of PASSPORT. The sequential application of PASSPORT followed by linuron will enhance the control of common ragweed.

Apply PASSPORI at the broadcast rate of 2 1/2 pints per acre. Refer to the metribuzin or linuron label for specific application rates, restrictions and precautionary instructions.

Under conditions of heavy grass pressure, the addition of a grass herbicide such as $PROWL^{\bullet}$ or trifluralin may be tank-mixed with PASSPORT.

When PASSPORT is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PASSPORT cannot be mixed with any product containing a label prohibiting such mixtures.

SFRAYING INSTRUCTIONS

DO NOT apply when wind velocity is greater than 10 mph, or when spray may be carried to sensitive crops. Sensitive crops include leafy vegetables, sugarbeets, and cotton.

GROUND APPLICATIONS:

Uniformly apply with properly calibrated ground equipment in 10 to 40 gallons of water per acre. A spray pressure of 20 to 40 psi is recommended. Adjust the boom height to ensure proper coverage.

APPLICATIONS WITH LIQUID FERTILIZER:

PASSPORT can be applied to the soil in liquid fertilizers. Follow all PASSPORT label recommendations regarding incorporation, timing of application, special instructions and precautions. Apply treatments in 20 or more gallons of liquid fertilizer per acre with ground equipment. Always test the compatibility of PASSPORT with liquid fertilizer before mixing in the spray tank.

PROWL is a registered trademark of American Cyanamid Company.

¹Sencor is a trademark of Mobay Corporation.

²Lexone and Lorox are trademarks of E.I. duPont de Nemours and Company.

³Linex is a trademark of Griffin Ag Products Co., Inc.

APPLICATIONS WITH DRY BULK FERTILIZER:

10-7/15

PASSPORT may be impregnated on dry bulk fertilizers. When applied as directed, PASSPORT/dry bulk fertilizer mixtures provide weed control equal to that provided by the same rates of PASSPORT applied in water or liquid fertilizer.

Follow all PASSPORT label recommendations regarding application and incorporation, special instructions and precautions. Apply PASSPORT/dry bulk fertilizer mixtures only with ground equipment.

All individual state regulations relating to dry bulk fertilizer, registration, labeling, and application are the responsibility of the individual and/or company selling the PASSPORT/dry bulk fertilizer mixtures.

A minimum of 200 pounds and a maximum of 450 pounds of dry bulk fertilizer impregnated with the recommended amount of PASSPOKT must be applied per acre.

DO NOT impregnate PASSPORT onto coated ammonium nitrate or limestone because these materials will not absorb the herbicide. Dry fertilizer blends containing mixtures of ammonium nitrate or limestone may be impregnated with PASSPORT. A minimum of 200 pounds of impregnated dry bulk fertilizer, excluding the weight of ammonium nitrate or limestone, must be applied per acre.

Apply PASSPORT at the rate of 2 and 1/2 pints per acre. Use the following table to determine the amount of PASSPORT to be impregnated on a ton of dry bulk fertilizer based on the rate of fertilizer which will be applied per acre.

RATE CHART FOR IMPREGNATION OF DRY BULK FERTILIZER WITH PASSPORT (Fints of PASSPORT per Ton of Fertilizer)

PASSPORT Rate Per Acre	Fertilizer Rate pounds per acre	Pints per Ton
2 1/2 pints	200	25
•	250	20
	300	16 2/3
	350	14 1/3
	400	12 1/2
	450	11

For those rates not listed in this table, calculate the pints of PASSPORT to be impregnated on a ton of dry bulk fertilizer using the following formula:

		2 1/2 Pints of		
2000		PASSPORT per		Pints of
Pounds of dry	X	acre	_	PASSPORT per
fertilizer per acre		(recommended rate)		ton of fertilizer

BEST AVAILABLE COPY

To impregnate PASSPORT on bulk fertilizer, use a closed rotary-drum mixer of other commonly used dry bulk fertilizer blender equipped with suitable spray equipment. Spray nozzles must be placed to provide uniform coverage of PASSPORT onto the fertilizer during mixing.

If Treflan is to be combined with the PASSPORT prior to impregnation, premix the Treflan with an equal volume of water before adding it to the PASSPORT. DO NOT mix undiluted Treflan with PASSPORT.


Apply the PASSPORT/dry bulk fertilizer mixture with an accurately calibrated dry fertilizer spreader. The PASSPORT/dry bulk fertilizer mixture must be spread uniformly on the soil surface. Uneven spreading can cause poor weed control and crop injury.

Refer to PREPLANT INCORPORATED APPLICATIONS section of this label for incorporation directions.

PASSPORT HERBICIDE USE AREA

PASSPORT may be applied to soybeans only in the states or parts of states shaded in the map below:

(In Minnesota PASSPORT may be applied south of state highway 210. In Texas PASSPORT may be applied east of state highway 83).



Apply PASSPORT at a broadcast rate of 2 1/2 pints per acre. At this broadcast rate, one gallon of PASSPORT will treat 3.2 acres of soybeans.

(2 1/2 PINTS PER ACRE)

NOTE: Only one application of FASSPORT may be made during the season.

WEEDS CONTROLLED

When applied as directed, PASSPORT will control or reduce competition from the weeds listed below.

NOTE: C = Control, R = Reduced Competition



Weeds Controlled	Preplant Incorporated		
Anoda, spurred	С		
Buffalobur	С		
Carpetweed	С		
Cocklebur, common*	С		
Devilsclaw	C		
Calinsoga	C		
Jimsonweed	C		
Kochia	С		
Lambquarters, common	С		
Mallow, Venice	R		
Morningglory			
entireleaf	R		
ivyleaf	R		
pitted	R		
smallflower	С		
tall	R		
Mustard species	С		
Nightshade			
black	С		
Eastern black	С		
hairy	c ,/		
Pigweed			
palmer	С		
redroot	С		
smooth	С		
spiny	С		
Puncturevine	С		
Purslane, common	С		
Pusley, Florida	С		
Ragweed,			
common	R		
giant	R		
Sida, prickly (teaweed)	С		
Smartweed			
ladysthumb	С		
Pennsylvania	С		
Spurge			
prostrate	С		
spotted	С		
Sunflower, common	С		
Velvetleaf	С		
Waterhemp, tall	С		

^{*} Cultivation and/or a postemergence herbicide may be required for season-long control.

GRASS WEEDS

Weeds Controlled	Preplant Incorporated
Barnyardgrass	С
Crabgrass	
large	C
smooth	С
Crowfootgrass	С
Cupgrass, woolly	С
Foxtail	
giant	С
green	С
yellow	C
Goosegrass	С
Itchgrass	ĸ
Johnsongrass	
seedling	С
rhizome	R
Millet, wild proso	R
Panicum	
browntop	С
fall	С
Texas	C
Sandbur, field	C
Shattercane	c /
Signalgrass, broadleaf	c /
Sorghum almum	c
Witchgrass	С
SEDGES	
Nutsedge	
yellow	R
purple	R
harbre	18



SOIL APPLICATION INSTRUCTIONS

PASSPORT may be soil applied and incorporated up to 45 days (early preplant) prior to soybean planting. DO NOT apply after planting soybeans.

PASSPORT controls weeds by uptake by weed roots and translocation to the growing points where it stops weed growth.

Adequate soil moisture is required for optimum activity. The amount of rainfall or irrigation required following application depends on existing soil moisture, soil texture and organic matter content. Sufficient water to moisten the soil to a depth of 2 inches is normally adequate. If dry soil conditions persist following a PASSPORT application, a cultivation is recommended to control escaped weeds. When adequate moisture is received after dry conditions, PASSPORT will provide residual control of susceptible germinating weeds; activity on established weeds will depend on the weed species and the location of its root system in the soil.

PREPLANT INCORPORATED APPLICATIONS

PASSPORT may be applied either prior to spring tillage, or following land preparation and should be thoroughly incorporated to a depth of 2 to 3 inches. When applied to beds, maintain PASSPORT in the surface 1-2 inches of the finished beds. Application may be made up to 45 days prior to planting (early preplant). Incorporate prior to soybean planting and within 24 hours of application. Mechanical incorporation can be achieved by the following:

- (a) Disk harrow set to cut 3 to 4 inches deep and operated at 4 to 6 mph with the second pass at an angle to the first.
- (b) PTO-driven equipment (tillers, cultivators, hoes) set to cut 2-3 inches deep and operated one time at 4 mph or less.
- (c) Rolling cultivator set to cut 2 to 4 inches deep and operated two times at 6 to 8 mph. Use only on coarse textured soils.
- (d) Field cultivator with 3 to 4 rows of sweeps (Do not use chisel points), spaced at intervals of 7 inches or less and staggered so that no soil is left unturned, set to cut 3 inches deep, and operated two times at more than 5 mph with the second pass made at an angle to the first.
- (e) LELY-Roterra⁴ set and operated according to the manufacture c's directions to thoroughly incorporate PASSPORT to a depth of 2 to 3 inches.
- (f) Do-All set and operated according to the manufacturer's directions to thoroughly incorporate PASSPORT to a depth of 2 to 3 inches.

If soybeans are planted on beds, apply and incorporate after bed formation using PTO-driven equipment or a rolling cultivator. For optimum weed control, FASSPORT should be maintained in the surface 1-2 inches of the finished bed.

Ì

)

⁴LELY-Roterra is a trademark of Lely.

ROTATIONAL CROPS

The following rotational crops may be planted after applying PASSPORT at the recommended rate in soybeans:

1. Anytime
Lima beans
Southern peas
Soybeans
Peanuts

Four months after PASSPORT application:
 Wheat
 Edible beans and peas (other than lima beans and Southern peas)

3. Nine and one-half months after PASSPORT application:

Barley Field corn Field corn grown for seed* Rye Tobacco

4. Eighteen months after PASSPORT application:

Alfalfa
Cotton
Lettuce
Oats
Popcorn
Safflower
Sorghum
Sweet corn

)

5. Twenty six months after PASSPORT a_P plication. All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.

 Forty months after PASSPORT application. Sugarbeets

*Several seed companies have tested a wide range of inbreds for sensitivity to PASSPORT soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PASSPORT the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PASSPORT shall be assumed by the user.

Use of PASSPORT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

EAR

If soybeans are furrow irrigated, till the soil prior to planting winter wheat or barley. The beds should be broken up and the soil mixed with tillage equipment set to cut 4-6 inches deep.

Only rotational crops harvested at maturity may be used for feed or food.

In the event of a crop loss due to weather, soybeans can be replanted. DO NOT work the soil deeper than 2 inches.

There should be an interval of at least 85 days between an application of PASSPORT and soybean harvest.

DO NOT graze or feed treated soybean forage, hay or straw to livestock.

DO NOT apply products containing chlorimuron ethyl (Classic⁵, Canopy⁵, Gemini⁵, Lorox Plus⁵, PREVIEW⁵, etc.) or imazaquin (SCEPTER[®], SQUADRON[®], or TRI-SCEPT[®], SCEPTER[®] O.T.[®], SCEPTER[®] 70DG,) or products containing imazethapyr (PURSUIT[®], or PURSUIT[®] PLUS) the same year as PASSPORT or injury to follow crops may occur.

)

For additional information regarding the use of PASSPORT herbicide, call telephone no. 800-942-0500.

American Cyanamid Company
Agricultural Division
Crop Protection Chemicals Department
Wayne, NJ 07470 •1992

⁵Classic, Canopy, Preview, Gemini and Lorox Plus are trademarks of E.I. duPont de Nemours and Company.

*Registered Trademarks of American Cyanamid Company.

Proposed PASSPORT[®] Label Changes EPA Reg. No. 241-325

The following changes are from the approved PASSPORT label dated 7/91 (approved by the EPA on 3/13/92) to the proposed PASSPORT label dated 7/92. The changes are listed below and highlight on the labels.

Froposed Label (page no.)	Approved Label (page no.)	Nature of change
1	1	Under ingredients, the statement "Contains petroleum distillates" has been added to comply with labeling regulations.
		The label date has been changed to 7/92.
3	3	The disclaimer under USES WITH OTHER PRODUCTS (TANK MIXES) has been modified to allow users to use tankmixes not specifically recommended by American Cyanamid, at their own risk.
7	7	New York state has been removed from the USE AFTA map.
8	8	The name of pusley weed has been corrected to read Florida pusley.
9	9	Foxtail species (giant green, robust purple and robust white) have been deleted.
		Sorghum almum has been added to GRASS WEEDS controlled.
11	11	A caution has been added to the incorporation, instructions in the first paragraph.

^{*}Registered trademark of American Cyanamid Company



Barley has been moved from the four month to the nine and one-half month interval. Rye has been placed under the nine and one-half month category (all cereals were previously approved at the four month restriction interval).

Specific crops are listed as safe at the eighteen month interval. (Previously all crops not otherwise listed fell under this timing).

All other crops not listed specifically under other plant-back intervals now restricted to twenty six months.

Sugarbeets are restricted to 40 months following a PASSPORT application to afford additional protection to sugarbeet growers. (The forty month restriction was previously approved by the EPA on the Supplemental Label for PURSUIT use in North Dakota and No. Minnesota - see Exhibit 1). This label change makes the restriction consistent on PURSUIT containing products.