

NOV 15 1993

Mrs. Desiree L. Little
American Cyanamid Co.
P.O. Box 400
Princeton, NJ 08543

Dear Mrs. Little:

Subject: Minor Label changes- Worker Protection
SQUADRON® Herbicide
EPA Reg. No. 241-327
Your application dated Oct. 27, 1993

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

Sincerely yours,

Robert J. Taylor
Product Manager 25
Fungicide-Herbicide Branch
Registration Division (H7505C)

Enclosure

BEST AVAILABLE COPY

SQUADRON®
herbicide
FOR USE IN SOYBEANS

ACTIVE INGREDIENTS:

Ammonium salt of imazaquin (2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-quinolinecarboxylic acid)*.....	3.65%
pendimethalin (N-1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine.....	21.96%
INERT INGREDIENTS**.....	74.39%
TOTAL.....	100.00%

*Equivalent to 3.46%, 2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-quinolinecarboxylic acid.

**Contains Petroleum Distillates

BEST AVAILABLE COPY

SQUADRON contains 2.33 pounds of active ingredients per gallon (2.00 pounds ai of pendimethalin and 0.33 pounds ai of imazaquin).

EPA Reg. No. 241-327

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN
DANGER!/PELIGRO!

ACCEPTED
with COMMENTS
in EPA Letter Dated:

NOV 15 1993

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

Label, pesticide, Fungicide, and Herbicide, as amended, for the year registered under EPA Reg. No.

241-327

In case of an emergency endangering life or property involving this product, call collect, day or night, Area Code 201-835-3100.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES; hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Call a physician.

IF SWALLOWED; DO NOT induce vomiting. Call a physician or Poison Control Center immediately.

IF ON SKIN; wash with plenty of soap and water. Get medical attention if skin irritation persists.

IF INVALED; Remove victim to fresh air. If not breathing, give artificial respiration. Get medical attention.

NOTE TO PHYSICIAN: Because of increased risk of chemical pneumonia or pulmonary edema caused by aspiration of the hydrocarbon solvent, vomiting should be induced only under professional supervision.

See Additional Precautionary Statements Inside.

AMERICAN CYANAMID COMPANY
AGRICULTURAL PRODUCTS DIVISION
CROP PROTECTION CHEMICALS DEPARTMENT
WAYNE, NJ 07470 ©1993

Net Contents:

10/93

®Registered Trademark of American Cyanamid Company

(PRECAUTIONARY STATEMENTS (3 7 25

HAZARDS TO HUMANS

DANGER

Causes irreversible eye damage. Harmful if inhaled or absorbed through the skin. DO NOT get in eyes or on clothing. Avoid breathing vapor or spray mist. Avoid contact with skin. Use with adequate ventilation.

Personal Protective Equipment (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions of category F on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- * Long-sleeved shirt and long pants
- * Chemical-resistant gloves, such as Barrier Laminate or Butyl Rubber ≥ 14 mils or Nitrile Rubber ≥ 14 mils or Viton ≥ 14 mils
- * Shoes plus socks
- * Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. DO NOT reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:

Users should:

- * Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- * Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. DO NOT apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. DO NOT contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of pesticide application.

BEST AVAILABLE COPY

Observe all cautions [limitations in this leaflet [on the labels of products used in combination with SQUADRON. DO NOT use SQUADRON other than in accordance with the instructions set forth on this label. The use of SQUADRON not consistent with this label can result in injury to crops, animals, or persons. Keep container closed to avoid spills and contamination.

DO NOT apply this product through any type of irrigation system.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: if the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- * Coveralls
- * Chemical-resistant gloves, such as Barrier Laminate or Butyl Rubber ≥14 mils or Nitrile Rubber ≥14 mils or Viton ≥14 mils.
- * Shoes plus socks
- * Protective eyewear

STORAGE AND DISPOSAL

STORAGE: KEEP FROM FREEZ'NG. DO NOT STORE BELOW 40°F. SHAKE WELL BEFORE USING.

DO NOT contaminate water, food, or feed by storage or disposal.

BEST AVAILABLE COPY

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL FOR 2.5 GALLONS:

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONTAINER DISPOSAL FOR MINI BULK:

If program offered by dealer, return to point of purchase for repackaging. If no repackaging program is available, triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.


DISCLAIMER

The label instructions for the use of this product reflect the opinion of experts based on research and field use. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, herbicide resistant weed populations, or the use of, or application of the product contrary to label instructions, all of which are beyond the control of American Cyanamid Company. All such risks shall be assumed by the user.

American Cyanamid Company shall not be responsible for losses or damages resulting from use of this product in any manner not set forth on this label. User assumes all risks associated with the use of this product in any manner not specifically set forth on this label.

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use, subject to the risks referred to above. CYANAMID DOES NOT MAKE OR AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED AND EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

BUYER'S EXCLUSIVE REMEDY AND AMERICAN CYANAMID'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF SQUADRON....In no case shall Cyanamid or the seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product.



If this product is used in combination with any other product except as specifically recommended in writing by AMERICAN CYANAMID COMPANY, then AMERICAN CYANAMID COMPANY shall have no liability for any loss, damage or injury arising out of its use in any such combination not so specifically recommended. If used in combination recommended by AMERICAN CYANAMID COMPANY, the liability of AMERICAN CYANAMID COMPANY shall in no manner extend to any damage, loss or injury not directly caused by the inclusion of the AMERICAN CYANAMID COMPANY product in such combination use, and in any event shall be limited to return of the amount of the purchase price of the product.

GENERAL INFORMATION

For broad-spectrum control of grass and broadleaf weeds in soybeans, SQUADRON may be applied as a preplant incorporated or preemergence treatment.

After SQUADRON is applied, some susceptible weeds emerge, growth stops, and then the weeds either die or are not competitive with the crop.

A timely cultivation may aid in the control of certain weeds or improve general weed control when adequate moisture is not received after application. Cultivation should be shallow.

SQUADRON reaches the growing points of susceptible weeds either by direct contact in the soil, or by root uptake and rapid translocation to the growing points. Therefore, adequate soil moisture is important for optimum SQUADRON activity. The amount of rainfall or irrigation required following application depends on existing soil moisture, soil texture and organic matter content. Sufficient water to moisten the soil to a depth of 2 inches is normally adequate. If adequate moisture is not received within 7 days after treatment, then a cultivation or postemergence herbicide application may be needed to improve weed control. When adequate moisture is received after dry conditions, SQUADRON will provide residual control of susceptible germinating weeds; activity on established weeds will depend on the weed species and the location of its root system in the soil.

A SQUADRON soil treatment followed by a SCEPTER[®], SCEPTER[®] 70DG, SCEPTER[®] O.T.[®] or Classic¹ postemergence treatment will control certain problem weeds. Use sequential treatments in recommended states only.

Occasionally, internode shortening of soybean plants may be observed with SQUADRON applications. This has no effect on soybean yields.

[®]Registered Trademark of American Cyanamid Company

¹Registered Trademark of E.I. du Pont de Nemours and Company

BEST AVAILABLE COPY

7 9 25

Within any weed population, there may be the potential for the occurrence of weeds resistant to herbicides when compounds that affect the same site of action are used continuously on the same acre. Therefore, any multiple year weed management program should include, when local agronomic practices allow, as many of the following practices as possible:

- use scouting and weed thresholds to determine the need for herbicide treatment
- utilize tank-mixes or sequential applications of different mode-of-action herbicides to control the same target weeds
- combine cultivation with herbicide treatments

See your Cyanamid representative for addition information.

PRECAUTIONS:

DO NOT apply SQUADRON postemergence to soybeans as crop injury may occur.

DO NOT APPLY SQUADRON OR ITS TANK MIXTURES PREEMERGENCE IN SOUTH DAKOTA.

DO NOT use SQUADRON other than in accordance with the instructions set forth on this label.

DO NOT use SQUADRON in any tank mixes other than those specifically provided in this label.

DO NOT use on crops other than soybeans. Crops other than soybeans, such as cotton, corn, grain sorghum, rice and vegetables, may be injured by spray drift or other indirect contact with SQUADRON.

To avoid injury to sensitive crops from spray drift, follow all use directions and precautions in SPRAYING INSTRUCTIONS on page 15 and 16.

To avoid injury to sensitive crops, spray equipment used for SQUADRON applications must be drained and thoroughly cleaned with water before being used to apply other products to these crops.

Apply SQUADRON prior to July 1 in Use Region 3, as defined in the USE AREA section of this label.

There should be an interval of 90 days between the SQUADRON application and soybean harvest.


DO NOT graze or feed treated soybean forage, hay, or straw to livestock.

Use of SQUADRON herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations, however, various environmental and agronomic factors make it impossible to eliminate all risks associated with use of this product and, therefore, rotational crop injury is always possible. See pages 20 to 22 for ROTATIONAL CROP RESTRICTIONS.

Replanting: If replanting is necessary in a field previously treated with SQUADRON, the field may be replanted to soybeans. Rework the soil no deeper than the treated zone. DO NOT APPLY A SECOND TREATMENT OF SQUADRON.

USE AREA

SQUADRON can be applied only in the states or parts of states shaded in the following map:



Use Region 1 includes eastern Oklahoma, Arkansas, the Missouri bootheel, Tennessee, North Carolina, South Carolina, Georgia, Florida, Alabama, Mississippi, Louisiana, and eastern Texas.

Use Region 2 includes eastern Kansas (east of U.S. 81; the counties of Cloud, Ellsworth, Harvey, Jewell, Lincoln, Mitchell, McPherson, Ottawa, Republic, Saline, Sedgewick, and Sumner), southeastern Nebraska (east of U.S. 81, south of U.S. 34), Missouri, Illinois (south of S.R. 116 west of Peoria; south of U.S. 24 east of Peoria), Indiana, Ohio, Michigan, Kentucky, Virginia, West Virginia, Pennsylvania, Maryland, Delaware, and New Jersey.

Use Region 3 includes Nebraska (east of U.S. 81, north of U.S. 34 and also that area east of U.S. 283, south of U.S. 30, and west of U.S. 81), South Dakota (east of U.S. 81), Illinois (north of S.R. 116 west of Peoria; north of U.S. 24 east of Peoria), Wisconsin, Iowa, and southern Minnesota.

Note: See USE RATES and ROTATIONAL CROP RESTRICTIONS sections for recommendations applying to each Use Region.

BEST AVAILABLE COPY

WEEDS CONTROLLED

The following grass and broadleaf weeds are controlled by recommended treatments of SQUADRON:


WEED	LEVEL OF CONTROL
GRASSES	
Barnyardgrass	Control
Corn, volunteer	Suppression
Crabgrass	Control
Crowfootgrass	Control
Cupgrass, Woolly	Control ¹
Foxtail	
Giant	Control
Green	Control
Robust	Control
Yellow	Control
Goosegrass	Control
Johnsongrass, seedling	Control
Panicum	
Browntop	Control
Fall	Control
Texas	Control
Sandbur, Field	Control
Shattercane	Control ¹
Signalgrass, Broadleaf	Control
Witchgrass	Control
BROADLEAF WEEDS	
Alligatorweed	Control
Beggarweed, Florida	Control ²
Bristly Starbur	Control
Burcucumber	Control ¹
Carpetweed	Control
Cocklebur, Common	Control
Copperleaf, hopcornbeam	Control ³
Jimsonweed	Control
Kochia	Control
Lambsquarters, Common	Control
Mallow, Venice	Control
Mexicanweed	Control ²
Morningglory	
Entireleaf	Control ⁴
Ivyleaf	Control ⁴
Palm Leaf	Control
Pitted	Control
Smallflower	Control
Tall	Control ⁴
Mustard Species	Control
Nightshade, Eastern Black	Control

4An application of SQUADRON will provide suppression of these weeds. For control of these morningglory species, the sequential program in Footnote #3 should be used.

5Some broadleaf weeds such as giant ragweed germinating deep in the soil or with subsequent flushes may require a cultivation or postemergence herbicide application for season long control.

6Only light to moderate infestations of sicklepod are controlled. Later sicklepod flushes may require a cultivation or a postemergence herbicide application. For heavy sicklepod infestations, the sequential program in footnote #2 may be used or Classic may be applied postemergence at 1/2 to 3/4 ounces per acre following a soil application of SQUADRON. Apply the higher rate of Classic if sicklepod seedlings are taller than 2 inches. A cultivation 14 days after Classic application may be required to control sicklepod escapes. Refer to the Rotational Crop Restrictions section of this label for instructions on planting follow crops when using this program. The Classic sequential program may be used only in Use Region 1 (except Oklahoma).

7A preplant incorporated application is recommended for velvetleaf control. A preemergence application of SQUADRON will aid in the control and reduce competition from velvetleaf.



DIRECTIONS FOR CONVENTIONAL, MINIMUM, AND NO-TILL APPLICATIONS

APPLICATION RATE

Apply SQUADRON at a rate of 3.0 pints per acre.

APPLICATION INSTRUCTIONS

SOIL APPLICATIONS

SQUADRON may be applied in conventional, minimum, or no-till as a preplant incorporated or preemergence application up to 45 days (30 days in Use Region 1) before planting. Preemergence applications may also be applied during or after planting but before the crop emerges. DO NOT apply postemergence (or at cracking) to soybeans.

Adequate moisture is required for activation of SQUADRON.

PREPLANT INCORPORATED APPLICATIONS

Apply SQUADRON before planting and incorporate uniformly into the top 1 to 2 inches of soil.

SQUADRON may be applied immediately before planting or up to 30 days prior to planting in Use Region 1. For Use Region 2 and Use Region 3, SQUADRON may be applied up to 45 days prior to planting. Incorporate within 7 days after application.

If soybeans are planted on beds, apply and incorporate after bed formation, using PTO-driven equipment or rolling cultivator.

PREEMERGENCE APPLICATIONS (including no-till)

Apply SQUADRON up to 45 days (30 days in USE REGION 1) before, during or after planting but before the crop emerges. As with other herbicides applied preemergence, rainfall or irrigation is necessary to activate SQUADRON in the soil. If sufficient rainfall or irrigation to activate SQUADRON is not received within 7 days after application, a thorough shallow tillage, cultivation, or postemergence herbicide treatment (as appropriate to the tillage system) may be required for control of emerged weeds.

HERBICIDE COMBINATIONS

For no-till uses, SQUADRON may be tank mixed with Gramoxone Extra², Roundup³, 2,4-D or 2,4-DB to kill existing vegetation. Use a minimum of 10 gallons of water per acre. Use higher gallonage for fields with dense vegetation or heavy crop residues. Use ground equipment only. Plant soybeans at least one inch deep and adjust planters to ensure adequate seed coverage.

²Registered Trademark of Zeneca Inc.

³Registered Trademark of Monsanto Agricultural Products Co.

BEST AVAILABLE COPY

When SQUADRON is tank mixed with Gramoxone Extra or Roundup, add SQUADRON to the tank first, next add PROWL 3.3EC (if used), then add Gramoxone Extra or Roundup. When Gramoxone Extra is used in a tank mixture, add 8 oz. of non-ionic surfactant per 100 gallons of spray mixture as the last ingredient in the tank.

SQUADRON should be used only in accordance with the directions on this label. Do not use SQUADRON in any tank mixes other than those specifically provided in this label. Always follow the more restrictive label limitations and precautions.

Soil Texture

The application rate of some herbicides which may be tank mixed with SQUADRON will vary with soil texture. Combination rate tables in this label refer to three soil texture groups: COARSE, MEDIUM, and FINE. The following table lists soil textures included in each of these three groups:

<u>COARSE</u>	<u>MEDIUM</u>	<u>FINE</u>
sands	sandy clay loams*	silty clay loams*
loamy sands	sandy clays	silty clays
sandy loams	loams	clay loams
	silt loams	clays
	silts	

*Sometimes considered transitional soils.

SQUADRON plus Additional PROWL 3.3EC herbicide Tank Mixture

PROWL 3.3EC can be tank mixed with SQUADRON and applied preplant incorporated or preemergence if heavy or difficult to control infestations are anticipated from grasses listed in this leaflet. SQUADRON plus PROWL 3.3EC must be applied preplant incorporated if shattercane or woolly cupgrass infestations are anticipated.

On coarse textured soils add PROWL 3.3EC at 0.6 pint per acre* and on medium and fine textured soils add PROWL 3.3EC at 0.6 to 1.2 pints per acre*.

See the PROWL 3.3EC label for soil types classified as coarse, medium and fine textured.

Apply this tank mixture preplant incorporated north of Interstate 80.

Observe all precautions and limitations on the PROWL 3.3EC label.

*PROWL 3.3EC rates are based on 3.3 pounds of active ingredient per gallon. If the 4.0 pounds of active ingredient per gallon formulation is used, multiply the above rates by 0.82.

BEST AVAILABLE COPY

SQUADRON plus Command 4EC Tank Mixture

Observe all precautionary statements in the Command⁴ 4EC label before using.

SQUADRON plus Command 4EC preplant incorporated tank mixture may be used when heavy infestations of velvetleaf are anticipated. Uniformly apply the tank mixture with properly calibrated ground equipment in 10 to 40 gallons of water or 20 or more gallons of liquid fertilizer per acre. Use sprayers equipped with nozzles that provide accurate and uniform application.

NOTE: The use of an agriculturally approved drift reducing agent is required at spray volumes of 10 to 15 gallons per acre.

Apply 3 pints per acre of SQUADRON plus 0.25 to 0.5 pint (4 to 8 oz.) per acre of Command 4EC to the soil surface and uniformly incorporate. Refer to the Command 4EC label for incorporation requirements. DO NOT apply aerially or through irrigation systems. DO NOT apply this tank mixture to overly moist or wet soils.

Command 4EC is a volatile compound. Off-site movement of spray drift or vapors of Command 4EC can cause foliar whitening or yellowing of some plants. Prior to using Command 4EC, read and strictly follow all precautions and application instructions on the Command 4EC label.

When applied as directed, tank mixtures of SQUADRON plus Command 4EC control heavy infestations of velvetleaf as well as the weeds listed in the WEEDS CONTROLLED section of this label.

Use the high rate of Command 4EC when severe infestations of velvetleaf are anticipated.

Follow all rotational crop restrictions on the SQUADRON and Command 4EC labels. Always follow the most restrictive label.

In the event of a crop loss due to weather conditions, soybeans can be replanted. DO NOT work the soil deeper than 2 inches.

SQUADRON Plus Metribuzin Tank Mixtures

Before using, observe all precautionary statements in the SQUADRON, Lexone⁵, and Sencor⁶ leaflet labels. SQUADRON may be applied in tank mix combination with Lexone or Sencor either preplant incorporated or preemergence, or SQUADRON may be applied preplant incorporated followed by a preemergence application of Lexone or Sencor. For preemergence applications, apply during or immediately after planting. These treatments will control Florida beggarweed, hemp sesbania, hophornbeam copperleaf and sicklepod in addition to the weeds controlled by a soil application of SQUADRON alone.

⁴Registered Trademark of FMC Corporation
⁵Registered Trademark of E.I. duPont de Nemours and Company
⁶Registered Trademark of Miles Company

BEST AVAILABLE COPY

Apply Lexone or Sencor at the broadcast rate per acre specified in the table below, in tank mix combination with 3 pints per acre of SQUADRON.

	<u>Lexone or Sencor Rate*</u>	
	4.0L	75%DF
Coarse Soils	1/2 pint	1/3 lb.
Medium Soils	1/2 pint	1/3 lb.
Fine Soils	2/3 pint	1/2 lb.

* DO NOT use Lexone or Sencor on coarse soils of less than 2% organic matter. Use 1/2 lb. or 2/3 pint of Lexone or Sencor on medium soils of greater than 4% organic matter.

Tank Mixtures with 2,4-D

2,4-D may be used with SQUADRON alone or in combination with other SQUADRON tank mixes prior to planting for control of some annual broadleaf weeds. Use the following rates of 2,4-D in tank mixtures with SQUADRON:

<u>2,4-D Formulation</u>	<u>Rate(lb a.i./A)</u>	<u>Minimum Days Before Planting</u>
Ester	0.5	7
Amine	0.5	15
Ester or Amine	1.0	30

Refer to 2,4-D label for weeds controlled. Observe all cautions and limitations on the 2,4-D label.

Tank Mixtures with 2,4-DB

2,4-DB may be used with SQUADRON alone or in combination with other SQUADRON tank mixes for control of some annual broadleaf weeds. Refer to the 2,4-DB label for specific use recommendations, rates and weeds controlled.

Observe all cautions and limitations on the 2,4-DB label.

Tank Mixtures with Roundup

Roundup may be used with SQUADRON alone or in combination with other tank mixes for the control of certain emerged grasses and broadleaf weeds. Refer to the Roundup label for specific use recommendations, rates and weeds controlled.

NOTE: The Roundup rate for tank mixes with SQUADRON is the same as the rate for Roundup used alone.

Observe all cautions and limitations on the Roundup label.

BEST AVAILABLE COPY

Tank Mixtures with Gramoxone Extra

Apply 1-1/2 to 2-1/2 pints of Gramoxone Extra per acre in SQUADRON tank mixtures. Use the 2-1/2 pint rate if weeds are 4 to 6 inches tall. Weeds over 6 inches may not be controlled with this treatment. Apply up to 14 days before, during or immediately after planting.

When Gramoxone Extra is included in a tank mixture, add a nonionic spreader surfactant at a rate of 8 fl. oz. per 100 gallons of spray mixture as the last ingredient in the tank.

Gramoxone Extra will control most annual emerged weeds and suppress many emerged perennials. Refer to the Gramoxone Extra label for specific use recommendations and weeds controlled.

Observe all precautions and limitations on Gramoxone Extra label.

MIXING INSTRUCTIONS

Fill the spray tank one-fourth to one-half full with clean water or liquid fertilizer. While agitating add the required amount of product and then fill the remainder of the tank with water or liquid fertilizer. Maintain agitation while spraying to ensure a uniform spray mixture.

When tank mixing SQUADRON with recommended herbicides, add the SQUADRON to the spray tank first and make sure it is thoroughly mixed before adding the other herbicides.

For mixing instructions for tank mixtures with Roundup or Gramoxone Extra see the NO-TILL SYSTEMS section of this leaflet.

SPRAYING INSTRUCTIONS


NOTE: DO NOT make applications when spray may be carried by wind to sensitive crops.

Avoid overlaps when spraying.

GROUND APPLICATIONS:

Uniformly apply with properly calibrated ground equipment in 10 to 40 gallons of water, or 20 or more gallons of liquid fertilizer per acre. Use higher gallonage for fields with dense vegetation or heavy crop residues. A spray pressure of 20 to 40 psi is recommended.

DO NOT apply with ground equipment when wind velocity is greater than 20 mph.




AERIAL APPLICATIONS:

Uniformly apply with properly calibrated aerial equipment in 5 or more gallons of water per acre.

To avoid injury to sensitive crops from drift, aerial application must adhere to the following SPECIAL AERIAL USE DIRECTIONS AND PRECAUTIONS:

- Use nozzles which produce a coarse spray.
- Nozzle height above ground must be a maximum of 10 feet.
- Nozzles must be pointed toward the rear of the aircraft. The downward angle of the nozzle should not be greater than 20 degrees.
- To minimize wing-tip vortex roll, nozzles or spray boom must not be located any closer to end of wing or rotor than three-fourths the distance from the center of the aircraft.
- Use a maximum spray pressure of 40 psi.
- A buffer zone must be established between the area to be sprayed and sensitive crops.
- DO NOT spray when wind velocity is greater than 5 mph.

Applicator is responsible for any loss or damage which results from spraying SQUADRON in a manner other than recommended in this label. In addition, applicator must follow all applicable state and local regulations and ordinances in regard to spraying.



APPLICATIONS WITH LIQUID FERTILIZERS

SQUADRON can be applied with liquid fertilizers, alone or in combination with PROWL 3.3EC or Command 4EC. Follow all SQUADRON label recommendations regarding incorporation, timing of application, special instructions and precautions. Apply treatments in 20 or more gallons of liquid fertilizer per acre with ground equipment. Maintain continuous agitation in the spray tank to prevent separation. Use only flood nozzles with no nozzle screens to prevent clogging.

All individual state regulations relating to fluid fertilizer mixing, registration, labeling and application are the responsibility of the individual and/or company selling the SQUADRON/liquid fertilizer mixture.

LIQUID FERTILIZER COMPATIBILITY DETERMINATIONS

If a liquid fertilizer and herbicide(s) mixture separates in the spray tank, clogged equipment and uneven application can result, which can cause poor weed control and crop injury. Always predetermine the compatibility of SQUADRON alone or with PROWL 3.3EC or Command 4EC in the specific liquid fertilizer to be used according to the following directions:

1. Add 1 pint of fertilizer to each of 2 one-quart jars.
2. Add 1/2 teaspoon of adjuvant* to one jar.
3. (a) When using SQUADRON alone, add to each jar the correct amount of SQUADRON as specified in the table below.
(b) When using SQUADRON plus PROWL 3.3EC or Command 4EC, first add the specified quantity of SQUADRON, then add the correct amount of PROWL 3.3EC or Command 4EC.
4. Close both jars and shake thoroughly for 10 seconds. Let them stand for 30 minutes and then observe the results. Look for signs of separation, an oily layer or globules, sludge, flakes or other precipitates.
5. Determine compatibility:
 - (a) If the mixture without adjuvant does not separate, use this mixture in your spray tank.
 - (b) If the mixture with adjuvant does not separate, but the one without adjuvant separates, use the adjuvant mixture in your spray tank. Add the adjuvant to the liquid fertilizer as directed on the manufacturer's label.
 - (c) If either mixture separates, but mixes readily with shaking, the mixture can be used providing good agitation is maintained in the spray tank.
 - (d) If separation of the mixture occurs, and agitation and/or adjuvant does not correct the problem, DO NOT use the herbicide(s) in that specific liquid fertilizer.

*Complex-Kalo Laboratories Inc, Kansas City MO; Sponto 168 D-Witco Chemical Co., Houston, TX; Unite-Hopkins Agricultural Chemical Co., Madison, WI; or other comparable adjuvants.

RATE CHART FOR IMPREGNATION OF DRY BULK FERTILIZER WITH SQUADRON

(Pints of SQUADRON per Ton of Fertilizer)

SQUADRON Rate Per Acre	Fertilizer Rate lbs/acre	Pints Per Ton
3 Pints	200	30
	250	24
	300	20
	350	17
	400	15
	450	13 1/3

For those rates not listed in this table, calculate the pints of SQUADRON to be impregnated on a ton of dry bulk fertilizer using the following formula:

$$\frac{2000}{\text{Pounds of dry fertilizer per acre}} \times \text{3 pints of SQUADRON per acre (recommended rate)} = \text{Pints of SQUADRON per ton of fertilizer}$$

To impregnate SQUADRON on bulk fertilizer, use a closed rotary-drum mixer or other commonly used dry bulk fertilizer blender equipped with suitable spray equipment. Spray nozzles must be placed to provide uniform coverage of SQUADRON onto the fertilizer during mixing.

If PROWL 3.3EC is to be combined with the SQUADRON prior to impregnation, premix the PROWL 3.3EC with an equal volume of water before adding it to the SQUADRON. DO NOT mix undiluted PROWL 3.3EC with SQUADRON.

Apply the SQUADRON/dry bulk fertilizer mixture with an accurately calibrated dry fertilizer spreader. The SQUADRON/dry bulk fertilizer mixture must be spread uniformly on the soil surface. Uneven spreading can cause poor weed control and crop injury.



21725

(ROTATIONAL CROP RESTRICTIONS)

The following rotational crops may be planted after applying SQUADRON in soybeans:

<u>CROP</u>	<u>USE REGION 1</u>	<u>USE REGION 2</u> (except Michigan ³)	<u>USE REGION 3</u>
Soybeans	No restrictions	No restrictions	No restrictions
Wheat	4 months	4 months ⁴	See FOOTNOTE 5 & 6
Rice	Spring following SQUADRON application	Spring following SQUADRON application ⁴	---
Barley	11 months	11 months ⁴	See FOOTNOTE 5
Field Corn	11 months ¹	11 months ^{2,4}	See FOOTNOTE 5 & 6
Edible Beans	11 months	11 months ⁴	11 months
Grain Sorghum	11 months	11 months ⁴	11 months
Oats	11 months	11 months ⁴	See FOOTNOTE 5
Peanuts	11 months	11 months ⁴	11 months
Tobacco	11 months	11 months ⁴	11 months
Sugar Beets	26 months	26 months	26 months
Other Crops	18 months	18 months	See FOOTNOTE 7

¹ For USE REGION 1 as defined by the USE AREA section of this label; field corn may be planted in the spring of the year following SQUADRON application unless extreme drought conditions develop (less than 15 inches of rainfall or irrigation is received within 6 months following the date of application).

² For USE REGION 2 (except Michigan³) as defined in the USE AREA section of this label, field corn may be planted as a rotation² crop in the spring of the year following SQUADRON application unless extreme drought conditions develop (less than 15 inches of rainfall or irrigation is received from two weeks prior to the date of SQUADRON application through November 15, of the same year).

If the minimum rainfall requirement is not met, only field corn hybrids which possess tolerance or resistance to SQUADRON and other imidazolinone herbicides may be planted the spring of the year following a SQUADRON application (SEE NOTE).

BEST AVAILABLE COPY

- 32 7 25
- 3 Growers in the counties of Berrien, Cass, St. Joseph, Branch, Hillsdale, Lenawee, Monroe, Van Buren, Kalamazoo, Calhoun, Jackson, Washtenaw, and Wayne may use the rotational crop restrictions for Use Region 2.

Growers in other Michigan counties may NOT plant oats or barley in the fall or spring of the year following a SQUADRON application. In this geography, only field corn hybrids which possess tolerance or resistance to SQUADRON and other imidazolinone herbicides may be planted the spring of the year following an application of SQUADRON (SEE NOTE).

- 4 ROTATIONAL CROP RESTRICTIONS for a sequential application of SCEPTER O.T. following a soil application of SQUADRON (Use Region 2 only):

For Use Region 2 as defined in the USE AREA section of this label; soybeans may be planted anytime. Barley, edible beans, grain sorghum, oats, peanuts, rice, tobacco, and wheat may be planted 15 months after the last herbicide application. Cotton may be planted 18 months after the last herbicide application.

Only field corn hybrids which possess tolerance or resistance to SQUADRON and other imidazolinone herbicides may be planted the spring of the year following a sequential application (SEE NOTE). Other field corn varieties may be planted 15 months after the last herbicide application.

- 5 For USE REGION 3 as defined in the USE AREA section of this label, DO NOT plant field corn, wheat, barley, or oats as rotational crops in the fall or in the spring of the year following SQUADRON application.

- 6 In Nebraska, east of U.S. 283, south of U.S. 30, and west of U.S. 81; wheat may be planted 4 months after a SQUADRON application. In this geography, only field corn hybrids which possess tolerance or resistance to SQUADRON and other imidazolinone herbicides may be planted the spring of the year following an application of SQUADRON (SEE NOTE).

- 7 For USE REGION 3 as defined in the USE AREA section of this label; canola, table beets, strawberries, cabbage, tomatoes, potatoes, carrots, celery, cole crops, garlic, onions, spinach, asparagus, cauliflower, and broccoli may be planted 26 months after a SQUADRON application. Other crops may be planted 18 months after a SQUADRON application.

ROTATIONAL CROP RESTRICTIONS for sequential Classic application following SQUADRON soil treatments:

Soybeans may be planted anytime. Barley, edible beans, field corn, grain sorghum, oats, peanuts, rice, tobacco, and wheat may be planted 15 months after the last herbicide application. Cotton may be planted 18 months after the last herbicide application. Refer to rotational crop restrictions listed in the Classic label. Always follow the more restrictive label.

NOTE: Contact your chemical dealer, seed supplier, or American Cyanamid to obtain information regarding the availability of imidazolinone tolerant field corn hybrids which are adapted to your area.

BEST AVAILABLE COPY

Use of SQUADRON herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with use of this product and, therefore, rotational crop injury is always possible.

Only rotational crops harvested at maturity may be used for feed or food.

Except where stated on a Supplemental Label or in this label, DO NOT apply products containing chlorimuron ethyl (e.g. Classic, Canopy⁷, Gemini⁷, Lorox Plus⁷, Preview⁷, etc.), imazaquin (e.g. SCEPTER[®], TRI-SCEPT[®], etc. [except as noted in this label]), or imazethapyr (e.g. PURSUIT, PURSUIT[®] PLUS, PASSPORT[®], etc.) the same year as SQUADRON or injury to follow crops may occur.

DO NOT graze or feed treated soybean forage, hay or straw to livestock.

⁷Registered Trademarks of E.I. du Pont de Nemours and Company.
[®]Registered Trademarks of American Cyanamid Company.



WEED SCIENTIFIC NAMES

GRASSES

Barnyardgrass	(<i>Echinochloa crus-galli</i>)
Corn, volunteer	(<i>Zea mays</i>)
Crabgrass	(<i>Digitaria</i> spp.)
Crowfootgrass	(<i>Dactyloctenium aegyptium</i>)
Cupgrass, Woolly	(<i>Eriochloa villosa</i>)
Foxtail	
Giant	(<i>Setaria faberi</i>)
Green	(<i>Setaria viridis</i>)
Robust	(<i>Setaria</i> spp.)
Yellow	(<i>Setaria glauca</i>)
Goosegrass	(<i>Eleusine indica</i>)
Johnsongrass, seedling	(<i>Sorghum halepense</i>)
Panicum	
Browntop	(<i>Panicum fasciculatum</i>)
Fall	(<i>Panicum ichotriflorum</i>)
Texas	(<i>Panicum texanum</i>)
Sandbur, Field	(<i>Cenchrus incertus</i>)
Shattercane	(<i>Sorghum bicolor</i>)
Signalgrass, Broadleaf	(<i>Brachiaria platyphylla</i>)
Witchgrass	(<i>Panicum capillare</i>)

BROADLEAF WEEDS

Alligatorweed	(<i>Alternanthera philoxeroides</i>)
Beggarweed, Florida	(<i>Desmodium tortuosum</i>)
Bristly Starbur	(<i>Acanthospermum hispidum</i>)
Burcucumber	(<i>Sicyos angulatus</i>)
Carpetweed	(<i>Mollugo verticillata</i>)
Cocklebur, Common	(<i>Xanthium strumarium</i>)
Copperleaf, hophornbeam	(<i>Acalypha ostryifolia</i>)
Jimsonweed	(<i>Datura stramonium</i>)
Kochia	(<i>Kochia scoparia</i>)
Lambsquarters, Common	(<i>Chenopodium album</i>)
Mallow, Venice	(<i>Hibiscus trionum</i>)
Mexicanweed	(<i>Caperonia castanifolia</i>)
Morningglory	
Entireleaf	(<i>Ipomoea hederacea</i> var. <i>intergriuscula</i>)
Ivyleaf	(<i>Ipomoea hederacea</i>)
Palm Leaf	(<i>Ipomoea wrightii</i>)
Pitted	(<i>Ipomoea lacunosa</i>)
Smallflower	(<i>Jacquemontia tamnifolia</i>)
Tall	(<i>Ipomoea purpurea</i>)
Mustard Species	(<i>Brassica</i> spp.)
Nightshade, Eastern Black	(<i>Solanum ptycanthum</i>)
Pigweed	
Palmer	(<i>Amaranthus palmeri</i>)
Redroot	(<i>Amaranthus retroflexus</i>)
Smooth	(<i>Amaranthus hybridus</i>)
Spiny	(<i>Amaranthus spinosus</i>)
Waterhemp, Tall	(<i>Amaranthus tuberculatus</i>)

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

Poinsettia, Wild	(<i>Euphorbia heterophylla</i>)
Puncturevine	(<i>Tribulus terrestris</i>)
Purslane	(<i>Portulaca oleracea</i>)
Pusley, Florida	(<i>Richardia scabra</i>)
Ragweed	
Common	(<i>Ambrosia artemisiifolia</i>)
Giant	(<i>Ambrosia trifida</i>)
Redweed	(<i>Melochia corchorifolia</i>)
Sesbania, hemp	(<i>Sesbania exaltata</i>)
Sicklepod	(<i>Cassia obtusifolia</i>)
Sida, Prickly (Teaweed)	(<i>Sida spinosa</i>)
Smartweed	
Ladysthumb	(<i>Polygonum persicaria</i>)
Pennsylvania	(<i>Polygonum pensylvanicum</i>)
Spurge	
Annual	(<i>Euphorbia</i> spp.)
Prostrate	(<i>Euphorbia humistrata</i>)
Spotted	(<i>Euphorbia maculata</i>)
Sunflower, Common	(<i>Helianthus annuus</i>)
Texasweed	(<i>Caperonia palustris</i>)
Velvetleaf	(<i>Abutilon theophrasti</i>)

SEDGES

Nutsedge, Yellow	(<i>Cyperus esculentus</i>)
------------------	-------------------------------

