)

)

CHOPPER[®] RTU herbicide

*Equivalent to 3.0% 2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1<u>H</u>imidazol-2-yl]-3-pyridinecarboxylic ac.3 or 0.255 pounds acid per gallon.

EPA Reg. No. 241-330

EPA Est. No ACCEPTED

KEEP OUT OF REACH OF CHILDREN

CAUTIONI

PRECAUCION

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada impliamente.

STATEMENT OF PRACTICAL TREATMENT

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF IN EYES: Flush with plenty of water.

In case of an emergency endangering life or property involving this product, call collect, day or night. Area Code 201-835-3100.

See next page for Additional Precautionary Statements

Net Contents:

AMERICAN CYANAMID COMPANY AGRICULTURAL DIVISION VEGETATION AND PEST CONTROL WAYNE, NJ 07470 •1990

Registered Trademark of American Cyanamid Company

12/90

Under the Federal Insecticity, Fungicide, and Rodenticity Act, as provided, for the pesicide registered under UPA Rog. No. 201-330

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

CAUTIONI

Harmful if inhaled or absorbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water or wetlands (swamps, bogs, marshes or potholes). Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. DO NOT contaminate water when disposing of equipment washwaters.

PHYSICAL AND CHEMICAL HAZARDS

)

)

COMBUSTIBLE! DO NOT use or store near heat or open flame.

DISCLAIMER

The label instructions for use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be remable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the use or application of the product contrary to label instructions, all of which are beyond the control of American Cyanamid Company. All such risks shall be assumed by the user.

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions of use, subject to the risks referred to above.

Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or values of any other special or indirect damages.

American Cyanamid Company makes no other express or implied warranty, including other express or implied warranty of FITNESS or of MERCHANTABILITY.

12/90

.

• • • •

.

Dogy

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Observe all cautions and limitations on this label. This labeling must be in the possession of the user at time of pesticide application. DO NOT apply this product through any type of irrigation system.

STORAGE AND DISPOSAL

STORAGE: Shake before using if product has been stored below 0°F.

PROHIBITIONS: DO NOT contaminate water, food or feed by storage or disposal.

)

)

- PESTICIDE DISPOSAL: Wastes esulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
- CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, by incineration or, if allowed by State and Local authorities by burning. If burned, stay out of smoke.

IMPORTANT

DO NOT use on food or feed crops. DO NOT treat irrigation ditches or water used for crop irrigation or domestic purposes. DO NOT apply where runoff water may flow onto agricultural land as injury to crops may result. Keep from contact with fertilizers, insecticides, fungicides, desirable vegetation and seeds. DO NOT apply or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. DO NOT use on lawns, walks, driveways, tennis courts, or similar areas. DO NOT use in California.

Thoroughly clean application equipment after use. Flush tank, pump, hoses and boom with several changes of water after removing nozzle tips and screens (clean these parts separately).

GENERAL INFORMATION

CHOPPER RTU herbicide controls several woody species as a ready-to-use product when applied as a basal or cut stump treatment. With proper application CHOPPER RTU can prevent the resprouting of cut trees in forests, utility areas, fence rows, roadsides, railroads, pipelines, wildlife management areas, and other noncrop areas.

CHOPPER RTU herbicide is readily absorbed through bark and roots and is translocated rapidly throughout the plants. With dormant applications, leaf-out is often prevented and dieback occurs as the plant depletes its energy stores.

-3- | BEST AVAILABLE COPY

12/90

BASAL BARK TREATMENTS

To control unwanted woody species with stems less than 4 inches DBH (tree diameter at breast height), apply undiluted CHOPPER RTU to the lower 12-18 inches of the plant stem. Apply with a backpack type sprayer or power spraying equipment using low pressure at the spray nozzle. Spray the basal parts of brush and tree stems thoroughly wetting the indicated area. To insure control with stems larger than 4 inches DBH, use cut stump treatments. To obtain maximum performance, apply CHOPPER RTU during late summer, fall and winter months. Reduced efficacy may occur with applications during sap rise or leaf expansion. DO NOT over apply causing dripping or puddling.

CUT STUMP TREATMENTS

Spray or brush the CHOPPER RTU undiluted solution onto the cambium area of the freshly cut stump surface and the bark of the cut stump. Insure that the CHOPPER RTU solution thoroughly wets the cambium area, the wood next to the bark of stump. DO NOT over apply causing puddling.

CHOPPER RTU herbicide will control the following species:

Ash (Fraxinus spp.) Aspen (Populus spp.) Autumn Olive (Elaeagnus umbellata) Bigleaf Maple (<u>Acer macrophyllum</u>) Black Cherry (Prunus serotina) Boxelder (Acer negundo) Cottonwood (Populus spp.) Hickory (Carva spp.) Laurel Oak (Quercus laurifolia) Live Oak (Quercus virginiana) Maple (Acer spp.) Multiflora rose (Rosa multiflora) Persimmon (Diospyros virginiana) Red Alder (<u>Alnus</u> <u>rubra</u>) Red Oak (Quercus rubra) Russian Olive (Elaeagnus angustifolia) Saltcedar (Tamarix ramosissima) Sassafras (Sassafras albidum) Sumac (Rhus spp.) Sweet gum (<u>Liquidambar</u> styraciflua) Tulip Poplar (Liriodendron tulipifera) Turkey Oak (Quercus laevis) Water Oak (Quercus nigra) White Oak (<u>Quercus</u> <u>alba</u>) Willow (<u>Salix</u> spp.)

)

)

