

FEB 25 1992

Mr. Mark Galley
American Cyanamid Co.
P.O. Box 400
Princeton, NJ 08543

Dear Mr. Galley:

Subject: Supplemental Labels- Navy, Great Northern, Red
Kidney, Black Turtle, Cranberry, and Small White
Dry Beans in part of Michigan
PURSUIT PLUS® EC
EPA Reg. No. 241-331
Your submission dated Feb. 10, 1992

The labeling referred to above, submitted in connection
with registration under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended, is acceptable. A stamped
copy is enclosed for your records.

Sincerely yours,

Robert J. Taylor
Product Manager 25
Fungicide-Herbicide Branch
Registration Division (H7505C)

Enclosure

233

EPA Reg. No. 241-331

FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE, CRANBERRY, AND SMALL WHITE TYPE DRY BEANS ONLY

USE ONLY IN MICHIGAN (EXCLUDING UPPER PENINSULA)

THIS LABEL EXPIRES DECEMBER 31, 1992

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT PLUS EC LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT PLUS EC application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT PLUS EC if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT PLUS EC if cold and/or wet conditions are present or predicted to occur within one week of application.

USE PURSUIT PLUS EC ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT PLUS EC herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT PLUS EC TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT PLUS EC LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.


DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply PURSUIT PLUS EC in any manner not specifically described in this label.

Sensitive crops such as sugarbeets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT PLUS EC.

Spray equipment used for PURSUIT PLUS EC application must be drained and thoroughly cleaned with water before being used to apply other products.



*Registered Trademark of American Cyanamid Company



DO NOT apply to Domino variety black turtle beans.

DO NOT apply PURSUIT PLUS EC to sand or loamy sand soils.

DO NOT apply PURSUIT PLUS EC after June 20.

Allow at least 60 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT PLUS EC within 1 week before planting. Applied preplant incorporated, PURSUIT PLUS EC may be tank-mixed with other registered herbicides. Refer to the following table for use rate.

Soil Type	PURSUIT PLUS EC Use Rate (oz/acre)
Sands, loamy sands	DO NOT USE
Sandy loams and all other soils with \leq 2% O.M.	20 (1.25 pts)
All other soils with $>$ 2% O.M.	30

For control of the annual grass weed species listed on the PROWL® herbicide label, add 0.75 to 1 pt/acre of PROWL herbicide to the 20 oz/acre rate of PURSUIT PLUS EC, and 0.25 to 0.5 pt/acre of PROWL to the 30 oz/acre rate of PURSUIT PLUS EC.

WEEDS CONTROLLED

PURSUIT PLUS EC applied to low organic matter soils at the broadcast rate of 20 ounces per acre preplant incorporated will control light to moderate infestations of:

Nightshade, black
Nightshade, Eastern black
Pigweed, redroot

PURSUIT PLUS EC applied at the broadcast rate of 30 ounces per acre preplant incorporated will control:

Mustard, wild
Nightshade, black
Nightshade, Eastern black
Nightshade, hairy
Pigweed, redroot

PE-17136

*Registered Trademark of American Cyanamid Company

Peanuts

Edible bean types listed on this label (Refer to heading on page 1)

2. Four months after PURSUIT PLUS EC application:

Barley
Edible bean types not listed on this label (Refer to heading on page 1)
Peas
Wheat

3. Nine and one-half months after PURSUIT PLUS EC application:

Field corn
Field corn grown for seed*
Tobacco

4. Twenty-six months after PURSUIT PLUS EC application:

Canola (Rapeseed) Onions
Carrots Potatoes
Celery Spinach
Cole crops Sugarbeets
Garlic Tomatoes


DO NOT plant other rotational crops prior to 18 months after a PURSUIT PLUS EC application.

In the event of a crop loss, the bean crop may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT PLUS EC soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT PLUS EC the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT PLUS EC shall be assumed by the user.

CONSULT THE PURSUIT PLUS EC LEAFLET LABEL FOR OTHER PRECAUTIONS.



BEST AVAILABLE COPY