

PM 2: 241-331

1991


PURSUIT® PLUS EC SUPPLEMENTAL LABELING

EPA Reg. No. 241-~~310~~331

FOR USE IN SOYBEANS

PURSUIT® PLUS EC TANK-MIXTURE WITH SCEPTER®  
OR  
PURSUIT PLUS FOLLOWED BY SCEPTER®  
(SEE USE AREA BELOW)

THIS LABEL EXPIRES DECEMBER 31, 1992



OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT PLUS EC AND SCEPTER® LEAFLET LABELS BEFORE USING. SEE THE PURSUIT PLUS EC LEAFLET LABEL FOR MIXING AND APPLICATION INFORMATION, ADDITIONAL WEEDS CONTROLLED AND ROTATIONAL CROP RESTRICTIONS. ALWAYS FOLLOW THE MORE RESTRICTIVE LABEL.

GENERAL INFORMATION

For control of a broad-spectrum of weeds in soybeans, SCEPTER may be applied:

- 1) Tank-mixed with PURSUIT PLUS EC in soil application, or
- 2) Applied postemergence following soil application of PURSUIT PLUS EC.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system.

USE AREA

The application of PURSUIT PLUS EC and SCEPTER may be used in Region 2 as described in the USE AREA of the SCEPTER label and the entire state of Illinois.

The approved USE AREA (Region 2) includes eastern Kansas (east of highway U.S. 81; the counties of Cloud, Ellsworth, Harvey, Jewell, Lincoln, Mitchell, McPherson, Ottawa, Republic, Saline, Sedgewick, and Sumner), southeastern Nebraska (east of U.S. 81, south of U.S. 34), Missouri, Illinois, Indiana, Ohio, Michigan, Kentucky, Virginia, West Virginia, Pennsylvania, Maryland, Delaware and New Jersey.

\*Registered Trademark of American Cyanamid Company

### APPLICATION RATES

Apply PURSUIT PLUS EC at the rate of 2 1/2 pints per acre. (Soil applied).

Apply SCEPTER at the rate of 1/3 pint per acre. (Either soil applied in tank-mixture with PURSUIT PLUS EC or postemergence following PURSUIT PLUS EC). For application timing, refer to the PURSUIT PLUS EC soil-application directions and the SCEPTER postemergence applications directions in the respective labels.

### WEEDS CONTROLLED

The combination of PURSUIT PLUS EC and SCEPTER provides enhanced control of common cocklebur and common ragweed. Refer to the PURSUIT PLUS EC leaflet label for additional weeds controlled.

### ROTATIONAL CROP GUIDELINES

When using the above tank mixture or sequential application, rotational crops may be planted at the following intervals:

1. Anytime:  
Soybeans
2. Eleven months after SCEPTER application:  
Field corn (resistant)\*  
Wheat
3. Eighteen months after PURSUIT PLUS EC application.  
All crops not listed elsewhere in the ROTATIONAL CROP GUIDELINES section
4. Twenty six months after PURSUIT PLUS EC application.

Canola (Rapeseed)	Onions
Carrots	Potatoes
Celery	Spinach
Cole crops	Sugarbeets
Garlic	Tomatoes

Refer to ROTATIONAL CROP restrictions on the SCEPTER label. Follow the more restrictive rotational crop recommendations on the PURSUIT PLUS EC and SCEPTER labels.

\*Only field corn hybrids which possess tolerance or resistance (resulting from specific, qualitative genetic factors) to SCEPTER and/or other imidazolinone herbicides may be planted in the spring of the year following an application including SCEPTER. Contact your chemical dealer, seed supplier, or American Cyanamid for information on suitable field corn hybrids. Other field corn hybrids may be planted 18 months after an application including PURSUIT PLUS EC and SCEPTER.

NOTE: Availability of imidazolinone resistant field corn seed cannot be assured.

2054

Use of PURSUIT PLUS EC and SCEPTER herbicides in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risk associated with the use of these products and, therefore, rotational crop injury is always possible.

This label must be in the possession of the user at the time of herbicide application.

PE-17139

1/92

American Cyanamid Company  
Agricultural Division  
Crop Protection Chemicals Department  
Wayne, NJ 07470 ©1992

7105 241-331

1/24

**PURSUIT® PLUS EC SUPPLEMENTAL LABELING**

EPA Reg. No. 241-331

FOR USE IN SOYBEANS

SURFACE APPLICATIONS UP TO 45 DAYS PRIOR TO PLANTING

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT PLUS EC LEAFLET LABEL BEFORE USING. SEE THE PURSUIT PLUS EC LEAFLET LABEL FOR USE AREA, MIXING AND APPLICATION INFORMATION, WEEDS CONTROLLED AND ROTATIONAL CROP RESTRICTIONS.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system.

PURSUIT PLUS EC will control emerged problem weeds such as velvetleaf, smartweed, common cocklebur, and pigweeds. The weed size should not exceed 3" for velvetleaf and smartweed, and 8" for cocklebur and pigweeds. Add surfactant to the spray mixture at the rate of 1 qt./100 gal. AND a nitrogen based fertilizer (such as 28%N, 32%N or 10-34-0) at the rate of 1-2 qt./acre for optimum activity. Ammonium sulfate (s tray grade) may be substituted for liquid fertilizer at the rate of 4 lbs. per acre.

If other vegetation is present (not controlled by PURSUIT PLUS EC), apply paraquat, Roundup<sup>1</sup> or 2,4-D in combination with PURSUIT PLUS EC. DO NOT apply 2,4-D low volatile ester within 30 days of planting. Refer to the respective labels for rates, methods of application, proper timing, weeds controlled, restrictions, and precautions. Use ground equipment only.

When PURSUIT PLUS EC is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT PLUS EC cannot be mixed with any product containing a label prohibiting such mixtures.

This label must be in the possession of the user at the time of pesticide application.

•Registered Trademark of American Cyanamid Company

<sup>1</sup>Roundup is a trademark of Monsanto Agricultural Products Company.

PE-17147 2/20/92

American Cyanamid Company  
Agricultural Division  
Crop Protection Chemicals Department  
Wayne, NJ 07470 ©1992

