

EPA Reg No. 241-331

FOR USE IN SOYBEANS

PURSUIT® PLUS EC TANK-MIXTURE WITH SCEPTER®

THIS LABEL EXPIRES DECEMBER 31, 1993

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT PLUS EC AND SCEPTER LEAFLET LABELS BEFORE USING. SEE THE PURSUIT PLUS EC LEAFLET LABEL FOR MIXING AND APPLICATION INFORMATION, ADDITIONAL WEEDS CONTROLLED AND ROTATIONAL CROP RESTRICTIONS. ALWAYS FOLLOW THE MORE RESTRICTIVE LABEL.

GENERAL INFORMATION

For control of a broad-spectrum of weeds in soybeans, SCEPTER may be applied:

Tank-mixed with PURSUIT PLUS EC in soil application.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system.

USE AREA

The application of PURSUIT PLUS EC and SCEPTER may be used in Region 2 as described in the USE AREA of the SCEPTER label and the entire state of Illinois.

The approved USE AREA (Region 2) includes eastern Kansas (east of highway U. S. 81; the counties of Cloud, Ellsworth, Harvey, Jewell, Lincoln, Mitchell, McPherson, Ottawa, Republic, Saline, Sedgewick, and Sumner), southeastern Nebraska (east of U.S. 81, south of U.S. 34), Missouri, Illinois, Indiana, Ohio, Michigan, Kentucky, Virginia, West Virginia, Pennsylvania, Maryland, Delaware and New Jersey.

BEST AVAILABLE COPY

*Registered Trademark of American Cyanamid Company

(see reverse side)

APPLICATION RATES

Apply PURSUIT PLUS EC at the rate of 2-1/2 pints per acre.

Apply SCEPTER at the rate of 1/3 pint per acre. (Applied in tank-mixture with PURSUIT PLUS EC in soil applications).

For application timing, refer to the PURSUIT PLUS EC and SCEPTER soil-application directions in the respective labels.

WEEDS CONTROLLED

The combination of PURSUIT PLUS EC and SCEPTER provides enhanced control of common cocklebur and common ragweed. Refer to the PURSUIT PLUS EC leaflet label for additional weeds controlled.

ROTATIONAL CROP GUIDELINES

When using the above tank-mixture, rotational crops may be planted at the following intervals:

1. Anytime:
Soybeans
2. Eleven months after SCEPTER application:
Field corn (resistant)
Wheat

ACCEPTED

JUN - 3 1993

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 241-331

Cotton
Lettuce
Oats

Sunflower
Sorghum
Sweet corn

4. Twenty-six months after PURSUIT PLUS EC application:
All crops not listed elsewhere in the ROTATIONAL CROP GUIDELINES.
5. Forty months after PURSUIT PLUS EC application:
Sugarbeets

Refer to ROTATIONAL CROP restrictions on the SCEPTER label. Follow the more restrictive rotational crop recommendations on the PURSUIT PLUS EC and SCEPTER labels.

*Only field corn hybrids which possess tolerance or resistance (resulting from specific, qualitative genetic factors) to SCEPTER and/or other imidazolinone herbicides may be planted in the spring of the year following an application including SCEPTER. Contact your chemical dealer, seed supplier, or American Cyanamid for information on suitable field corn hybrids. Other field corn hybrids may be planted 18 months after an application including PURSUIT PLUS EC and SCEPTER.

NOTE: Availability of imidazolinone resistant field corn seed cannot be assured.

Use of PURSUIT PLUS EC and SCEPTER herbicides in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risk associated with the use of these products and, therefore, rotational crop injury is always possible.

This label must be in the possession of the user at the time of herbicide application.

PE-17139 Rev. 1 3/93 SD

 **CYANAMID**
American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1993