

SEP - 1 1993

Mrs. Desiree L. Little  
American Cyanamid Co.  
P.O. Box 400  
Princeton, NJ 08543

Dear Mrs. Little:

Subject: Supplemental Label for Fall Applications in S. Dakota  
PURSUIT® PLUS EO Herbicide  
EPA Reg. No. 241-331  
Your submission dated Aug. 27, 1993

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

This acceptance of your label does not relieve you of any obligation to comply with the Worker Protection Standard (WPS). Under the WPS labeling regulations at 40 CFR part 158, subpart K, § 156.200(c)(3), you are prohibited from distributing or selling any product within the scope of the WPS requirements after April 21, 1994, without amended labeling accepted by the Agency.

Sincerely yours,

Robert J. Taylor  
Product Manager 25  
Fungicide-Herbicide Branch  
Registration Division (H7505C)

Enclosure

RD:RTaylor:eja:9/1/93:3056800

PURSUIT® PLUS EC SUPPLEMENTAL LABELING

2 9 3  
ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

EPA Reg. No. 241-331

FOR USE IN SOYBEANS

FALL APPLICATIONS OF PURSUIT PLUS EC  
FOR USE ONLY IN THE STATE OF SOUTH DAKOTA

This Label Expires October 31, 1994

1993  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the  
registered use of EPA  
241-331

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT PLUS EC LEAFLET LABEL BEFORE USING. SEE THE PURSUIT PLUS EC LEAFLET LABEL FOR MIXING AND APPLICATION INFORMATION, WEEDS CONTROLLED AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system.

GENERAL INFORMATION

Late fall applications of PURSUIT PLUS EC may be made for control of weeds in NO-TILL soybeans, rather than in the spring prior to planting soybeans. Apply PURSUIT PLUS EC after October 31, and prior to ground freeze-up in the winter. Fall and winter precipitation will activate the herbicide for control of most winter annual weeds and spring germinating weeds in no-till soybeans.

When no-till planting soybeans following small grain harvest, apply a burndown application of Gramoxone<sup>1</sup> Extra or Roundup<sup>2</sup> to the small grain stubble within three weeks of harvest to control weeds present after harvest.

APPLICATION RATES

Apply PURSUIT PLUS EC at the rate of 2 1/2 pints/acre. If heavy grass pressure is expected, add PROWL® 3.3EC herbicide to the spray mixture at the rate of 1.25 pint/acre.

WEEDS CONTROLLED

Refer to the PURSUIT PLUS EC label for a list of weeds controlled.

®Registered Trademarks of American Cyanamid Company

<sup>1</sup>Trademark of ICI Americas, Inc.

<sup>2</sup>Trademark of Monsanto Company

3 7 3

## ROTATIONAL CROPS

---

The following rotational crops may be planted after applying PURSUIT PLUS EC at the recommended rate in soybeans:

1. Anytime
  - Lima beans
  - Southern peas
  - Soybeans
  - Peanuts
2. Four months after PURSUIT PLUS EC application:
  - Edible beans and peas (other than lima beans and Southern peas)
3. Nine and one-half months after PURSUIT PLUS EC application:
  - Barley
  - Field corn
  - Field corn grown for seed\*
  - Rye
  - Tobacco
  - Wheat
4. Eighteen months after a PURSUIT PLUS EC application.
  - Alfalfa
5. Twenty six months after PURSUIT PLUS EC application.
  - All crops not listed elsewhere in this ROTATIONAL CROPS guideline
6. Forty months after PURSUIT PLUS EC application.
  - Sugar beets

\* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT PLUS EC soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT PLUS EC the previous season. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT PLUS EC shall be assumed by the user.

Use of PURSUIT PLUS EC herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

This label must be in the possession of the user at the time of herbicide application.

PE- 7/93

American Cyanamid Company  
Agricultural Products Division  
Crop Protection Chemicals Department  
Wayne, NJ 07470 ©1993

SEP - 1 1993

Mrs. Desiree L. Little  
American Cyanamid Co.  
P.O. Box 400  
Princeton, NJ 08543

Dear Mrs. Little:

Subject: Supplemental Label for Fall Applications in S. Dakota  
PURSUIT® PLUS EO Herbicide  
EPA Reg. No. 241-331  
Your submission dated Aug. 27, 1993

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

This acceptance of your label does not relieve you of any obligation to comply with the Worker Protection Standard (WPS). Under the WPS labeling regulations at 40 CFR part 158, subpart K, § 156.200(c)(3), you are prohibited from distributing or selling any product within the scope of the WPS requirements after April 21, 1994, without amended labeling accepted by the Agency.

Sincerely yours,

Robert J. Taylor  
Product Manager 25  
Fungicide-Herbicide Branch  
Registration Division (H7505C)

Enclosure

RD:RTaylor:eja:9/1/93:3056800

**BEST AVAILABLE COPY**

PURSUIT® PLUS EC SUPPLEMENTAL LABELING

ACCEPTED  
with COMMENTS  
by EPA Letter dated

EPA Reg. No. 241-331

11-1-93

FOR USE IN SOYBEANS

FALL APPLICATIONS OF PURSUIT PLUS EC  
FOR USE ONLY IN THE STATE OF SOUTH DAKOTA

Under the Federal Insecticide, Fungicide, and Plant Growth Regulator Act, this product is approved for use in the State of South Dakota.  
241-331

This Label Expires October 31, 1994

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT PLUS EC LEAFLET LABEL BEFORE USING. SEE THE PURSUIT PLUS EC LEAFLET LABEL FOR MIXING AND APPLICATION INFORMATION, WEEDS CONTROLLED AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system.

GENERAL INFORMATION

Late fall applications of PURSUIT PLUS EC may be made for control of weeds in NO-TILL soybeans, rather than in the spring prior to planting soybeans. Apply PURSUIT PLUS EC after October 31, and prior to ground freeze-up in the winter. Fall and winter precipitation will activate the herbicide for control of most winter annual weeds and spring germinating weeds in no-till soybeans.

When no-till planting soybeans following small grain harvest, apply a burndown application of Gramoxone<sup>1</sup> Extra or Roundup<sup>2</sup> to the small grain stubble within three weeks of harvest to control weeds present after harvest.

APPLICATION RATES

Apply PURSUIT PLUS EC at the rate of 2 1/2 pints/acre. If heavy grass pressure is expected, add PROWL<sup>3</sup> 3.3EC herbicide to the spray mixture at the rate of 1.25 pint/acre.

WEEDS CONTROLLED

Refer to the PURSUIT PLUS EC label for a list of weeds controlled.

<sup>3</sup>Registered Trademark of American Cyanamid Company

<sup>1</sup>Trademark of ICI Americas, Inc.

<sup>2</sup>Trademark of Monsanto Company

BEST AVAILABLE COPY

## ROTATIONAL CROPS

The following rotational crops may be planted after applying PURSUIT PLUS EC at the recommended rate in soybeans:

1. Anytime
  - Lima beans
  - Southern peas
  - Soybeans
  - Peanuts
2. Four months after PURSUIT PLUS EC application:
  - Edible beans and peas (other than lima beans and Southern peas)
3. Nine and one-half months after PURSUIT PLUS EC application:
  - Barley
  - Field corn
  - Field corn grown for seed\*
  - Rye
  - Tobacco
  - Wheat
4. Eighteen months after a PURSUIT PLUS EC application.
  - Alfalfa
5. Twenty six months after PURSUIT PLUS EC application.
  - All crops not listed elsewhere in this ROTATIONAL CROPS guideline
6. Forty months after PURSUIT PLUS EC application.
  - Sugar beets

\* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT PLUS EC soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT PLUS EC the previous season. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT PLUS EC shall be assumed by the user.

Use of PURSUIT PLUS EC herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

This label must be in the possession of the user at the time of herbicide application.

PE- 1/93

American Cyanamid Company  
Agricultural Products Division  
Crop Protection Chemicals Department  
Wayne, NJ 07470 ©1993

**BEST AVAILABLE COPY**