

PM 10:00 AM 10/7/93

10/16

OCT 7 1993

Mrs. Desiree L. Little
American Cyanamid Co.
P.O. Box 400
Princeton, NJ 08543

Dear Mrs. Little:

Subject: Minor Changes
PURSUIT® PLUS EC Herbicide
EPA Reg. No. 241-331
Your submission dated Sept. 20, 1993

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

This acceptance of your label does not relieve you of any obligation to comply with the Worker Protection Standard (WPS). Under the WPS labeling regulations at 40 CFR part 158, subpart K, § 156.200(c)(3), you are prohibited from distributing or selling any product within the scope of the WPS requirements after April 21, 1994, without amended labeling accepted by the Agency.

Sincerely yours,

Robert J. Taylor
Product Manager 25
Fungicide-Herbicide Branch
Registration Division (H7505C)

Enclosure

BEST AVAILABLE COPY

200/16

PURSUIT® PLUS EC
herbicide

FOR USE IN SOYBEANS

ACTIVE INGREDIENTS:

| | |
|--|---------------|
| Imazethapyr (+)-2-[4,5-dihydro-4-methyl-4-(1-methyl-ethyl)-5-oxo-1H-imidazol-2-yl]-5-ethyl-3-pyridinecarboxylic acid | 2.25% |
| pendimethalin (N-1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine | 30.73% |
| INERT INGREDIENTS* | <u>67.02%</u> |
| TOTAL | 100.00% |

PURSUIT PLUS contains 2.9 pounds of active ingredients per gallon (2.7 pounds ai of pendimethalin and 0.2 pounds acid equivalent of imazethapyr).

*Contains petroleum distillates

EPA Reg. No. 241-331

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN

.CAUTION!/PRECAUCION!

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Do not induce vomiting. Call a physician or Poison Control Center immediately.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

NOTE TO PHYSICIAN: Because of increased risk of chemical pneumonia or pulmonary edema caused by aspiration of the hydrocarbon solvent, vomiting should be induced only under professional supervision.

In case of an emergency endangering life or property involving this product, call collect, day or night, area code 201-835-3100.

See Next Page for Additional Precautionary Statements

American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
WAYNE, NJ 07470 ©1993

ACCEPTED
with COMMENTS
In EPA Letter Dated:

Net Contents: 2.5 gallons
(9.45 liters)

OCT 7 1993

®Registered Trademark of American Cyanamid Company

Under the Federal Insecticide, Fungicide, and Plant Growth Regulator Act, this product is registered under EPA Reg. No. 241-331.

BEST AVAILABLE COPY

241 331/6/93

207/6

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)**

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. DO NOT apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. DO NOT contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of pesticide application.

Observe all cautions and limitations in this leaflet. Do not use PURSUIT PLUS EC other than in accordance with the instructions set forth on this label. The use of PURSUIT PLUS EC not consistent with this label can result in injury to crops, animals, or persons. Keep container closed to avoid spills and contamination.

DO NOT apply this product through any type of irrigation system.

STORAGE AND DISPOSAL

STORAGE: KEEP FROM FREEZING. DO NOT STORE BELOW 40°F. SHAKE WELL BEFORE USING.

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

BEST AVAILABLE COPY

DISCLAIMER

The label instructions for the use of this product reflect the opinion of experts based on research and field use. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, herbicide resistant weed populations, or the use of, or application of the product contrary to label instructions, all of which are beyond the control of American Cyanamid Company. All such risks shall be assumed by the user.

American Cyanamid Company shall not be responsible for losses or damages resulting from use of this product in any manner not set forth on this label. User assumes all risks associated with the use of this product in any manner not specifically set forth on this label.

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use, subject to the risks referred to above. **CYANAMID DOES NOT MAKE OR AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED AND EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.**

BUYER'S EXCLUSIVE REMEDY AND AMERICAN CYANAMID'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF PURSUIT PLUS EC. In no case shall Cyanamid or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product.

USES WITH OTHER PRODUCTS (TANK-MIXES)

If this product is used in combination with any other product except as specifically recommended in writing by American Cyanamid Company then American Cyanamid Company shall have no liability for any loss, damage or injury arising out of its use in any such combination not so specifically recommended. If used in combination recommended by American Cyanamid Company, the liability of American Cyanamid Company shall in no manner extend to any damage, loss or injury not directly caused by the inclusion of the American Cyanamid Company product in such combination use, and in any event shall be limited to return of the account of the purchase price of the American Cyanamid Company product.

20710

GENERAL INFORMATION

Entry into treated fields is not permitted without protective clothing until sprays have dried.

PURSUIT PLUS EC is effective in providing weed control in conservation tillage systems. PURSUIT PLUS EC may be applied in minimum tillage, no-till, or conventional tillage soybean production. Apply PURSUIT PLUS EC preplant or preplant incorporated from 45 days prior to planting up to planting. Incorporate PURSUIT PLUS EC within 7 days of application if rainfall is not received. PURSUIT PLUS EC may be surface applied (up to 45 days) prior to planting soybeans. After PURSUIT PLUS EC is applied, some susceptible weeds emerge, growth stops, and the weeds either die or are not competitive with the crop.

The weed killing activity of PURSUIT PLUS EC involves herbicide uptake by weed roots and rapid translocation to the growing points. Therefore, adequate soil moisture is important for optimum PURSUIT PLUS EC activity. When adequate soil moisture is present, PURSUIT PLUS EC will provide residual control of susceptible germinating weeds.

Occasionally, internode shortening and/or temporary yellowing of soybean plants may occur following PURSUIT PLUS EC applications. This will not affect soybean yields.

Use of PURSUIT PLUS EC herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

Within any weed population, there may be the potential for the occurrence of weeds resistant to herbicides when compounds that affect the same site of action are used continuously on the same acre. Therefore, any multiple year weed management program should include, when local agronomic practices allow, as many of the following practices as possible:

- use scouting and weed thresholds to determine the need for herbicide treatment
- utilize tank-mixtures or sequential applications of different mode-of-action herbicides to control the same target weeds
- combine cultivation with herbicide treatments

See your Cyanamid representative for additional information.

Replanting: If replanting is necessary in a field previously treated with PURSUIT PLUS EC, the field may be replanted to soybeans. Rework the soil no deeper than the treated zone. Do not apply a second treatment of PURSUIT PLUS EC.

BEST AVAILABLE COPY

MIXING INSTRUCTIONS

Fill the spray tank one-quarter to one-half full with clean water. While agitating add the required amount of product and then fill the remainder of the tank with water. Maintain agitation while spraying to ensure a uniform spray mixture.

TANK MIX COMBINATIONS WITH OTHER HERBICIDES

If other herbicides are tankmixed with PURSUIT PLUS EC, while agitating, add components in the following order:

- 1) Fill spray tank 1/4 to 1/2 full with clean water.
- 2) Add PURSUIT PLUS EC.
- 3) Add other EC (emulsifiable concentrate) products.
- 4) Add soluble packet products and thoroughly mix.
- 5) Add WP (wetttable powder), DG (dispersible granules), DF (dry flowable) formulations not in soluble packets, or liquid flowables .
- 6) Add aqueous solution products.
- 7) Add liquid dispersible products.
- 8) While agitating, fill the remainder of the tank with water.

BEST AVAILABLE COPY

When paraquat is included in a tank-mixture, add 8 ounces of non-ionic surfactant per 100 gallons of spray mixture as the last ingredient in the tank.

To avoid injury to sensitive crops, spray equipment used for PURSUIT PLUS EC applications must be drained and thoroughly cleaned with water before being used to apply other products.

HERBICIDE COMBINATIONS

PURSUIT PLUS EC may be applied either preplant, preplant incorporated or preemergence in tank-mix combination with metribuzin (Sencor² or Lexone³) or preemergence with linuron (Lorox³ or Linex⁴). PURSUIT PLUS EC may also be applied preplant incorporated, followed by a preemergence application of metribuzin or linuron. The tank-mix or sequential application of PURSUIT PLUS EC and metribuzin will enhance the control of common cocklebur and common ragweed in addition to the weeds controlled by a soil application of PURSUIT PLUS EC. The addition of linuron to PURSUIT PLUS EC will enhance the control of common ragweed.

Under conditions of heavy grass pressure, the addition of a grass herbicide such as PROWL 3.3 EC or trifluralin may be tank-mixed with PURSUIT PLUS EC.

See recommendation for addition of Roundup or paraquat to the spray solution under the NO-TILL OR REDUCED TILLAGE section.

When PURSUIT PLUS EC is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT PLUS EC cannot be mixed with any product containing a label prohibiting such mixtures.

¹Roundup is a trademark of Monsanto Agricultural Products Company.

²Sencor is a trademark of Mobay Corporation.

³Lexone and Lorox are trademarks of E.I. duPont de Nemours and Company.

⁴Linex is a trademark of Griffin Ag Products Co., Inc.

[®]Registered trademark of American Cyanamid Company.

SPRAYING INSTRUCTIONS

DO NOT apply when wind velocity is greater than 10 mph, or when spray may be carried to sensitive crops. Sensitive crops include leafy vegetables, sugarbeets, and cotton.

GROUND APPLICATIONS:

Uniformly apply with properly calibrated ground equipment in 10 to 40 gallons of water per acre. A spray pressure of 20 to 40 psi is recommended.

APPLICATIONS WITH LIQUID FERTILIZER:

PURSUIT PLUS EC can be applied to the soil in liquid fertilizers, alone or in combination with PROWL 3.3 EC, trifluralin, Lasso⁵ or Dual⁶. Follow all PURSUIT PLUS EC label recommendations regarding incorporation, timing of application, special instructions and precautions. Apply treatments in 20 or more gallons of liquid fertilizer per acre with ground equipment. Always test the compatibility of PURSUIT PLUS EC with liquid fertilizer before mixing in the spray tank.

APPLICATIONS WITH DRY BULK FERTILIZER:

PURSUIT PLUS EC may be impregnated on dry bulk fertilizers. When applied as directed, PURSUIT PLUS EC/dry bulk fertilizer mixtures provide weed control equal to that provided by the same rates of PURSUIT PLUS EC applied in water or liquid fertilizer.


Follow all PURSUIT PLUS EC label recommendations regarding application and incorporation, special instructions and precautions. Apply PURSUIT PLUS EC/dry bulk fertilizer mixtures only with ground equipment.

All individual state regulations relating to dry bulk fertilizer, registration, labeling, and application are the responsibility of the individual and/or company selling the PURSUIT PLUS EC/dry bulk fertilizer mixtures.

A minimum of 200 pounds and a maximum of 450 pounds of dry bulk fertilizer impregnated with the recommended amount of PURSUIT PLUS EC must be applied per acre.

DO NOT impregnate PURSUIT PLUS EC onto coated ammonium nitrate or limestone because these materials will not absorb the herbicide. Dry fertilizer blends containing mixtures of ammonium nitrate or limestone may be impregnated with PURSUIT PLUS EC. A minimum of 200 pounds of impregnated dry bulk fertilizer, excluding the weight of ammonium nitrate or limestone, must be applied per acre.

Apply PURSUIT PLUS EC at the rate of 2-1/2 pints per acre. Use the following table to determine the amount of PURSUIT PLUS EC to be impregnated on a ton of dry bulk fertilizer based on the rate of fertilizer which will be applied per acre.



⁵Lasso is a trademark of Monsanto Agricultural Products Company.

⁶Dual is a trademark of Ciba Geigy Corporation.

(12-12)

RATE CHART FOR IMPREGNATION OF DRY BULK FERTILIZER WITH PURSUIT PLUS EC
(Pints of PURSUIT PLUS EC per Ton of Fertilizer)

| <u>PURSUIT PLUS EC Rate Per Acre</u> | <u>Fertilizer Rate pounds per acre</u> | <u>Pints per Ton</u> |
|--|--|----------------------|
| 2-1/2 pints | 200 | 25 |
| | 250 | 20 |
| | 300 | 16-2/3 |
| | 350 | 14-1/3 |
| | 400 | 12-1/2 |
| | 450 | 11 |

For those rates not listed in this table, calculate the pints of PURSUIT PLUS EC to be impregnated on a ton of dry bulk fertilizer using the following formula:

$$\frac{2000}{\text{Pounds of dry fertilizer per acre}} \times \frac{2-1/2 \text{ Pints of PURSUIT PLUS EC per acre (recommended rate)}}{1} = \text{Pints of PURSUIT PLUS EC per ton of fertilizer}$$

To impregnate PURSUIT PLUS EC on bulk fertilizer, use a closed rotary-drum mixer or other commonly used dry bulk fertilizer blender equipped with suitable spray equipment. Spray nozzles must be placed to provide uniform coverage of PURSUIT PLUS EC onto the fertilizer during mixing.

If PROWL 3.3 EC is to be combined with the PURSUIT PLUS EC prior to impregnation, premix the PROWL 3.3 EC with an equal volume of water before adding it to the PURSUIT PLUS EC. DO NOT mix undiluted PROWL 3.3 EC with PURSUIT PLUS EC.

Apply the PURSUIT PLUS EC/dry bulk fertilizer mixture with an accurately calibrated dry fertilizer spreader. The PURSUIT PLUS EC/dry bulk fertilizer mixture must be spread uniformly on the soil surface. Uneven spreading can cause poor weed control and crop injury.

Refer to PREPLANT INCORPORATED APPLICATIONS section of this label for incorporation directions.

7-17-14

AERIAL APPLICATIONS:

Uniformly apply with properly calibrated aerial equipment in 5 or more gallons of water per acre.

To avoid injury to sensitive crops from drift, aerial applicators must adhere to the following SPECIAL AERIAL USE DIRECTIONS AND PRECAUTIONS:

- Use nozzles which produce a coarse spray.
- Nozzle height above ground must be a maximum of 10 feet.
- Nozzles must be pointed toward the rear of the aircraft. The downward angle of the nozzle should not be greater than 20 degrees.
- To minimize wing-tip vortex roll, nozzles or spray boom must not be located any closer to end of wing or rotor than three-fourths the distance from the center of the aircraft.
- Use a maximum spray pressure of 40 psi.
- A buffer zone must be established between the area to be sprayed and sensitive crops.
- DO NOT spray when wind velocity is greater than 5 mph.

Applicator is responsible for any loss or damage which results from spraying PURSUIT PLUS EC in manner other than recommended on this label. In addition, applicator must follow all applicable state and local regulations and ordinances in regard to spraying.

PURSUIT PLUS EC HERBICIDE USE AREA

Do not apply PURSUIT PLUS EC to soybeans in North Dakota or in Minnesota north of state highway 210.

USE RATE
(2-1/2 PINTS PER ACRE)

Apply PURSUIT PLUS EC at a broadcast rate of 2 1/2 pints per acre for all methods of application: preplant, preplant incorporated and preemergence (including minimum and no-till). At this broadcast rate, one gallon of PURSUIT PLUS EC will treat 3.2 acres of soybeans.

NOTE: Only one application of PURSUIT PLUS EC may be made during the season.

WEEDS CONTROLLED

When applied as directed, PURSUIT PLUS EC will control or reduce competition from the weeds listed below.

NOTE: C = Control, R = Reduced Competition

BEST AVAILABLE COPY

0.17.2

BROADLEAF WEEDS

| Weeds Controlled | Preplant Incorporated | Preemergence |
|-------------------------|--------------------------|--------------|
| Anoda, spurred | C | C |
| Buffalobur | C | C |
| Carpetweed | C | C |
| Cocklebur, common | R | |
| Devilsclaw | C | |
| Galinsoga | C | C |
| Jimsonweed | C | |
| Kochia | C | C |
| Lambquarters, common | C | |
| Mallow, Venice | R | |
| Morningglory | | |
| entireleaf | R | |
| ivyleaf | R | |
| pitted | R | |
| smallflower | C | C |
| tall | R | |
| Mustard species | C | C |
| Nightshade | | |
| black | C | C |
| Eastern black | C | C |
| hairy | C | C |
| Pigweed | | |
| palmer | C | C |
| redroot | C | C |
| smooth | C | C |
| spiny | C | C |
| Puncturevine | C | C |
| Purslane, common | C | C |
| Pusley, Florida | C | C |
| Ragweed, | | |
| common | R | R |
| giant | R | R |
| Sida, prickly (teaweed) | C | |
| Smartweed | | |
| ladysthumb | C | C |
| Pennsylvania | C | C |
| Spurge | | |
| prostrate | C | C |
| spotted | C | C |
| Sunflower | C | |
| Velvetleaf | C | |
| Waterhemp, tall | C | C |

11/27/10

GRASS WEEDS

| Weeds Controlled | Preplant Incorporated | Preemergence |
|------------------------|-----------------------|--------------|
| Barnyardgrass | C | C |
| Crabgrass | | |
| large | C | C |
| smooth | C | C |
| Crowfootgrass | C | C |
| Cupgrass, woolly | C | |
| Foxtail | | |
| giant | C | C |
| green | C | C |
| yellow | C | C |
| Goosegrass | C | C |
| Itchgrass | R | |
| Johnsongrass | | |
| seedling | C | C |
| rhizome | R | |
| Millet, wild proso | R | R |
| Panicum | | |
| browntop | C | C |
| fall | C | C |
| Texas | C | C |
| Sandbur, field | C | C |
| Shattercane | C | |
| Signalgrass, broadleaf | C | C |
| Sorghum almum | C | R |
| Witchgrass | C | C |
| SEDGES | | |
| Nutsedge | | |
| yellow | R | |
| purple | R | |

20110

SOIL APPLICATION INSTRUCTIONS

PURSUIT PLUS EC provides effective weed control in conservation tillage systems designed to meet conservation compliance requirements. PURSUIT PLUS EC can be applied as an early preplant, preplant incorporated, or preemergence treatment in soybeans. It can also be applied in conventional, minimum tillage and no-till production systems. The application method of choice will depend on the anticipated weed spectrum and the preference of the applicator. If weeds have emerged, see instructions under NO-TILL OR REDUCED TILLAGE section.

PURSUIT PLUS EC controls weeds by uptake by weed roots, and translocation to the growing points where it stops weed growth.

Adequate soil moisture is required for optimum activity. For surface applications, rainfall or overhead irrigation is necessary to move PURSUIT PLUS EC into the weed germination zone. The amount of rainfall or irrigation required following application depends on existing soil moisture, soil texture and organic matter content. Sufficient water to moisten the soil to a depth of 2 inches is normally adequate. If adequate moisture is not received within 7 days after a surface-applied treatment, then a cultivation is recommended to control escaped weeds. When adequate moisture is received after dry conditions, PURSUIT PLUS EC will provide residual control of susceptible germinating weeds; activity on established weeds will depend on the weed species and the location of its root system in the soil.

In ridge-till plantings, PURSUIT PLUS EC may be applied early preplant or at soybean planting. If the herbicide is banded over the row, cultivation will be required for weed control between the beds. If cultivation is not possible or if weed pressure is heavy, apply PURSUIT PLUS EC in a broadcast application. Use proportionally less PURSUIT PLUS EC per acre in a band versus a broadcast application. If rainfall does not occur within 7 days of application, a rotary hoe incorporation will enhance weed control. See PREEMERGENCE APPLICATIONS and PREPLANT INCORPORATED APPLICATIONS for further information.

PREEMERGENCE APPLICATIONS

SURFACE APPLICATIONS MADE BEFORE PLANTING

PURSUIT PLUS EC may be surface applied prior to soybean planting (up to 45 days). If sufficient rain does not occur before planting to activate PURSUIT PLUS EC, shallow incorporation before planting will enhance weed control. PURSUIT PLUS EC may be surface applied prior to soybean planting both north and south of highway I-80.

SURFACE APPLICATIONS MADE AFTER PLANTING

PURSUIT PLUS EC may be surface applied up to 2 days after soybean planting (before crop emergence) South of Interstate Highway I-80 only. DO NOT APPLY PURSUIT PLUS EC AFTER SOYBEAN PLANTING North of Interstate Highway I-80.

BEST AVAILABLE COPY

1- 15716

NO-TILL OR REDUCED TILLAGE:

PURSUIT PLUS EC is effective in controlling weeds in conservation tillage production systems. Apply PURSUIT PLUS EC treatments up to 45 days prior to planting (early preplant) but before crop emergence.

To ensure thorough coverage, use a minimum of 20 gallons of water per acre. Use higher gallonage for fields with dense vegetation or heavy crop residues. Adjust the boom height to ensure proper coverage of weed foliage (according to the manufactures recommendation). The nozzle spacing on the boom should be 20 inches. Use only standard flat-fan nozzle tips. Use ground equipment only.

PURSUIT PLUS EC will control emerged problem weeds such as velvetleaf, smartweed, common cocklebur, and pigweeds. The weed size should not exceed 3" for velvetleaf and smartweed, and 8" for cocklebur and pigweeds. Add surfactant to the spray mixture at the rate of 1 qt./100 gal. AND a nitrogen based fertilizer (such as 28%N, 32%N or 10-34-0) at the rate of 1-2 qt./acre for optimum activity. Ammonium sulfate (spray grade) may be substituted for liquid fertilizer at the rate of 2.5 lbs. per acre.

If other vegetation is present (not controlled by PURSUIT PLUS EC), apply paraquat, Roundup or 2,4-D in combination with PURSUIT PLUS EC. Refer to the respective labels for rates, methods of application, proper timing, weeds controlled, restrictions, and precautions. Use ground equipment only.

NOTE: Adjust planters to ensure adequate seed coverage.

110210

PREPLANT INCORPORATED APPLICATIONS

PURSUIT PLUS EC may be applied following land preparation and should be thoroughly incorporated to a depth of 1 to 2 inches. When applied to beds, maintain PURSUIT PLUS EC in the surface 1 to 2 inches of the finished beds. Application may be made up to 45 days prior to planting (early preplant). Incorporate prior to soybean planting and within 7 days of application. Mechanical incorporation can be achieved by the following:

- (a) Disk harrow set to cut 3 to 4 inches deep and operated at 4 to 6 mph with the second pass at an angle to the first.
- (b) PTO-driven equipment (tillers, cultivators, hoes) set to cut 2-3 inches deep and operated one time at 4 mph or less.
- (c) Rolling cultivator set to cut 2 to 4 inches deep and operated two times at 6 to 8 mph. Use only on coarse textured soils.
- (d) Field cultivator with 3 to 4 rows of sweeps (Do not use chisel points), spaced at intervals of 7 inches or less and staggered so that no soil is left unturned, set to cut 3 inches deep, and operated two times at more than 5 mph with the second pass made at an angle to the first.
- (e) LELY-Roterra⁷ set and operated according to the manufacturer's directions to thoroughly incorporate PURSUIT PLUS EC to a depth of 1 to 3 inches.
- (f) Do-All set and operated according to the manufacturer's directions to incorporate PURSUIT PLUS EC 1 to 3 inches deep and operated at 4 to 6 mph.

If soybeans are planted on beds, apply and incorporate after bed formation using PTO-driven equipment or a rolling cultivator. For optimum weed control, PURSUIT PLUS EC should be maintained in the surface 1-2 inches of the finished bed.

⁷LELY-Roterra is a trademark of Lely.

12-24-11

ROTATIONAL CROPS

The following rotational crops may be planted after applying PURSUIT PLUS EC at the recommended rate in soybeans:

1. Anytime
 - Lima beans
 - Southern peas
 - Soybeans
 - Peanuts

2. Four months after PURSUIT PLUS EC application:
 - Wheat
 - Edible beans and peas (other than lima beans and Southern peas)

4. Nine and one-half months after PURSUIT PLUS EC application:
 - Barley
 - Field corn
 - Field corn grown for seed*
 - Rye
 - Tobacco

5. Eighteen months after PURSUIT PLUS EC application:
 - Alfalfa
 - Cotton
 - Lettuce
 - Oats
 - Popcorn
 - Safflower
 - Sorghum
 - Sweet corn

6. Twenty six months after PURSUIT application.
 - All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.

7. Forty months after PURSUIT application.
 - Sugarbeets

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT PLUS EC soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT PLUS EC the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT PLUS EC shall be assumed by the user.

Use of PURSUIT PLUS EC herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

PRECAUTIONS

If soybeans are furrow irrigated, till the soil prior to planting winter wheat or barley. The beds should be broken up and the soil mixed with tillage equipment set to cut 4-6 inches deep.

Only rotational crops harvested at maturity may be used for feed or food.

In the event of a crop loss due to weather, soybeans can be replanted. DO NOT work the soil deeper than 2 inches.

There should be an interval of at least 85 days between an application of PURSUIT PLUS EC and soybean harvest.

DO NOT graze or feed treated soybean forage, hay or straw to livestock.

DO NOT apply products containing chlorimuron ethyl (Classic⁸, Canopy⁸, Gemini⁸, Lorox Plus⁸, PREVIEW⁸, etc.) or imazaquin (SCEPTER[®], SQUADRON[®], or TRI-SCEPT[®], SCEPTER[®] O.T.[®], or SCEPTER[®] 70DG,) or products containing imazethapyr (PURSUIT[®], PURSUIT[®] DG or PASSPORT[®]) the same year as PURSUIT PLUS EC or injury to follow crops may occur.

DO NOT tank-mix PURSUIT PLUS EC with clomazone containing herbicides (Command⁹ or Commence⁹).

⁸Classic, Canopy, Preview, Gemini and Lorox Plus are trademarks of E.I. duPont de Nemours and Company.

⁹Command and Commence are trademarks of FMC Corporation.

[®]Registered Trademarks of American Cyanamid Company.

For additional information regarding the use of PURSUIT PLUS EC herbicide, call telephone no. 800-942-0500.

American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1993