

FM 25

241-331

107-3

FEB 14 1994

Mrs. Desiree L. Little
American Cyanamid Co.
P.O. Box 400
Princeton, NJ 08540

Dear Mrs. Little:

Subject: Supplemental Label for Use on Navy, Great
Northern, Red Kidney, Black Turtle, Cran-
Berry, and Small White Type Dry Beans
For Distribution and Use in Michigan
Pursuit® Plus Herbicide
EPA Reg. No. 241-331
Your submission dated Jan. 20, 1994

The labeling referred to above, submitted in connection
with registration under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended, is acceptable. A stamped
copy is enclosed for your records.

Sincerely yours,

Robert J. Taylor
Product Manager 25
Fungicide-Herbicide Branch
Registration Division (H7505C)

Enclosure

BEST AVAILABLE COPY

2013

EPA Reg. No. 241-331

FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE, CRANBERRY, AND SMALL WHITE TYPE DRY BEANS ONLY

USE ONLY IN MICHIGAN (EXCLUDING UPPER PENINSULA)

THIS LABEL EXPIRES DECEMBER 31, 1994

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT PLUS EC LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT PLUS EC application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. DO NOT apply PURSUIT PLUS EC if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT PLUS EC if cold and/or wet conditions are present or predicted to occur within one week of application

USE PURSUIT PLUS EC ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT PLUS EC herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT PLUS EC TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT PLUS EC LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT PLUS EC in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT PLUS EC.

Spray equipment used for PURSUIT PLUS EC application must be drained and thoroughly cleaned with water before being used to apply other products.

**ACCEPTED
WITH COMMENTS
by EPA Letter Draft**

*Registered Trademark of American Cyanamid Company

FEB 14 1994

Under the Federal Insecticide, Fungicide, and Rodenticide Act, this product is registered under

241-331

(see reverse side)

BEST AVAILABLE COPY

DO NOT apply PURSUIT PLUS EC after June 20.

Allow at least 60 days between application and harvest.

Preplant Incorporated Applications. Apply PURSUIT PLUS EC within 1 week before planting. Applied preplant incorporated, PURSUIT PLUS EC may be tank-mixed with other registered herbicides. Refer to the following table for use rate.

Soil Type	PURSUIT PLUS EC Use Rate (oz/acre)
Sands, loamy sands	DO NOT USE
Sandy loams and all other soils with \leq 2% O.M.	20 (1.25 pts)
All other soils with > 2% O.M.	30

For control of the annual grass weed species listed on the PROWL® herbicide label, add 0.75 to 1 pt/acre of PROWL herbicide to the 20 oz/acre rate of PURSUIT PLUS EC, and 0.25 to 0.5 pt/acre of PROWL to the 30 oz/acre rate of PURSUIT PLUS EC.

WEEDS CONTROLLED

PURSUIT PLUS EC applied to low organic matter soils at the broadcast rate of 20 ounces per acre preplant incorporated will control light to moderate infestations of:

Nightshade, black
Nightshade, Eastern black
Pigweed, redroot

PURSUIT PLUS EC applied at the broadcast rate of 30 ounces per acre preplant incorporated will control:

Mustard, wild
Nightshade, black
Nightshade, Eastern black
Nightshade, hairy
Pigweed, redroot

PE-17136 Rev 2 11/1993 SD

®Registered Trademark of American Cyanamid Company

Soybeans

Peanuts

Edible bean types listed on this label (Refer to heading on page 1)

IMI-Corn

2. Four months after PURSUIT PLUS EC application:

Barley

Edible bean types not listed on this label (Refer to heading on page 1)

Peas

Wheat

3. Nine and one-half months after PURSUIT PLUS EC application:

Field corn

Field corn grown for seed*

Tobacco

4. Eighteen months after PURSUIT PLUS EC application:

Alfalfa

Oats

Sorghum

Cotton

Popcorn

Sweet Corn

Lettuce

Safflower

5. Twenty-six months after PURSUIT PLUS EC application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.

6. Forty months after PURSUIT PLUS EC application:

Sugar beets

In the event of a crop loss, the bean crop may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT PLUS EC soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT PLUS EC the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT PLUS EC shall be assumed by the user.

CONSULT THE PURSUIT PLUS EC LEAFLET LABEL FOR OTHER PRECAUTIONS.



American Cyanamid Company
Agriculture & Markets Division
Crop Protection Chemicals Department
Wayne, NJ 07470

BEST AVAILABLE COPY