


PURSUIT® PLUS EC SUPPLEMENTAL LABELING



EPA Reg. No. 241-331

FOR USE IN SOYBEANS

PURSUIT® PLUS EC TANK-MIXTURE WITH SCEPTER® 70DG

THIS LABEL EXPIRES DECEMBER 31, 1994

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT PLUS EC AND SCEPTER® 70DG LEAFLET LABELS BEFORE USING. SEE THE PURSUIT PLUS EC LEAFLET LABEL FOR MIXING AND APPLICATION INFORMATION, ADDITIONAL WEEDS CONTROLLED AND ROTATIONAL CROP RESTRICTIONS. ALWAYS FOLLOW THE MORE RESTRICTIVE LABEL.

GENERAL INFORMATION

For control of a broad-spectrum of weeds in soybeans, SCEPTER 70DG may be applied tank-mixed with PURSUIT PLUS EC in soil applications only.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system.

USE AREA

The application of PURSUIT PLUS EC and SCEPTER 70DG may be used in Region 2 as described in the USE AREA of the SCEPTER 70DG label and the entire state of Illinois.

The approved USE AREA (Region 2) includes eastern Kansas (east of highway U.S. 81; the counties of Cloud, Ellsworth, Harvey, Jewell, Lincoln, Mitchell, McPherson, Ottawa, Republic, Saline, Sedgewick, and Sumner), southeastern Nebraska (east of U.S. 81, south of U.S. 34), Missouri, Illinois, Indiana, Ohio, Michigan, Kentucky, Virginia, West Virginia, Pennsylvania, Maryland, Delaware and New Jersey.



®Registered Trademark of American Cyanamid Company

2 of 11

APPLICATION RATES

Apply PURSUIT PLUS EC at the rate of 2 1/2 pints per acre.

Apply SCEPTER 70DG at the rate of 1.4 ounces per acre. (One (1) bag of SCEPTER 70DG treats 10 acres. Apply in tank-mixture with PURSUIT PLUS EC in soil applications).

For application timing, refer to the PURSUIT PLUS EC and SCEPTER 70DG soil-application directions in the respective labels.

WEEDS CONTROLLED

The combination of PURSUIT PLUS EC and SCEPTER 70DG provides enhanced control of common cocklebur and common ragweed. Refer to the PURSUIT PLUS EC leaflet label for additional weeds controlled.

ROTATIONAL CROP GUIDELINES

When using the above tank mixture, rotational crops may be planted at the following intervals:

- 1. Anytime:
Soybeans
- 2. Eleven months after PURSUIT PLUS EC + SCEPTER 70DG application:
Field corn (resistant)*
Field corn (Michigan only)**
Wheat
- 4. Eighteen months after PURSUIT PLUS EC + SCEPTER 70DG application:
Alfalfa
Cotton
Lettuce
Oats
Popcorn
Safflower
Sorghum
Sweet corn
- 5. Twenty six months after PURSUIT PLUS EC + SCEPTER 70DG application.
All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.
- 6. Forty months after PURSUIT PLUS EC + SCEPTER 70DG application.
Sugarbeets

BEST AVAILABLE COPY

Refer to ROTATIONAL CROP restrictions on the SCEPTER 70DG label. Follow the more restrictive rotational crop recommendations on the PURSUIT PLUS EC and SCEPTER 70DG labels.

*Only field corn hybrids which possess tolerance or resistance (resulting from specific, qualitative genetic factors) to SCEPTER 70DG and/or other imidazolinone herbicides may be planted in the spring of the year following an application including SCEPTER 70DG. Contact your chemical dealer, seed supplier, or American Cyanamid for information on suitable field corn hybrids. Other field corn hybrids may be planted 18 months after an application including PURSUIT PLUS EC and SCEPTER 70DG.

**In the state of Michigan, field corn (non-resistant) may be planted the year following an application of SCEPTER 70DG unless extreme drought conditions develop (less than 15 inches of rainfall or irrigation is received from two weeks prior to the date of last application through November 15 of the same year).

If the minimum rainfall requirements is not met, only IMI-Corn hybrids (hybrids classified as tolerant or resistant to imidazolinone herbicides) may be planted the spring of the year following PURSUIT PLUS EC and SCEPTER 70DG applications. Contact your chemical dealers, seed supplier or American Cyanamid to obtain a list of IMI-Corn (resistant/tolerant) hybrids.

NOTE: Availability of imidazolinone resistant field corn seed cannot be assured.

Use of PURSUIT PLUS EC and SCEPTER 70DG herbicides in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risk associated with the use of these products and, therefore, rotational crop injury is always possible.

This label must be in the possession of the user at the time of herbicide application.

BEST AVAILABLE COPY

ACCEPTED

APR - 9 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 241-331

EPA Reg. No. 241-331

**FOR USE IN FIELD CORN
(APPLY ONLY ON IMI-CORN HYBRIDS)**

**DIRECTIONS FOR THE USE OF PURSUIT PLUS EC FOR WEED CONTROL
IN IMI-CORN HYBRIDS**

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT PLUS EC LEAFLET LABEL BEFORE USING. SEE THE PURSUIT PLUS EC LEAFLET LABEL FOR PRECAUTIONS, WORKER PROTECTION STANDARD REQUIREMENTS, APPLICATION INFORMATION, USES WITH OTHER PRODUCTS AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of pesticide application.

The use of PURSUIT PLUS EC herbicide not consistent with this label can result in injury to crops, animals, or persons. Keep container closed to avoid spills and contamination.

DO NOT apply this product through any type of irrigation system.

This label expires December 31, 1994.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT PLUS EC TO IMI-CORN.

GENERAL INFORMATION

Apply PURSUIT PLUS EC only on selected field corn hybrids (IMI-Corn) warranted by the seed company to possess resistance/tolerance to direct application of PURSUIT. **DO NOT** apply PURSUIT PLUS EC to corn hybrids which lack genetic resistance/tolerance to PURSUIT herbicide. Contact your chemical dealer, seed supplier or American Cyanamid to obtain a list of IMI-Corn hybrids. With the exception of reduced tillage systems (see NO-TILL or REDUCED TILLAGE section) plant into a firm seed bed that is free of clods and crop residue. Use only where adequate tillage is practiced to provide good soil coverage of the corn seed. Plant corn **AT LEAST** 1-1/2 inches deep to ensure good seed coverage.

PURSUIT PLUS EC herbicide can be applied as a preemergence or postemergence treatment to IMI-Corn hybrids. PURSUIT PLUS EC provides effective weed control in conventional and reduced tillage (conservation tillage, minimum tillage, and no-till) systems. After PURSUIT PLUS EC is applied, some susceptible weeds emerge, growth stops, and the weeds either die or are not competitive with the crop.

Crops growing under stressful environmental conditions can exhibit various injury symptoms which may be more pronounced if herbicides are used. Stunting, leaf curling or temporary yellowing of the field corn plants occasionally may occur following PURSUIT PLUS EC applications. Normal growth and appearance should resume within 1 to 2 weeks.

Within any weed population, there may be the potential for the occurrence of weeds resistant to herbicides when compounds that affect the same site of action are used continuously on the same acre. Therefore, any multiple year weed management program should include, when local agronomic practices allow, as many of the following practices as possible:

- use scouting and weed thresholds to determine the need for herbicide treatment
- utilize tank-mixes or sequential applications of different mode-of-action herbicides to control the same target weeds
- combine cultivation with herbicide treatments

See your Cyanamid representative for additional information.

Replanting: If replanting is necessary in a field previously treated with PURSUIT PLUS EC herbicide, the field may be replanted to soybeans, peanuts, lima beans or southern peas. Re-work the soil no deeper than the treated zone. Do not apply a second treatment of PURSUIT PLUS EC.

BEST AVAILABLE COPY

agitation while spraying to ensure a uniform spray mixture.

When tank-mixing PURSUIT PLUS EC with recommended herbicides, add the other herbicides and adjuvants in the following order while agitating:

1. Fill spray tank 1/4 to 1/2 full with clean water.
2. Add PURSUIT PLUS EC.
3. Add other EC (emulsifiable concentrate) products.
4. Add soluble packet products and thoroughly mix.
5. Add WP (wettable powder), DG (dispersible granule), DF (dry flowable) or LF (liquid flowable) formulations.
6. Add aqueous solution products.
7. Add surfactant or crop oil to the spray tank.
8. Add liquid fertilizer.
9. While agitating, fill the remainder of the tank with water.

To avoid injury to sensitive crops, spray equipment used for PURSUIT PLUS EC herbicide applications must be drained and thoroughly cleaned with water before being used to apply other products.

USE RATE (2-1/2 PINTS PER ACRE)

Apply PURSUIT PLUS EC at a broadcast rate of 2-1/2 pints per acre for all methods of application: preemergence and postemergence. At this broadcast rate, one gallon of PURSUIT PLUS EC will treat 3.2 acres of IMI-Corn.

NOTE: Only one application of PURSUIT PLUS EC may be made during the growing season.

PURSUIT PLUS EC HERBICIDE USE AREA

DO NOT apply PURSUIT PLUS EC to corn in North Dakota or in Minnesota north of state highway 210.

SPRAYING INSTRUCTIONS

DO NOT apply when wind velocity is greater than 10 mph, or when spray may be carried to sensitive crops. Sensitive crops include leafy vegetables, sugar beets, and cotton.

APPLICATION INSTRUCTIONS POST EMERGENCE

Apply PURSUIT PLUS EC herbicide as a postemergence treatment to IMI-Corn when crop and weeds are actively growing. Apply PURSUIT PLUS EC before weeds exceed a height of 3", unless otherwise indicated. More restrictive crop growth stage limitations of tank-mix partners MUST be followed.

DO NOT apply PURSUIT PLUS EC with a liquid fertilizer as a carrier.

DO NOT apply PURSUIT PLUS EC when air temperatures are expected to reach or stay below 40° F for 10 or more hours or when extended cold, wet conditions are predicted.

DO NOT tank-mix PURSUIT PLUS EC herbicide with Accent™ or Beacon™.

ADDITIVES:

POSTEMERGENCE APPLICATIONS OF PURSUIT PLUS EC REQUIRE THE ADDITION OF A SURFACTANT AND A LIQUID FERTILIZER.

I. SURFACTANTS

Use a non-ionic surfactant containing at least 80% active ingredient. Apply the surfactant at the rate of 1 quart per 100 gallons of spray solution

AND

II. LIQUID FERTILIZER SOLUTION¹

Recommended nitrogen based fertilizers include liquid fertilizers (such as 28%N, 32%N or 10-34-0) at the rate of 1-2 quarts per Acre. Use the higher rate when weeds are under moisture or temperature stress. Instead of a liquid fertilizer, spray grade ammonium sulfate may be used at the rate of 2.5 lbs per Acre.

SOIL APPLICATION INSTRUCTIONS

DO NOT INCORPORATE PURSUIT PLUS EC.

PURSUIT PLUS EC provides effective weed control in conservation tillage systems designed to meet conservation compliance requirements. PURSUIT PLUS EC herbicide can be applied as a preemergence treatment in IMI-Corn. It can also be applied in conventional, minimum tillage and no-till production systems. The application method of choice will depend on the anticipated weed spectrum and the preference of the applicator. If weeds have emerged, see instructions under NO-TILL OR REDUCED TILLAGE section.

SURFACE APPLICATIONS MADE AFTER PLANTING:

Apply PURSUIT PLUS EC alone preemergence (surface treatment only) after planting. DO NOT INCORPORATE PURSUIT PLUS EC or serious corn injury may result.

¹PURSUIT PLUS EC applications may be made with a non-ionic surfactant only (liquid fertilizer not required) in the states of Arkansas, Bootheel of Missouri, Tennessee, South Carolina, Louisiana, Mississippi, Alabama, and Georgia.

NO-TILL OR REDUCED TILLAGE:

PURSUIT PLUS EC is effective in controlling weeds in conservation tillage production systems.

PURSUIT PLUS EC herbicide will control emerged problem weeds such as velvetleaf, smartweeds, common cocklebur, and pigweeds. The weed size should not exceed 3" for velvetleaf and smartweed, and 8" for cocklebur and pigweeds. Add surfactant to the spray mixture at the rate of 1 quart per 100 gallons AND a nitrogen based fertilizer (such as 28%N, 32%N, or 10-34-0) at the rate of 1-2 quarts per acre for optimum activity. Ammonium sulfate (spray grade) may be substituted for liquid fertilizer at the rate of 2.5 lbs. per acre.

NOTE: Plant corn at least 1-1/2 inches deep. Adjust planters to ensure adequate seed coverage.

The use of no-till planters in minimum tillage corn under conditions which do not allow good soil coverage of the corn seed can result in reduced crop stand or injury if PURSUIT PLUS EC contacts the germinating corn seed. Check equipment to ensure good seed coverage.

AERIAL APPLICATIONS

Uniformly apply with properly calibrated aerial equipment in 5 or more gallons of water per acre. When applied POSTEMERGENCE, the addition of a non-ionic surfactant AND fertilizer solution are required for optimum weed control. Apply a non-ionic surfactant at the rate of 1 quart per 100 gallons of spray mixture AND a liquid fertilizer at the rate of 1 quart per acre. (See instructions under APPLICATION INSTRUCTIONS - POSTEMERGENCE.)

To avoid injury to sensitive crops from drift, aerial applicators must adhere to the following SPECIAL AERIAL USE DIRECTIONS AND PRECAUTIONS.

- Nozzle height above ground must be a maximum of 10 feet.
- Nozzles must be pointed toward the rear of the aircraft. The downward angle of the nozzle should not be greater than 20 degrees.
- To minimize wing-tip vortex roll, nozzles or spray boom must not be located any closer to end of wing or rotor than three-fourths the distance from the center of the aircraft.
- Use a maximum spray pressure of 40 psi.
- A buffer zone must be established between the area to be sprayed and sensitive crops.
- DO NOT spray when wind velocity is greater than 5 mph. Coarse sprays (larger droplets) are less likely to drift.

Applicator is responsible for any loss or damage which results from spraying PURSUIT PLUS EC in a manner other than recommended in this label. In addition, applicator must follow all applicable state and local regulations and ordinances in regard to spraying.

SEQUENTIAL HERBICIDE COMBINATIONS WITH PURSUIT PLUS EC

PURSUIT PLUS EC FOLLOWING EARLY PRE-PLANT APPLICATION OF:

Bicep™, Bicep Lite™	Frontier™
Bronco™	Gramoxone Extra™
Bullet™	Lasso EC™, Lasso MT™
Dual™	Roundup™
Extrazine II™	

PURSUIT PLUS EC FOLLOWING PREPLANT INCORPORATED APPLICATION OF:

Atrazine ¹	Extrazine II
Bicep, Bicep Lite	Frontier
Bullet	Lariat™
Dual	Lasso EC, Lasso MT
Eradicane Extra™	Sutan+™

PURSUIT PLUS EC FOLLOWING PREEMERGENCE APPLICATION OF:

Atrazine	Extrazine II
Banvel™, 2	Frontier
Bicep, Bicep Lite	Lariat
Bullet	Lasso EC, Lasso MT
Clarity™, 2	Marksman™, 3
Dual	PROWL ³ 3.3EC
Eradicane Extra	Sutan+

TANK-MIX HERBICIDE COMBINATIONS WITH PURSUIT PLUS EC

PREEMERGENCE APPLICATION OF PURSUIT PLUS EC TANK-MIXED WITH:

Atrazine	Extrazine II
Banvel	Frontier
Bicep, Bicep Lite	Lariat
Bullet	Lasso EC, Lasso MT
Clarity	Marksman
Dual	PROWL 3.3EC
Eradicane Extra	Sutan+

POSTEMERGENCE APPLICATION OF PURSUIT PLUS EC TANK MIXED WITH:

Atrazine ¹	Clarity
Banvel ²	Marksman
Buctril™, 2, 3	PROWL 3.3EC
Basagran™, 2	Senco: DF™, Senco: 3L™

¹In some cases the grass activity of PURSUIT PLUS EC will be reduced when used in combination with atrazine, Buctril, Banvel, Basagran, Clarity, or Marksman postemergence.

²Some corn leaf burn may result with Buctril or atrazine postemergence combinations with PURSUIT PLUS EC.

³Applications of Banvel or Clarity herbicide to corn during periods of rapid growth may result in temporary leaf burn.



When PURSUIT PLUS EC herbicide is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions.

WEEDS CONTROLLED

When applied as directed, PURSUIT PLUS EC will control or reduce competition from the weeds listed below.

NOTE: C = Control, R = Reduced Competition

BROADLEAF WEEDS

Weeds Controlled	PREEMER- GENCE	Leaf Stage	POSTEMERGENCE MAX. SIZE (Inches)
Alligator weed		4	1 - 3
Anoda, spurred	C	2	1 - 2
Artichoke, Jerusalem		8	6 - 10
Buffalobur	C*	R	1 - 3
Carpetweed	C		
Cocklebur, common	R	8	1-8
Galinsoga	C		
Jimsonweed	C*	4	1 - 3
Kochia	C	4	1 - 3
Lambsquarters, common	C*	2	1 - 2
Mallow, Venice	R		
Marshelder	C	4	1 - 3
Morningglory,			
entireleaf	R	2	1 - 2
ivyleaf	R	2	1 - 2
pitted	R	2	1 - 2
smallflower	C	4	1 - 3
tall	R	2	1 - 2
Mustard species	C	4	1 - 3
Nightshade,			
black	C	4	1 - 3
Eastern black	C	4	1 - 3
hairy	C	4	1 - 3
Pigweed,			
palmer	C	8	1 - 8
redroot	C	8	1 - 8
smooth	C	8	1 - 8
spiny	C	8	1 - 8
Poinsettia, wild	C		
Puncturevine	C		
Purslane, common	C		
Pusley, Florida	C		
Ragweed,			
common	R	4	1 - 3
giant	R	4	1 - 3
Sage, barnyard	R	R	1 - 3
Sida, prickly (teaweed)	C*		
Smartweed,			
lady	C	4	1 - 3
Pennsylvania	C	4	1 - 3

BROADLEAF WEEDS

Weeds Controlled	PREEMER- GENCE	Leaf Stage	POSTEMERGENCE MAX. SIZE (Inches)
Spurge,			
prostrate	C	4	1 - 3
spotted	C	4	1 - 3
Sunflower	C*	4	1 - 3
Thistle, Canada		R	1 - 3
Velvetleaf	C*	4	1 - 3
Waterhemp, tall	C	4	1 - 3

* Cultivation and/or a postemergence herbicide may be required for season-long control.

The number under Maximum Leaf Stage indicates the MAXIMUM number of leaves at which weeds should be sprayed postemergence. DO NOT count cotyledon leaves when determining weed stage of growth.

GRASS WEEDS

Weeds Controlled	PREEMER- GENCE	Leaf Stage	POSTEMERGENCE MAX. SIZE (Inches)
Barnyardgrass	R	3	1 - 3
Crabgrass,			
large	C	3	1 - 3
smooth	C	3	1 - 3
Crowfootgrass	C		
Cupgrass, woolly	C	3	1 - 3
Foxtail,			
giant	C	6	1 - 6
green	C	3	1 - 3
yellow	C	3	1 - 3
Goosegrass	R		
Johnson grass,			
seedling	C	6	1 - 8
rhizome		R	1 - 8
Millet, wild proso	R	R	
Panicum,			
browntop	C		
fall	C		
Texas	C		
Red rice		3	1 - 3
Sandbur, field	C		
Shattercane	R	6	1 - 8
Signalgrass, broadleaf	R	4	1 - 8
Sorghum alnum	R		
Witchgrass	C	6	1 - 3

SEDGES

Nutsedge,			
yellow	R	R	1 - 3
purple	R	R	1 - 3



ROTATIONAL CROPS

The following rotational crops may be planted after applying PURSUIT PLUS EC herbicide at the recommended rate in field corn:

1. Anytime

Lima beans	Peanuts
Southern peas	Soybeans
2. Four months after PURSUIT PLUS EC application:

Wheat
Edible beans and peas (other than lima beans and Southern peas)
4. Nine and one-half months after PURSUIT PLUS EC application:

Barley	Rye
Field corn	Tobacco
Field corn grown for seed*	
5. Eighteen months after PURSUIT PLUS EC application:

Alfalfa	Popcorn
Cotton	Safflower
Lettuce	Sorghum
Oats	Sweet corn
6. Twenty six months after PURSUIT PLUS EC application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.
7. Forty months after PURSUIT PLUS EC application:

Sugar beets

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT PLUS EC herbicide soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT PLUS EC the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT PLUS EC shall be assumed by the user.

Use of PURSUIT PLUS EC herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

PRECAUTIONS

In the event of a crop loss due to weather, soybeans, lima beans, southern peas or peanuts can be replanted. DO NOT work the soil deeper than 2 inches.

There should be an interval of at least 45 days between an application of PURSUIT PLUS EC herbicide and corn harvest (silage, fodder, or grain).

DO NOT graze or feed treated corn, forage, silage, fodder, or grain to livestock for at least 45 days after an application of PURSUIT PLUS EC.

If field corn is furrow irrigated, till the soil prior to planting winter wheat or barley. The beds should be broken up and the soil mixed with tillage equipment set to cut 4-6 inches deep.

Only rotational crops harvested at maturity may be used for feed or food.

All soil insecticides, including labelled banded or in-furrow applications, may be used in combination with Pioneer imidazolinone resistant (IR) corn hybrids.

Imidazolinone tolerant hybrids from other seed companies may occasionally exhibit injury symptoms when soil insecticides are used in combination with PURSUIT PLUS EC. USE banded applications of COUNTER® 15G, COUNTER® 20CR® or THIMET® in combination with PURSUIT PLUS EC application; DO NOT USE COUNTER 15G soil insecticide products in-furrow with imidazolinone tolerant hybrids. American Cyanamid has not tested all hybrids in which the imidazolinone tolerance trait is claimed and cannot be responsible for factors which are beyond its control, such as growing conditions, environmental conditions, grower practices and the specific genetics of each hybrid which may influence tolerance. Growers should contact the seed company for information regarding specific hybrid tolerance to herbicide and insecticide applications.

For additional information regarding the use of PURSUIT PLUS EC herbicide, call telephone no. (800) 942-0500.

PE-17189 Rev. 1 8/3/93 SD


*Registered Trademark of American Cyanamid Company.
 *Roundup, Bronco, Bullet, Lariat, Lasso EC, and Lasso MT are Trademarks of Monsanto Agricultural Products.
 *Accent and Extrazine II are Trademarks of E.I. DuPont de Nemours and Company, Inc.
 *Banvel, Clarity, Frontier, and Marksman are Trademarks of Sandoz Crop Protection Corporation
 *Buctril is a Trademark of Rhone Poulenc Ag Company.
 *Basagran is a Trademark of BASF Corporation
 *Sencor is a Trademark of Mohay Corporation.
 *Beacon, Bicep, Bicep Linc, and Dual are Trademarks of Ciba Geigy Corporation.
 *Gramoxone Extra, Eradicane Extra and Sutan+ are Trademarks of ICI Americas, Inc



FOR USE IN SOYBEANS

FALL APPLICATIONS OF PURSUIT PLUS EC
FOR USE ONLY IN THE STATE OF SOUTH DAKOTA

This Label Expires October 31, 1994



OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT PLUS EC LEAFLET LABEL BEFORE USING. SEE THE PURSUIT PLUS EC LEAFLET LABEL FOR MIXING AND APPLICATION INFORMATION, WEEDS CONTROLLED, AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system.

GENERAL INFORMATION

Late fall applications of PURSUIT PLUS EC may be made for control of weeds in NO-TILL soybeans, rather than in the spring prior to planting soybeans. Apply PURSUIT PLUS EC after October 31, and prior to ground freeze-up in the winter. Fall and winter precipitation will activate the herbicide for control of most winter annual weeds and spring germinating weeds in no-till soybeans.

When no-till planting soybeans following small grain harvest, apply a burndown application of Gramoxone¹ Extra or Roundup² to the small grain stubble within three weeks of harvest to control weeds present after harvest.

¹Registered Trademarks of American Cyanamid Company
¹Trademark of ICI Americas, Inc.
²Trademark of Monsanto Agricultural Company

APPLICATION RATES

Apply PURSUIT PLUS EC at the rate of 2-1/2 pints/acre. If heavy grass pressure is expected, add PROWL[®] 3.3EC herbicide to the spray mixture at the rate of 1.25 pint/acre.

WEEDS CONTROLLED


Refer to the PURSUIT PLUS EC label for a list of weeds controlled.

ROTATIONAL CROPS

The following rotational crops may be planted after applying PURSUIT PLUS EC at the recommended rate in soybeans:

1. Anytime

- Lima beans
- Southern peas
- Soybeans
- Peanuts



2. Four months after PURSUIT PLUS EC application:

- Edible beans and peas (other than lima beans and Southern peas)

(see reverse side)



Barley
Field corn
Field corn grown for seed*

Rye
Tobacco
Wheat

4. Eighteen months after a PURSUIT PLUS EC application:
Alfalfa
5. Twenty six months after PURSUIT PLUS EC application.
All crops not listed elsewhere in this ROTATIONAL CROPS guideline.
6. Forty months after PURSUIT PLUS EC application:
Sugar beets

*Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT PLUS EC soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT PLUS EC the previous season. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT PLUS EC shall be assumed by the user.

Use of PURSUIT PLUS EC herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

This label must be in the possession of the user at the time of herbicide application.

PE-17200 7/93 SD

BEST AVAILABLE COPY

 **CYANAMID**
American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 4-1993