

EPA Reg. No. 241-331

FOR USE IN SOYBEANS

VARIABLE RATES OF PURSUIT PLUS EC FOR ENHANCED WEED CONTROL

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT PLUS EC LEAFLET LABEL BEFORE USING. SEE THE PURSUIT PLUS EC LEAFLET LABEL FOR PRECAUTIONS, WORKER PROTECTION STANDARD REQUIREMENTS, APPLICATION INFORMATION, USES WITH OTHER PRODUCTS AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at the time of herbicide application. The use of PURSUIT PLUS EC herbicide not consistent with this labeling may result in injury to crops. Keep container closed to avoid spills and contamination. DO NOT apply this product through any type of irrigation system.

GENERAL INFORMATION

PURSUIT PLUS EC provides control of many broadleaf and grass weeds when applied preplant and preemergence to soybeans when applied at 2 1/2 pints per acre. If difficult to control weeds are present, or if a high population of weeds is anticipated, PURSUIT PLUS EC may be applied up to 3 3/4 pints per acre. The higher rate will provide more consistent control of velvetleaf, sunflower and giant ragweed. It will also provide more consistent control of late germinating grasses.

MIXING INSTRUCTIONS

Fill the spray tank one-fourth to one-half full with clean water or liquid fertilizer. Add the required amount of PURSUIT PLUS EC to the spray tank first and make sure it is thoroughly mixed. When tank-mixing PURSUIT PLUS EC with Roundup¹ products add Roundup (or other glyphosate containing products) to the tank while agitating. Add adjuvants to the spray mixture (for no-till burndown applications). Then fill the remainder of the tank with clean water. Maintain agitation while spraying to ensure a uniform spray mixture. To avoid injury to sensitive crops, spray equipment used for PURSUIT PLUS EC herbicide applications must be drained and thoroughly cleaned with water before being used to apply other products.

USE RATE

Apply PURSUIT PLUS EC at the rate of 2 1/2 pints to 3 3/4 pints per acre.
Refer to the Roundup (glyphosate) or other tank mixture product label for recommended use rates.

NOTE: Only one application of PURSUIT PLUS EC may be made during the growing season.

ACCEPTED
~~with COMMENTS~~
In EPA Letter Dated

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

241-331

preemergence application up to 45 days before planting. Preemergence applications may also be applied during or after planting but before the crop emerges. DO NOT apply PURSUIT PLUS EC postemergence (or at cracking) to soybeans.

Adequate moisture is required for activation of PURSUIT PLUS EC.

PREPLANT INCORPORATED APPLICATIONS

Apply PURSUIT PLUS EC before planting and incorporate uniformly into the top 1 to 2 inches of soil.

PURSUIT PLUS EC may be applied immediately before planting or up to 45 days prior to planting. Incorporate within 7 days after application.

If soybeans are planted on beds, apply and incorporate after bed formation, using PTO-driven equipment or rolling cultivator.

SURFACE APPLICATIONS (including no-till)

Apply PURSUIT PLUS EC up to 45 days before, during or after planting but before the crop emerges. As with other herbicides applied preemergence, rainfall or irrigation is necessary to activate PURSUIT PLUS EC in the soil. If sufficient rainfall or irrigation to activate PURSUIT PLUS EC is not received within 7 days after application, a thorough shallow tillage, cultivation, or postemergence treatment may be required for control of emerged weeds.

Refer to the PURSUIT PLUS EC label (and other product labels when tank-mixing with other products) for additional information regarding WEEDS CONTROLLED, MIXING AND SPRAYING INSTRUCTIONS, PRECAUTIONARY STATEMENTS, RESTRICTIONS AND WORKER PROTECTION INFORMATION.

When tank-mixing herbicides, always follow the most restrictive label.

ROTATIONAL CROP GUIDELINES

The following rotational crops may be planted after applying PURSUIT PLUS EC at the recommended rate:

1. Anytime

Lima beans	Soybeans
Southern peas	Peanuts

2. Four months after PURSUIT PLUS EC application:
 - Wheat
 - Edible beans and peas (other than lima beans and Southern peas)

3. Eight and one-half months after PURSUIT PLUS EC application:
 - Field corn*
 - Field corn grown for seed*

4. Nine and one-half months after PURSUIT PLUS EC application:
 - Barley
 - Rye
 - Tobacco

5. Eighteen months after PURSUIT PLUS EC application:

Alfalfa	Safflower
Cotton	Sorghum
Lettuce	Sunflower
Oats	Sweet corn
Popcorn	

6. Twenty six months after PURSUIT PLUS EC application.
 - Potatoes

7. Forty months after PURSUIT PLUS EC application
 - All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.

NOTE: Following forty months after a PURSUIT PLUS EC application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop

* Field corn may be planted 8 1/2 months following an application of PURSUIT PLUS EC when applied at 2 1/2 pints per acre. If more than 2 1/2 pints per acre is applied (up to the maximum of 3 3/4 pints per acre) field corn may be planted 8 1/2 months following an application of PURSUIT PLUS EC unless extreme drought conditions develop (less than 15 inches of rainfall or irrigation is received from two weeks prior to the date of PURSUIT PLUS EC application through November 15 of the same year). If the minimum rainfall requirement is not met, only field corn hybrids (IMI-Corn™) which possess tolerance or resistance to PURSUIT PLUS EC may be planted the spring of the year following a PURSUIT PLUS EC application.

Use of PURSUIT PLUS EC herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

©Registered Trademark of American Cyanamid Company

™IMI-Corn is a Trademark of American Cyanamid Company

‡Trademark of Monsanto Agricultural Products Company

PE xxxx 12/95 SD

American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 1995©

IN EPA LABEL 12/89

MAR 28 1996

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
241-331

EPA Reg. No. 241-331

FOR USE IN SOYBEANS

5715

PURSUIT PLUS EC HERBICIDE AND STATUS¹ HERBICIDE FOR WEED CONTROL IN SOYBEANS

OBSERVE ALL PRECAUTIONARY AND WORKER PROTECTION STATEMENTS IN THE PURSUIT PLUS EC LEAFLET LABEL BEFORE USING. SEE THE PURSUIT PLUS EC LEAFLET LABEL FOR USE AREA, MIXING AND APPLICATION INFORMATION, WEEDS CONTROLLED AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of herbicide application. DO NOT apply this product through any type of irrigation system.

GENERAL INFORMATION

When soil applied for weed control in soybeans, PURSUIT PLUS EC controls many problem broadleaf and grass weeds. Under adverse weather conditions or when heavy weed populations exist, some problem weeds may not be completely controlled by PURSUIT PLUS EC. STATUS herbicide provides fast postemergence burn-down of susceptible broadleaf weeds. Temporary bronzing of the soybean leaves may occur soon after application. Soybeans soon outgrow early symptoms; yield will not be effected by applying STATUS following an application of PURSUIT PLUS EC.

WEEDS CONTROLLED

A postemergence application of STATUS herbicide following PURSUIT PLUS EC at the recommended rates will control common and giant ragweeds, pigweed species and waterhemp. (Refer to the STATUS label for additional weeds controlled).

APPLICATION RATES

Sequential Application:

When PURSUIT PLUS EC is soil applied, followed by STATUS applied postemergence apply PURSUIT PLUS EC at the rate of 2-12 pints per acre. Apply STATUS postemergence at the rate indicated in the table below, depending on weed size:

Common ragweed Pigweed species, Waterhemp, tall commor			
Giant Ragweed		1 - 6"	6 - 8"***

*Use the higher rate if common ragweed is present or the weed population is high.

**Use the 24 ounce/acre rate if giant ragweed is 6 - 8 inches tall.

MIXING INSTRUCTIONS

POSTEMERGENCE APPLICATIONS OF STATUS REQUIRE THE ADDITION OF AN ADJUVANT TO ACHIEVE CONSISTENT WEED CONTROL.

ADJUVANTS

- I. **SURFACTANTS:** Use a non-ionic surfactant containing at least 80% active ingredient. Apply the surfactant at the rate of 1 quart per 100 gallons.

OR

CROP OIL CONCENTRATE: Instead of a surfactant, a petroleum or vegetable seed based oil concentrate (such as SUN-IT II²) may be used. SUN-IT II is recommended when weeds are under moisture or temperature stress. Use SUN-IT II or other methylated seed oils at the rate of 1.5 to 2 pints per acre (use the higher rate when weeds are at the maximum label size), or use a crop oil concentrate at 2 pints per acre. A non-ionic surfactant is recommended when crop response is of concern (i.e., under conditions of high relative humidity or the presence of dew on the crop leaves).

AND

II. LIQUID FERTILIZER SOLUTION

For more consistent weed control, the addition of a nitrogen based fertilizer to the spray solution is recommended. Include liquid fertilizers (such as 28%N, 32%N or 10-34-0) at the rate of 1-2 quarts per acre. Use the higher rate when weeds are under moisture or temperature stress. Instead of a liquid fertilizer, spray grade ammonium sulfate may be used at the rate of 2.5 lbs. per acre.

Refer to the STATUS label for additional information regarding application rates, restrictions, precautions, weeds controlled, adjuvants recommended and other information

¹Trademark of BASF Corporation

²Trademark of Agsco, Inc.

Registered Trademark of American Cyanamid Company

© 1996 American Cyanamid Company

American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 01/96

7/15
8

EPA Reg. No. 241-331

FOR USE IN SOYBEANS

**PURSUIT PLUS EC HERBICIDE AND PRESTIGE¹ HERBICIDE FOR
WEED CONTROL IN SOYBEANS**

OBSERVE ALL PRECAUTIONARY AND WORKER PROTECTION STATEMENTS IN THE PURSUIT PLUS EC LEAFLET LABEL BEFORE USING. SEE THE PURSUIT PLUS EC LEAFLET LABEL FOR USE AREA, MIXING AND APPLICATION INFORMATION, WEEDS CONTROLLED AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of herbicide application. DO NOT apply this product through any type of irrigation system.

GENERAL INFORMATION

When soil applied for weed control in soybeans, PURSUIT PLUS EC controls many problem broadleaf and grass weeds. Under adverse weather conditions or when heavy weed populations exist, some problem weeds may not be completely controlled by PURSUIT PLUS EC. PRESTIGE herbicide provides postemergence control of many grass weeds. Refer to the table below for a partial list of weeds and the recommended use rate.

WEEDS CONTROLLED

A postemergence application of PRESTIGE herbicide following PURSUIT PLUS EC at the recommended rates will control the grasses listed below. (Refer to the PRESTIGE label for additional weeds controlled).

USE RATES

Apply PURSUIT PLUS EC at the broadcast rate of 2 and 1/2 pints per acre for all methods of application: preplant, preplant incorporated, and preemergence. At this broadcast rate, one gallon of PURSUIT PLUS EC will treat 3.2 acres.

Refer to the following table for the recommended rate of PRESTIGE.

ACCEPTED
~~WITH COMMENTS~~
IN EPA LETTER DATED
11/28/79

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

241-331

Grass	(Inches)	Acre (oz.)	(Inches)	Acre (oz.)	(Inches)	Acre (oz.)
Barnyardgrass	4	18	8	24	12	36
Crabgrass, Large	-	-	6	24	8	36
Smooth	-	-	6	24	8	36
Cupgrass, Woolly	-	-	8	24	-	-
Foxtail, Giant	4	18	8	24	16	36
Green	4	18	8	24	16	36
Yellow	-	-	8	24	16	36
Goosegrass	3	18	6	24	8	36
Johnsongrass Seedling	-	-	8	24	16	36
Millet, Wild Proso	10	12	10	12	24	24
Panicum, Browntop	-	-	8	24	-	-
Fall	4	18	8	24	12	36
Texas	4	18	8	24	12	24
Shattercane	-	-	18	24	-	-
Signalgrass Broadleaf	4	18	8	24	12	36
Volunteer, Barley	-	-	4	36	-	-
Corn	12	18	20	24	-	-
Oats	-	-	4	36	-	-
Rye	-	-	4	36	-	-
Wheat	-	-	4	36	-	-
Witchgrass	-	-	8	24	-	-

*If a mixture of grasses are present, use the highest rate indicated for the grasses present.

ADDITIVES

POSTEMERGENCE APPLICATIONS OF PRESTIGE REQUIRE THE ADDITION OF A CROP OIL CONCENTRATE AND A LIQUID FERTILIZER SOLUTION.

- I. **CROP OIL CONCENTRATE:** A petroleum or vegetable seed based oil concentrate (such as SUN-IT II²) may be used. SUN-IT II is recommended when weeds are under moisture or temperature stress. Use SUN-IT II or other methylated seed oils at the rate of 1 to 1.5 pints per acre (use the higher rate when weeds are at the maximum label size), or use a crop oil concentrate at 2 pints per acre.

9415
1

AND

II. **LIQUID FERTILIZER SOLUTION**

Recommended nitrogen based fertilizers include liquid fertilizers (such as 28%N, 32%N or 10-34-0) at the rate of 1-2 quarts per acre. Use the higher rate when weeds are under moisture or temperature stress. Instead of a liquid fertilizer, spray grade ammonium sulfate may be used at the rate of 2.5 lbs. per acre.

This label must be in the possession of the user at the time of herbicide application.

Refer to the PRESTIGE label for additional information regarding application rates, restrictions, precautions, weeds controlled, adjuvants recommended and other information.

¹Trademark of BASF Corporation

²Trademark of Agsco, Inc.

©Registered Trademark of American Cyanamid Company

PE-xxxxx 1/96 SD

American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1996

EPA Reg. No. 241-331

10/15

FOR USE IN SOYBEANS

PURSUIT PLUS EC AND ROUNDUP™ FOR WEED CONTROL IN SOYBEANS AND PURSUIT PLUS EC USE IN ROUNDUP READY™ SOYBEANS

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PURSUIT PLUS EC LEAFLET LABEL BEFORE USING. SEE THE PURSUIT PLUS EC LEAFLET LABEL FOR PRECAUTIONS, WORKER PROTECTION STANDARD REQUIREMENTS, APPLICATION INFORMATION, USES WITH OTHER PRODUCTS AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at the time of herbicide application. The use of PURSUIT PLUS EC herbicide not consistent with this labeling may result in injury to crops. Keep container closed to avoid spills and contamination. **DO NOT** apply this product through any type of irrigation system.

GENERAL INFORMATION

PURSUIT PLUS EC provides control of many broadleaf and grass weeds when applied preplant and preemergence. PURSUIT PLUS EC can be applied to either conventional or no-till soybeans. When applied as a burndown application, Roundup (or other glyphosate containing products) may be tank-mixed with PURSUIT PLUS EC for control of weeds common to no-till production, such as mat. stail. Always include a nonionic surfactant and liquid fertilizer in the spray solution when applied to no-till soybeans.

PURSUIT PLUS EC may also be soil applied to Roundup Ready soybeans. If weeds should escape the PURSUIT PLUS EC treatment, a glyphosate containing product may be applied postemergence for weed control.

Apply glyphosate containing products postemergence only to Roundup Ready soybean varieties (these varieties contain the glyphosate resistant gene). Use only glyphosate products that are registered for use on Roundup Ready soybeans.

MIXING INSTRUCTIONS

Fill the spray tank one-fourth to one-half full with clean water or liquid fertilizer. Add the required amount of PURSUIT PLUS EC to the spray tank first and make sure it is thoroughly mixed before adding glyphosate products. When tank-mixing PURSUIT PLUS EC with Roundup, add the Roundup (or other glyphosate containing product) to the tank while agitating. Add adjuvants to the spray mixture (for no-till burndown applications). Then fill the remainder of the tank with clean water. Maintain agitation while spraying to ensure a uniform spray mixture. To avoid injury to sensitive crops, spray equipment used for PURSUIT PLUS EC herbicide applications must be drained and thoroughly cleaned with water before being used to apply other products.

USE RATE

Apply PURSUIT PLUS EC at the rate of 2 1/2 pints per acre. Refer to the glyphosate product label for recommended use rates.

NOTE: Only one application of PURSUIT PLUS EC may be made during the growing season.

(See other side)

ACCEPTED
with COMMENTS
By EPA Letter Dated

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

241-331

11/15

PURSUIT PLUS EC may be applied in conventional, minimum, or no-till soybeans as a preplant or preplant incorporated application up to 45 days before planting. DO NOT apply PURSUIT PLUS EC postemergence (or at cracking) to soybeans.

Adequate moisture is required for activation of PURSUIT PLUS EC.

PREPLANT INCORPORATED APPLICATIONS

Apply PURSUIT PLUS EC before planting and incorporate uniformly into the top 1 to 2 inches of soil.

PURSUIT PLUS EC may be applied immediately before planting or up to 45 days prior to planting. Incorporate within 7 days after application.

If soybeans are planted on beds, apply and incorporate after bed formation, using PTO-driven equipment or rolling cultivator.

SURFACE APPLICATIONS (including no-till)

Apply PURSUIT PLUS EC up to 45 days before, during or after planting but before the crop emerges. As with other herbicides applied preemergence, rainfall or irrigation is necessary to activate PURSUIT PLUS EC in the soil. If sufficient rainfall or irrigation to activate PURSUIT PLUS EC is not received within 7 days after application, a thorough shallow tillage, cultivation, or postemergence treatment may be required for control of emerged weeds.

Refer to the PURSUIT PLUS EC label and glyphosate product label for additional information regarding WEEDS CONTROLLED, MIXING AND SPRAYING INSTRUCTIONS, PRECAUTIONARY STATEMENTS AND WORKER PROTECTION INFORMATION. Refer to the glyphosate label for additional information regarding application rates, restrictions, precautions, weeds controlled, adjuvants recommended and other information.

When tank-mixing herbicides, always follow the most restrictive label.

Use of PURSUIT PLUS EC herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

®Registered Trademark of American Cyanamid Company

™Trademarks of Monsanto Agricultural Products Company

PE xxxx 12/95 SD

American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 1995©

MAR 28 1988

EPA Reg. No. 241-331

10715

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

**FOR USE IN SUCCULENT PEAS, DRY EDIBLE PEAS,
LENTILS, CHICKPEAS, AND LIMA BEANS**

PURSUIT PLUS EC can be used on succulent peas, dry edible peas (including chickpeas) lentils, and lima beans in Washington (Adams, Asotin, Columbia, Garfield, Spokane, Walla Walla and Whitman counties), Idaho (Benewah, Clearwater, Idaho, Kootenai, Latah, Lewis, and Nez Perce counties), Oregon (Union, Umatilla and Wallowa counties) and Montana only.

OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT PLUS EC LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.

GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT PLUS EC application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT PLUS EC if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT PLUS EC if cold and/or wet conditions are present or predicted to occur within one week of application

USE PURSUIT PLUS EC ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT PLUS EC herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT PLUS EC TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT PLUS EC LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT PLUS EC in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT PLUS EC.

Spray equipment used for PURSUIT PLUS EC application must be drained and thoroughly cleaned with water before being used to apply other products.

DO NOT make more than one application of PURSUIT PLUS EC per year.

DO NOT apply PURSUIT PLUS EC to sand, loamy sand, or sandy loam soils.

DO NOT apply PURSUIT PLUS EC after June 20.

Allow at least 30 days between application and harvest of succulent peas

Allow at least 60 days between application and harvest of dry edible peas, chickpeas, lentils, and lima beans

mixed with any product containing a label promoting such mixtures.

Preplant Incorporated Applications. Apply PURSUIT PLUS EC at the broadcast rate of 30 ounces per acre within 1 week before planting. Applied preplant incorporated, PURSUIT PLUS EC may be tank-mixed with other registered herbicides.

For control of the annual grass weed species listed on the PROWL® herbicide label, add 8 oz/acre of PROWL herbicide to PURSUIT PLUS EC. Do not use on sand, loamy sand, or sandy loam soils.

WEEDS CONTROLLED

PURSUIT PLUS EC applied at the broadcast rate of 30 ounces per acre preplant incorporated will control:

Buckwheat, wild	Nightshade, Eastern black
Kochia	Nightshade, hairy
Lambsquarters, common	Pigweed, redroot
Mustard, wild	Shepherdspurse
Nightshade, black	Thistle, Russian

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT PLUS EC as recommended in this label:

- Anytime:

Soybeans	Peas
Peanuts	Lima beans
Chickpeas	
Lentils	
- Four months after PURSUIT PLUS EC application:
 - Barley
 - Dry edible beans
 - Rye
 - Snap beans (three months)
 - Wheat
- Nine and one-half months after PURSUIT PLUS EC application:
 - Field corn
 - Field corn grown for seed*
 - Tobacco

*Registered trademark of American Cyanamid Company
PL XXXXX 1/5/76 SD



American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1976

5. Twenty-six months after PURSUIT PLUS EC application:

potatoes

- Forty months after PURSUIT PLUS EC application:

All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT PLUS EC application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

In the event of a crop loss, the crops listed may be replanted.

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT PLUS EC soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT PLUS EC the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT PLUS EC shall be assumed by the user.

CONSULT THE PURSUIT PLUS EC LABEL FOR OTHER PRECAUTIONS.

EPA Reg. No. 241-331

10/15
1

**FOR USE IN NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE,
CRANBERRY, AND SMALL WHITE TYPE DRY BEANS ONLY**

USE ONLY IN MICHIGAN (EXCLUDING UPPER PENINSULA)

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN
THE PURSUIT PLUS EC LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a PURSUIT PLUS EC application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply PURSUIT PLUS EC if planting is delayed and chance of frost prior to maturity is likely.

DO NOT apply PURSUIT PLUS EC if cold and/or wet conditions are present or predicted to occur within one week of application.

USE PURSUIT PLUS EC ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of PURSUIT PLUS EC herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF PURSUIT PLUS EC TO EDIBLE LEGUME VEGETABLE CROPS.

REFER TO THE PURSUIT PLUS EC LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

DO NOT apply by air.

DO NOT apply PURSUIT PLUS EC in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with PURSUIT PLUS EC.

Spray equipment used for PURSUIT PLUS EC application must be drained and thoroughly cleaned with water before being used to apply other products.

DO NOT make more than one application of PURSUIT PLUS EC per year.

DO NOT apply to Domino variety black turtle beans

DO NOT apply PURSUIT PLUS EC to sand or loamy sand soils

DO NOT apply PURSUIT PLUS EC after June 20

Allow at least 60 days between application and harvest

**ACCEPTED
with COMMENTS
In EPA Letter Dated**

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
241-331

*Registered Trademark of American Cyanamid Company

(See reverse side)

Soil Type

Use Rate (oz/acre)

Soil Type	Use Rate (oz/acre)
Sands, loamy sands	DO NOT USE
Sandy loams and all other soils with $\leq 2\%$ O.M.	20 (1.25 pts)
All other soils with $> 2\%$ O.M.	30

For control of the annual grass weed species listed on the PROWL[®] herbicide label, add 0.75 to 1 pt/acre of PROWL herbicide to the 20 oz/acre rate of PURSUIT PLUS EC, and 0.25 to 0.5 pt/acre of PROWL to the 30 oz/acre rate of PURSUIT PLUS EC.

WEEDS CONTROLLED

PURSUIT PLUS EC applied to low organic matter soils at the broadcast rate of 20 ounces per acre preplant incorporated will control light to moderate infestations of:

Nightshade, black
Nightshade, Eastern black
Pigweed, redroot

PURSUIT PLUS EC applied at the broadcast rate of 30 ounces per acre preplant incorporated will control:

Mustard, wild
Nightshade, black
Nightshade, Eastern black
Nightshade, hairy
Pigweed, redroot

ROTATIONAL CROP GUIDELINE

The following crops may be planted after applying PURSUIT PLUS EC as recommended in this label:

- Anytime:
 - Soybeans
 - Peanuts
 - Edible bean types listed on this label (Refer to heading on page 1)
- Four months after PURSUIT PLUS EC application
 - Barley
 - Edible bean types not listed on this label (Refer to heading on page 1)
 - Peas
 - Wheat

Cotton
Lettuce
Popcorn
Safflower
Sunflower
Sweet Corn

- Twenty-six months after PURSUIT PLUS EC application:
 - potatoes
- Forty months after PURSUIT PLUS EC application:
 - All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE. Following forty months after a PURSUIT PLUS EC application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

In the event of a crop loss, the bean crop may be replanted

DO NOT feed treated forage, hay, or straw to livestock.

* Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT PLUS EC soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT PLUS EC the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT PLUS EC shall be assumed by the user.

CONSULT THE PURSUIT PLUS EC LEAFLET LABEL FOR OTHER PRECAUTIONS.

[®]Registered Trademark of American Cyanamid Company
E-12136 Rev. 4-1-79



American Cyanamid Company
Agricultural Products Division
Crop Protection Chemical Department
Wayne, NJ 07090