

PM 05

241-331

page 1 of 26

PURSUIT® PLUS EC
herbicide

FOR USE IN SOYBEANS AND FIELD CORN*

***(APPLY ONLY TO IMI-CORN™ HYBRIDS)**

ACTIVE INGREDIENTS:

Imazethapyr (±)-2-[4,5-dihydro-4-methyl-4-(1-methyl-ethyl)-5-oxo-1H-imidazol-2-yl]-5-ethyl-3-pyridinecarboxylic acid	2.24%
pendimethalin (N-1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine	30.24%
INERT INGREDIENTS*	67.52%
TOTAL	100.00%

PURSUIT PLUS EC contains 2.9 pounds of active ingredients per gallon (2.7 pounds ai of pendimethalin and 0.2 pounds acid equivalent of imazethapyr).

*Contains petroleum distillates

Patent No. 4,798,619
EPA Reg. No. 241-331

ACCEPTED
W/ COMMENTS
In EPA Letter Dated
OCT 10 1996

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. EPA Est. No.

KEEP OUT OF REACH OF CHILDREN

241-331

CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand this label, find someone to explain it to you in detail).

STATEMENT OF PRACTICAL TREATMENT

- IF SWALLOWED:** Do not induce vomiting. Call a physician or Poison Control Center immediately.
- IF IN EYES:** Flush eyes with plenty of water. Call a physician if irritation persists.
- IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.

NOTE TO PHYSICIAN: Because of increased risk of chemical pneumonia or pulmonary edema caused by aspiration of the hydrocarbon solvent, vomiting should be induced only under professional supervision.

In case of an emergency endangering life or property involving this product, call collect, day or night, area code 201-835-3100.

See Next Page for Additional Precautionary Statements

American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1996

Net Contents: 2.5 gallons
(9.45 liters)

®Registered Trademark of American Cyanamid Company
™Trademark of American Cyanamid Company

REC'D EPA/DPP/DPD1
96 SEP 27 P 2:19

TABLE OF CONTENTS

Ingredients	cover
Statement of Practical Treatment.....	cover
Precautionary Statements.....	3
Environmental Hazards.....	3
Directions for Use	3
Agricultural Use Requirements.....	4
Storage and Disposal.....	4
Disclaimer	5
General Information.....	6
Mixing Instructions.....	7
Spraying Instructions	8
Ground Applications	8
Aerial Applications	10
Soil Application Instructions.....	10
Preemergence (Surface Applications).....	11
Surface Applications Made Before Planting	11
Surface Applications Made After Planting (Field Corn).....	11
Surface Applications Made After Planting (Soybeans).....	11
No-Till or Reduced Tillage (Soybeans).....	11
Preplant Incorporated Applications (Soybeans only)	12
Application Instructions - Postemergence (Field corn only).....	12
Herbicide Use Area	13
Use Rate	13
IMI-CORN - Directions For Use	14
Tank-Mix Herbicide Combinations With PURSUIT PLUS EC.....	14
Weeds Controlled.....	14
Soybeans - Directions For Use.....	17
Herbicide Combinations.....	17
PURSUIT PLUS EC Followed by STATUS™ herbicide	17
PURSUIT PLUS EC Followed by PRESTIGE™ herbicide.....	17
PURSUIT PLUS EC + SCEPTER® 70DG Tankmixture	19
PURSUIT PLUS EC Plus Metribuzin herbicide	20
PURSUIT PLUS EC Plus 2,4-D.....	21
Fall Applications (South Dakota).....	21
Weeds Controlled	21
Rotational Crop Guideline	24
Exception to Rotational Crop Guideline.....	25
Precautions	25

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)**

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing.

Personal Protective Equipment (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate or viton \geq 14 mils
- Shoes plus socks.

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. DO NOT apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. DO NOT contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

This labeling must be in the possession of the user at the time of pesticide application.

Observe all cautions and limitations in this leaflet. Do not use PURSUIT PLUS EC other than in accordance with the instructions set forth on this label. The use of PURSUIT PLUS EC not consistent with this label can result in injury to crops. Keep container closed to avoid spills and contamination.

DO NOT apply this product through any type of irrigation system.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Exception: if the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical resistant gloves, such as barrier laminate or viton \geq 14 mils
- Shoes plus socks

STORAGE AND DISPOSAL

STORAGE: KEEP FROM FREEZING. DO NOT STORE BELOW 40°F. SHAKE WELL BEFORE USING.

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL FOR 2.5 GALLONS: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

CONTAINER DISPOSAL FOR MINI-BULK AND BULK: Return empty container to point of purchase.

DISCLAIMER

The label instructions for the use of this product reflect the opinion of experts based on research and field use. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the use of, or application of the product contrary to label instructions, all of which are beyond the control of American Cyanamid Company. All such risks shall be assumed by the user.

American Cyanamid Company shall not be responsible for losses or damages resulting from use of this product in any manner not set forth on this label. User assumes all risks associated with the use of this product in any manner not specifically set forth on this label.

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use, subject to the risks referred to above. **CYANAMID DOES NOT MAKE OR AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED AND EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.**

BUYER'S EXCLUSIVE REMEDY AND AMERICAN CYANAMID'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF PURSUIT® PLUS EC herbicide. In no case shall Cyanamid or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product.

USES WITH OTHER PRODUCTS (TANK-MIXES)

If this product is used in combination with any other product except as specifically recommended in writing by American Cyanamid Company then American Cyanamid Company shall have no liability for any loss, damage or injury arising out of its use in any such combination not so specifically recommended. If used in combination recommended by American Cyanamid Company, the liability of American Cyanamid Company shall in no manner extend to any damage, loss or injury not directly caused by the inclusion of the American Cyanamid Company product in such combination use, and in any event shall be limited to return of the account of the purchase price of the American Cyanamid Company product.

GENERAL INFORMATION

PURSUIT PLUS EC is effective in providing weed control in conservation tillage systems. PURSUIT PLUS EC may be applied in minimum tillage, no-fill, or conventional tillage to field corn (IMI-CORN hybrids) or soybeans.

Field Corn: Apply PURSUIT PLUS EC only on selected field corn hybrids (IMI-CORN) warranted by the seed company to possess resistance/tolerance to direct application of PURSUIT® herbicide. DO NOT apply PURSUIT PLUS EC to corn hybrids which lack genetic resistance/tolerance to PURSUIT herbicide. Contact your seed supplier, chemical dealer or American Cyanamid to obtain information regarding IMI-CORN hybrids. With the exception of reduced tillage systems (see No-till or Reduced Tillage section) plant into a firm seed bed that is free of clods and crop residue. Use only where adequate tillage is practiced to provide good soil coverage of the corn seed. Plant corn AT LEAST 1-1/2 inches deep to ensure good seed coverage. Make sure furrows close to prevent seed contact with the herbicide. Wait at least 7-10 days before cultivating postemergence treatments.

Crops growing under stressful environmental conditions can exhibit various injury symptoms which may be more pronounced if herbicides are used. Stunting, leaf curling or temporary yellowing of the field corn plants may occur following PURSUIT PLUS EC applications. Normal growth and appearance should resume within 1 to 2 weeks.

Soybeans: Apply PURSUIT PLUS EC preplant or preplant incorporated from 45 days prior to planting up to planting. Incorporate within 7 days of application if rainfall is not received.

Occasionally, internode shortening and/or temporary yellowing of soybean plants may occur following PURSUIT PLUS EC applications. This will not affect soybean yields.

After PURSUIT PLUS EC is applied, some susceptible weeds emerge, growth stops, and the weeds either die or are not competitive with the crop.

PURSUIT PLUS EC kills weeds by herbicide uptake by weed roots and rapid translocation to the growing points. Therefore, adequate soil moisture is important for optimum PURSUIT PLUS EC activity. When adequate soil moisture is present, PURSUIT PLUS EC will provide residual control of susceptible germinating weeds.

Use of PURSUIT PLUS EC herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

Naturally occurring biotypes* of some of the weeds listed on this label may not be effectively controlled by this and/or other products with either the ALS/AHAS enzyme inhibiting mode of action or the mitotic inhibiting mode of action. Other herbicides with the ALS/AHAS enzyme inhibiting mode of action include the sulfonylureas (e.g., Accent™, Basic™, Classic™, Concert™, Exceed™, Permit™, Pinnacle™, etc.), the sulfonamides (e.g., Broadstrike™, etc.) and the pyrimidyl benzoates (e.g. Staple™, etc.). Herbicides with the mitotic inhibiting mode of action include the other dinitroaniline herbicides such as PROWL® herbicide, TREFLAN™ herbicide TRI-4® herbicide and Sonolan™. If naturally occurring ALS/AHAS resistant biotypes are present in a field, PURSUIT PLUS EC and/or any other ALS/AHAS enzyme inhibiting mode of action herbicide should be tank-mixed or applied sequentially with an appropriate registered herbicide having a different mode of action to ensure control.

*A weed biotype is a naturally occurring plant within a given species that has a slightly different, but distinct, genetic makeup from other plants.

Replanting: If replanting is necessary in a field previously treated with PURSUIT PLUS EC, the field may be replanted to soybeans, peanuts, lima beans or southern peas. Rework the soil no deeper than the treated zone. IMI-CORN may also be replanted, but do not rework the soil. Do not apply a second treatment of PURSUIT PLUS EC.

MIXING INSTRUCTIONS

Fill the spray tank one-fourth to one-half full with clean water. While agitating add the required amount of product and then fill the remainder of the tank with clean water. Maintain agitation while spraying to ensure a uniform spray mixture.

When tank-mixing PURSUIT PLUS EC with recommended herbicides, add the other herbicides and adjuvants in the following order while agitating.

1. Fill spray tank 1/4 to 1/2 full with clean water.
2. Add soluble packet products and thoroughly mix.
3. Add WP (wetttable powder), DG (dispersible granule), DF (dry flowable) or LF (liquid flowable) formulations.
4. Add aqueous solution products.
5. Add PURSUIT PLUS EC
6. Add other EC (emulsifiable concentrate) products.
7. Add surfactant to the spray tank (postemergence applications to IMI-CORN only).
8. Add liquid fertilizer.
9. While agitating, fill the remainder of the tank with water.

When paraquat is included in a tank-mixture, add 8 ounces of non-ionic surfactant per 100 gallons of spray mixture as the last ingredient in the tank.

To avoid injury to sensitive crops, spray equipment used for PURSUIT PLUS EC applications must be drained and thoroughly cleaned with water before being used to apply other products.

8 7 26

SPRAYING INSTRUCTIONS

DO NOT apply when wind velocity is greater than 10 mph, or when spray may be carried to sensitive crops. Sensitive crops include leafy vegetables, sugarbeets, and cotton.

GROUND APPLICATIONS:

Uniformly apply with properly calibrated ground equipment in 10 to 40 gallons of water per acre. A spray pressure of 20 to 40 psi is recommended.

Applications With Liquid Fertilizer:

PURSUIT PLUS EC can be applied to the soil in liquid fertilizers, alone or in combination with PROWL 3.3 EC, trifluralin (soybeans only), Lasso™ or Dual™. Follow all PURSUIT PLUS EC label recommendations regarding incorporation, timing of application, special instructions and precautions. Apply treatments in 20 or more gallons of liquid fertilizer per acre with ground equipment. Always test the compatibility of PURSUIT PLUS EC with liquid fertilizer before mixing in the spray tank.

Applications With Dry Bulk Fertilizer:

PURSUIT PLUS EC may be impregnated on dry bulk fertilizers. When applied as directed, PURSUIT PLUS EC/dry bulk fertilizer mixtures provide weed control equal to that provided by the same rates of PURSUIT PLUS EC applied in water or liquid fertilizer.

Follow all PURSUIT PLUS EC label recommendations regarding application and incorporation, special instructions and precautions. Apply PURSUIT PLUS EC/dry bulk fertilizer mixtures only with ground equipment.

All individual state regulations relating to dry bulk fertilizer, registration, labeling, and application are the responsibility of the individual and/or company selling the PURSUIT PLUS EC/dry bulk fertilizer mixtures.

A minimum of 200 pounds and a maximum of 450 pounds of dry bulk fertilizer impregnated with the recommended amount of PURSUIT PLUS EC must be applied per acre.

DO NOT impregnate PURSUIT PLUS EC onto coated ammonium nitrate or limestone because these materials will not absorb the herbicide. Dry fertilizer blends containing mixtures of ammonium nitrate or limestone may be impregnated with PURSUIT PLUS EC. A minimum of 200 pounds of impregnated dry bulk fertilizer, excluding the weight of ammonium nitrate or limestone, must be applied per acre.

Apply PURSUIT PLUS EC at the rate of 2-1/2 pints per acre. Use the following table to determine the amount of PURSUIT PLUS EC to be impregnated on a ton of dry bulk fertilizer based on the rate of fertilizer which will be applied per acre.

9 8 26

RATE CHART FOR IMPREGNATION OF DRY BULK FERTILIZER WITH PURSUIT PLUS EC
(Pints of PURSUIT PLUS EC per Ton of Fertilizer)

<u>PURSUIT PLUS EC Rate Per Acre</u>	<u>Fertilizer Rate pounds per acre</u>	<u>Pints per Ton</u>
2-1/2 pints	200	25
	250	20
	300	16-2/3
	350	14-1/3
	400	12-1/2
	450	11

For those rates not listed in this table, calculate the pints of PURSUIT PLUS EC to be impregnated on a ton of dry bulk fertilizer using the following formula:

$$\frac{2000}{\text{Pounds of dry fertilizer per acre}} \times \begin{matrix} 2-1/2 \text{ Pints of} \\ \text{PURSUIT PLUS EC} \\ \text{per acre} \\ \text{(recommended rate)} \end{matrix} = \begin{matrix} \text{Pints of} \\ \text{PURSUIT PLUS EC} \\ \text{per ton of fertilizer} \end{matrix}$$

To impregnate PURSUIT PLUS EC on bulk fertilizer, use a closed rotary-drum mixer or other commonly used dry bulk fertilizer blender equipped with suitable spray equipment. Spray nozzles must be placed to provide uniform coverage of PURSUIT PLUS EC onto the fertilizer during mixing.

If PROWL 3.3 EC is to be combined with the PURSUIT PLUS EC prior to impregnation, premix the PROWL 3.3 EC with an equal volume of water before adding it to the PURSUIT PLUS EC. DO NOT mix undiluted PROWL 3.3 EC with PURSUIT PLUS EC.

Apply the PURSUIT PLUS EC/dry bulk fertilizer mixture with an accurately calibrated dry fertilizer spreader. The PURSUIT PLUS EC/dry bulk fertilizer mixture must be spread uniformly on the soil surface. Uneven spreading can cause poor weed control and crop injury.

Refer to PREPLANT INCORPORATED APPLICATIONS section of this label for incorporation directions.

AERIAL APPLICATIONS:

Uniformly apply with properly calibrated aerial equipment in 5 or more gallons of water per acre. When applied POSTEMERGENCE, (to IMI-CORN HYBRIDS ONLY) the addition of a non-ionic surfactant AND fertilizer are required for optimum weed control. Add a non-ionic surfactant at the rate of 1 quart per 100 gallons of spray mixture AND a liquid fertilizer at the rate of 1 quart per acre. (See instructions under APPLICATIONS INSTRUCTIONS - POSTEMERGENCE).

To avoid injury to sensitive crops from drift, aerial applicators must adhere to the following SPECIAL AERIAL USE DIRECTIONS AND PRECAUTIONS:

- Use nozzles which produce a coarse spray.
- Nozzle height above ground must be a maximum of 10 feet.
- Nozzles must be pointed toward the rear of the aircraft. The downward angle of the nozzle should not be greater than 20 degrees.
- To minimize wing-tip vortex roll, nozzles or spray boom must not be located any closer to end of wing or rotor than three-fourths the distance from the center of the aircraft.
- Use a maximum spray pressure of 40 psi.
- A buffer zone must be established between the area to be sprayed and sensitive crops.
- DO NOT spray when wind velocity is greater than 5 mph.

Applicator is responsible for any loss or damage which results from spraying PURSUIT PLUS EC in a manner other than recommended on this label. In addition, applicator must follow all applicable state and local regulations and ordinances in regard to spraying.

SOIL APPLICATION INSTRUCTIONS

PURSUIT PLUS EC provides effective weed control in conservation tillage systems designed to meet conservation compliance requirements. PURSUIT PLUS EC can be applied preemergence or postemergence treatment to IMI-CORN hybrids. (DO NOT apply preplant incorporated to IMI-CORN). It can be applied early preplant, preplant incorporated or preemergence in soybeans (DO NOT apply postemergence to soybeans). It can also be applied in conventional, minimum tillage and no-till production systems. The application method of choice will depend on the anticipated weed spectrum and the preference of the applicator. If weeds have emerged, see instructions under NO-TILL OR REDUCED TILLAGE section.

PURSUIT PLUS EC controls weeds by uptake by weed roots, and translocation to the growing points where it stops weed growth.

Adequate soil moisture is required for optimum activity. For surface applications, rainfall or overhead irrigation is necessary to move PURSUIT PLUS EC into the weed germination zone. The amount of rainfall or irrigation required following application depends on existing soil moisture, soil texture and organic matter content. Sufficient water to moisten the soil to a depth of 2 inches is normally adequate. If adequate moisture is not received within 7 days after a surface-applied treatment, then a cultivation is recommended to control escaped weeds. In no-till situations where cultivation is not practical, a postemergence application is required to control escaped weeds. When adequate moisture is received after dry conditions, PURSUIT PLUS EC will provide residual control of susceptible germinating weeds; activity on established weeds will depend on the weed species and the location of its root system in the soil.

In ridge-till plantings, PURSUIT PLUS EC may be applied early preplant (soybeans only) or preemergence. If the herbicide is banded over the row, cultivation will be required for weed control between the beds. If cultivation is not possible or if weed pressure is heavy, apply PURSUIT PLUS EC in a broadcast application. Use proportionally less PURSUIT PLUS EC per acre in a band versus a broadcast application. If rainfall does not occur within 7 days of application, a rotary hoe incorporation will enhance weed control. See PREEMERGENCE APPLICATIONS and PREPLANT INCORPORATED APPLICATIONS for further information.

PREEMERGENCE APPLICATIONS

SURFACE APPLICATIONS MADE BEFORE PLANTING: (Soybeans)

PURSUIT PLUS EC may be surface applied prior to soybean planting (up to 45 days). If sufficient rain does not occur before planting to activate PURSUIT PLUS EC, shallow incorporation before planting will enhance weed control. (PURSUIT PLUS EC may be surface applied prior to soybean planting both north and south of highway I-80).

SURFACE APPLICATIONS MADE AFTER PLANTING: (IMI-CORN)

Apply PURSUIT PLUS EC alone preemergence (surface treatment only) after planting. DO NOT INCORPORATE PURSUIT PLUS EC or corn injury may result.

NOTE: Plant corn at least 1-1/2 inches deep. Adjust planters to ensure adequate seed coverage.

The use of no-till planters in minimum tillage corn under conditions which do not allow good soil coverage of the corn seed can result in reduced crop stand or injury if PURSUIT PLUS EC contacts the germinating corn seed. Check equipment to ensure good seed coverage.

SURFACE APPLICATIONS MADE AFTER PLANTING: (Soybeans)

PURSUIT PLUS EC may be surface applied up to 2 days after soybean planting (before crop emergence) South of Interstate Highway I-80 only. DO NOT APPLY PURSUIT PLUS EC AFTER SOYBEAN PLANTING North of Interstate Highway I-80.

NO-TILL OR REDUCED TILLAGE: (Soybeans)

PURSUIT PLUS EC is effective in controlling weeds in conservation tillage production systems. Apply PURSUIT PLUS EC treatments up to 45 days prior to planting (early preplant) but before crop emergence.

To ensure thorough coverage, use a minimum of 20 gallons of water per acre. Use higher gallonage for fields with dense vegetation or heavy crop residues. Adjust the boom height to ensure proper coverage of weed foliage (according to the manufacturers recommendation). The nozzle spacing on the boom should be 20 inches. Use only standard flat-fan nozzle tips. Use ground equipment only.

PURSUIT PLUS EC will control emerged problem weeds such as velvetleaf, smartweed, common cocklebur, and pigweeds. The weed size should not exceed 3" for velvetleaf and smartweed, and 8" for cocklebur and pigweeds. Add surfactant to the spray mixture at the rate of 1 qt./100 gal. AND a nitrogen based fertilizer (such as 28%N, 32%N or 10-34-0) at the rate of 1-2 qt./acre for optimum activity. Ammonium sulfate (spray grade) may be substituted for liquid fertilizer at the rate of 2.5 lbs. per acre.

If other vegetation is present (not controlled by PURSUIT PLUS EC), apply paraquat, Roundup™ or 2,4-D in combination with PURSUIT PLUS EC. Refer to the respective labels for rates, methods of application, proper timing, weeds controlled, restrictions, and precautions. Use ground equipment only.

12 9 26

PREPLANT INCORPORATED APPLICATIONS (Soybeans only)

PURSUIT® PLUS EC herbicide may be applied following land preparation and should be thoroughly incorporated to a depth of 1 to 2 inches. When applied to beds, maintain PURSUIT PLUS EC in the surface 1 to 2 inches of the finished beds. Application may be made up to 45 days prior to planting (early preplant). Incorporate prior to soybean planting and within 7 days of application.

If soybeans are planted on beds, apply and incorporate after bed formation using PTO-driven equipment or a rolling cultivator. For optimum weed control, PURSUIT PLUS EC should be maintained in the surface 1-2 inches of the finished bed.

APPLICATION INSTRUCTIONS - POSTEMERGENCE (IMI-CORN)

Apply PURSUIT PLUS EC herbicide as a postemergence treatment to IMI-CORN when crop and weeds are actively growing. Apply PURSUIT PLUS EC before weeds exceed a height of 3", unless otherwise indicated. **More restrictive crop growth stage limitations of tank-mix partners MUST be followed.**

RESTRICTIONS FOR POSTEMERGENCE APPLICATIONS:

To avoid serious crop injury with postemergence applications observe the following precautions:

- DO NOT apply PURSUIT PLUS EC with a liquid fertilizer as a carrier.
- DO NOT apply PURSUIT PLUS EC when air temperatures are expected to reach or stay below 40° F for 10 or more hours or when extended cold, wet conditions are predicted.
- DO NOT apply PURSUIT PLUS EC postemergence to soybeans.

ADDITIVES:

POSTEMERGENCE APPLICATIONS OF PURSUIT PLUS EC REQUIRE THE ADDITION OF A SURFACTANT AND A LIQUID FERTILIZER/SOLUTION.

I. SURFACTANTS

Use a non-ionic surfactant containing at least 80% active ingredient. Apply the surfactant at the rate of 1 quart per 100 gallons of spray solution.

AND

II. LIQUID FERTILIZER SOLUTION¹

Recommended nitrogen based fertilizers include liquid fertilizers (such as 28%N, 32%N, or 10-34-0) at the rate of 1-2 quarts per acre. Use the higher rate when weeds are under moisture or temperature stress. Instead of a liquid fertilizer, spray grade ammonium sulfate may be used at the rate of 2.5 lbs per acre.

¹PURSUIT PLUS EC applications may be made with a non-ionic surfactant only. (Liquid fertilizer not required in the states of Arkansas, Bootheel of Missouri, Tennessee, South Carolina, Louisiana, Mississippi, Alabama and Georgia.)

13 7 26

When STATUS herbicide is applied following an application of PURSUIT PLUS EC, a surfactant OR a crop oil concentrate AND a nitrogen-based fertilizer must be added to the spray solution.

When PRESTIGE herbicide is applied following an application of PURSUIT PLUS EC, a crop oil concentrate and a nitrogen-based fertilizer must be added to the spray solution.

PURSUIT PLUS EC HERBICIDE USE AREA

Do not apply PURSUIT PLUS EC to soybeans or IMI-CORN in North Dakota or in Minnesota north of state highway 210.

USE RATE (2-1/2 PINTS PER ACRE)

Apply PURSUIT PLUS EC at a broadcast rate of 2 1/2 pints per acre (to soybeans or IMI-CORN hybrids) for all methods of application: preplant, preplant incorporated (soybeans only) and preemergence (including minimum and no-till) or postemergence (IMI-CORN hybrids only). At this broadcast rate, one gallon of PURSUIT PLUS EC will treat 3.2 acres.

NOTE: Only one application of PURSUIT PLUS EC may be made during the season.

CROPS

IMI-CORN - DIRECTIONS FOR USE

TANK-MIX HERBICIDE COMBINATIONS WITH PURSUIT PLUS EC

Preemergence Applications of PURSUIT PLUS EC Tank-Mixed With:

Atrazine	Frontier™
Bicep™, Bicep Lite™	Lariat™
Banvel™	Harness Plus™
Bullet™	Lasso EC, Lasso MT
Clarity™	Marksman™
Dual	PROWL 3.3 EC
Eradicane™ Extra	Surpass™
Extrazine™ II	Sutan+™

Postemergence Applications of PURSUIT PLUS EC Tank-Mixed With:

Atrazine ^{1,2}	Clarity ^{1,3}
Banvel ^{1,3}	Marksman ¹
Basagran™,1	PROWL 3.3 EC
Buctril™,1,2	Sencor DF™,Sencor
2,4-D	3L™

When PURSUIT PLUS EC herbicide is used in combination with another herbicide, refer to the respective labels for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions.

WEEDS CONTROLLED

When applied as directed, PURSUIT PLUS EC will control or reduce competition from the weeds listed below.

NOTE: C = Control, R = Reduced Competition

¹ In some cases the grass activity of PURSUIT PLUS EC will be reduced when used in combination with atrazine, Buctril, Banvel, Basagran, Clarity, or Marksman postemergence.

² Some corn leaf burn may result with Buctril or atrazine postemergence combinations with PURSUIT PLUS EC

³ Applications of Banvel or Clarity herbicide to corn during periods of rapid growth may result in temporary leaning.

15 9 26

BROADLEAF WEEDS

Weeds Controlled	PREEMERGENCE	POSTEMERGENCE	
		Maximum Leaf Stage***	Size (Inches)
Alligator weed		4	1-3
Anoda, spurred	C	2	1-2
Artichoke, Jerusalem		8	6-10
Buffalobur	C*	R	1-3
Carpetweed	C		
Cocklebur, common	R	8	1-8
Galinsoga	C		
Jimsonweed	C*	4	1-3
Kochia	C	4	1-3
Lambquarters, common	C*	R	1-2
Mallow, Venice	C		
Marshelder	C	4	1-3
Morningglory			
entireleaf	R	2	1-2
ivyleaf	R	2	1-2
pitted	R	2	1-2
smallflower	C	4	1-3
tall	R	2	1-2
Mustard species	C	4	1-3
Nightshade			
black	C	4	1-3
Eastern black	C	4	1-3
hairy	C	4	1-3
Pigweed			
palmer	C		
redroot	C	8	1-8
smooth	C	8	1-8
spiny	C	8	1-8
Poinsettia, wild	C		
Puncturevine	C		
Purslane, common	C		
Pusley, Florida	C		
Ragweed,			
common	R	4	1-3
giant	R	4	1-3
Sage, barnyard	R	R	1-3
Sida, prickly (teaweed)	C*		
Smartweed			
ladysthumb	C	4	1-3
Pennsylvania	C	4	1-3
Spurge			
prostrate	C	4	1-3
spotted	C	4	1-3
Sunflower	C*	4	1-3
Thistle, Canada		R	1-3
Velvetleaf	C	4	1-3
Waterhemp, tall**	C		

GRASS WEEDS

Weeds Controlled	PREEMERGENCE	POSTEMERGENCE	
		Maximum Leaf Stage***	Size (Inches)
Barnyardgrass	R	3	1-3
Crabgrass			
large	C	3	1-3
smooth	C	3	1-3
Crowfootgrass	C		
Cupgrass, woolly	C	3	1-3
Foxtail			
giant	C	6	1-6
green	C	3	1-3
yellow	C	3	1-3
Goosegrass	R		
Johnsongrass			
seedling	C	6	1-8
rhizome		R	1-8
Millet, wild proso	R		
Panicum			
browntop	C		
fall	C		
Texas	C		
Red rice		3	1-3
Sandbur, field	C		
Shattercane	R	6	1-8
Signalgrass, broadleaf	R	4	1-8
Sorghum almum	R		
Witchgrass...	C	6	1-3
SEDGES			
Nutsedge			
yellow	R	R	1-3
purple	R	R	1-3

* Cultivation and/or a postemergence herbicide may be required for season-long control.

** If a heavy infestation of waterhemp species is anticipated a tank mixture of PURSUIT PLUS EC plus additional PROWL 3.3EC is required for control. Add PROWL 3.3EC to the PURSUIT PLUS EC mixture at the following rates, depending on soil type:

Coarse textured soils: add PROWL 3.3EC at 0.6 pints/acre.

Medium texture soils: add PROWL 3.3EC at 1.2 to 1.8 pints/acre.

Fine texture soils: add PROWL 3.3EC at 1.8 pints/acre.

Refer to the PROWL 3.3EC label for specific use rates, application methods and application timings based on soil texture and soil organic matter content. A postemergence herbicide such as Blazer, Cobra or Reflex may be needed to control waterhemp species escapes. Refer to individual product labels for specific uses and recommendations.

*** The number under Maximum Leaf Stage indicates the MAXIMUM number of leaves at which weeds should be sprayed postemergence). DO NOT count cotyledon leaves when determining weed stage of growth.

SOYBEANS - DIRECTIONS FOR USE

HERBICIDE COMBINATIONS

PURSUIT PLUS EC FOLLOWED BY STATUS™ HERBICIDE

When soil applied for weed control in soybeans, PURSUIT PLUS EC controls many problem broadleaf and grass weeds. Under adverse weather conditions or when heavy weed populations exist, some problem weeds may not be completely controlled by PURSUIT PLUS EC. STATUS herbicide provides fast postemergence burn-down of susceptible broadleaf weeds. Temporary bronzing of the soybean leaves may occur soon after application. Soybeans soon outgrow early symptoms; yield will not be effected by applying STATUS following an application of PURSUIT PLUS EC.

Weeds Controlled

A postemergence application of STATUS herbicide following PURSUIT PLUS EC at the recommended rates will control common and giant ragweeds, pigweed species and waterhemp. (Refer to the STATUS label for additional weeds controlled).

Application Rates

Sequential Application:

When PURSUIT PLUS EC is soil applied, followed by STATUS applied postemergence, apply PURSUIT PLUS EC at the rate of 2 1/2 pints per acre. Apply STATUS postemergence at the rate indicated in the table below, depending on weed size:

Weeds/Weed Size	STATUS Rate (ounces per acre):*		
	10 - 12 oz.	14 - 16 oz.	18 - 24 oz.
Common ragweed Pigweed species, Waterhemp, tall common	1 - 4"	4 - 6"	6 - 8"
Giant Ragweed		1 - 6"	6 - 8"***

*Use the higher rate if common ragweed is present or the weed population is high.

**Use the 24 ounce/acre rate if giant ragweed is 6 - 8 inches tall.

Refer to the STATUS label for additional information regarding application rates, restrictions, precautions, weeds controlled, adjuvants recommended and other information.

PURSUIT PLUS EC FOLLOWED BY PRESTIGE™ HERBICIDE FOR ENHANCED GRASS CONTROL.

Under adverse weather conditions or when heavy weed populations exist, some problem weeds may not be completely controlled by PURSUIT PLUS EC. PRESTIGE herbicide provides postemergence control of many grass weeds. Refer to the table below for a partial list of weeds and the recommended use rate.

Use Rates

Apply PURSUIT PLUS EC at the broadcast rate of 2 and 1/2 pints per acre for all methods of application: preplant, preplant incorporated, and preemergence. At this broadcast rate, one gallon of PURSUIT PLUS EC will treat 3.2 acres.

Weeds Controlled

A postemergence application of PRESTIGE herbicide following PURSUIT PLUS EC at the recommended rates will control the grasses listed below. (Refer to the PRESTIGE label for additional weeds controlled).

Refer to the following table for the recommended rate of PRESTIGE.

WEEDS CONTROLLED
(PRESTIGE Herbicide following PURSUIT PLUS EC)

PRESTIGE Rate and Maximum Weed Height at Application

Grass	Early Postemergence		Mid - Postemergence		Late Postemergence	
	Max. Ht. (Inches)	Rate per Acre (oz.)	Max. Ht. (Inches)	Rate per Acre (oz.)	Max. Ht. (Inches)	Rate per Acre (oz.)
Barnyardgrass	4	18	8	24	12	36
Crabgrass, Large	-	-	6	24	8	36
Smooth	-	-	6	24	8	36
Cupgrass, Woolly	-	-	8	24	-	-
Foxtail, Giant	4	18	8	24	16	36
Green	4	18	8	24	16	36
Yellow	-	-	8	24	16	36
Goosegrass	3	18	6	24	8	36
Johnsongrass Seedling	-	-	8	24	16	36
Millet, Wild Proso	10	12	10	12	24	24
Panicum, Browntop	-	-	8	24	-	-
Fall	4	18	8	24	12	36
Texas	4	18	8	24	12	24
Shattercane	-	-	18	24	-	-
Signalgrass Broadleaf	4	18	8	24	12	36
Volunteer, Barley	-	-	4	36	-	-
Corn	12	18	20	24	-	-
Oats	-	-	4	36	-	-
Rye	-	-	4	36	-	-
Wheat	-	-	4	36	-	-
Witchgrass	-	-	8	24	-	-

*If a mixture of grasses are present, use the highest rate indicated for the grasses present.

Refer to the PRESTIGE label for additional information regarding application rates, restrictions, precautions, weeds controlled, adjuvants recommended and other information.

PURSUIT PLUS EC + SCEPTER® 70DG FOR ENHANCED WEED CONTROL

For control of a broad-spectrum of weeds in soybeans, SCEPTER® 70DG herbicide may be applied tank-mixed with PURSUIT PLUS EC in soil applications only.

Use Area

The application of PURSUIT PLUS EC and SCEPTER 70DG may be used in Region 2 as described in the USE AREA of the SCEPTER 70DG label and the entire state of Illinois.

The approved USE AREA (Region 2) includes eastern Kansas (east of highway U.S. 81; the counties of Cloud, Ellsworth, Harvey, Jewell, Lincoln, Mitchell, McPherson, Ottawa, Republic, Saline, Sedgewick, and Sumner), southeastern Nebraska (east of U.S. 81, south of U.S. 34), Missouri, Illinois, Indiana, Ohio, Michigan, Kentucky, Virginia, West Virginia, Pennsylvania, Maryland, Delaware and New Jersey.

Application Rates

Apply PURSUIT PLUS EC at the rate of 2 1/2 pints per acre.

Apply SCEPTER 70DG at the rate of 1.4 ounces per acre. (One (1) 14 ounce bag of SCEPTER 70DG treats 10 acres. Apply in tank-mixture with PURSUIT PLUS EC in soil applications.)

For application timing, refer to the PURSUIT PLUS EC and SCEPTER 70DG soil-application directions in the respective labels.

Weeds Controlled

The combination of PURSUIT PLUS EC and SCEPTER 70DG provides enhanced control of common cocklebur and common ragweed in addition to the weeds controlled by PURSUIT PLUS EC.

For rotational crop information, refer to both the PURSUIT PLUS EC and the SCEPTER 70DG labels. Follow the more restrictive rotational crop guidelines.

PURSUIT PLUS EC + METRIBUZIN FOR ENHANCED CONTROL OF COMMON RAGWEED

PURSUIT PLUS EC may be applied either preplant, preplant incorporated or preemergence in tank-mix combination with metribuzin (Sencor™ or Lexone™) or preemergence with linuron (Lorox™ or Linex™). PURSUIT PLUS EC may also be applied preplant incorporated, followed by a preemergence application of metribuzin or linuron. The tank-mix or sequential application of PURSUIT PLUS EC and metribuzin will enhance the control of common cocklebur and common ragweed in addition to the weeds controlled by a soil application of PURSUIT PLUS EC. The addition of linuron to PURSUIT PLUS EC will enhance the control of common ragweed.

Under conditions of heavy grass pressure, the addition of a grass herbicide such as PROWL 3.3 EC or trifluralin may be tank-mixed with PURSUIT PLUS EC.

See recommendation for addition of Roundup, paraquat or 2,4-D to the spray solution under the NO-TILL OR REDUCED TILLAGE section.

When PURSUIT PLUS EC is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. No label dosages should be exceeded. PURSUIT PLUS EC cannot be mixed with any product containing a label prohibiting such mixtures.

PURSUIT PLUS EC + 2,4-D EARLY PREPLANT APPLICATIONS

PURSUIT PLUS EC will control emerged problem weeds such as velvetleaf, smartweed, common cocklebur, and pigweeds. The weed size should not exceed 3" for velvetleaf and smartweed, and 8" for cocklebur and pigweeds. Add surfactant to the spray mixture at the ratio of 1 qt./100 gal. spray solution AND a nitrogen based fertilizer (such as 28%, 32%N or 10-34-0) at the rate of 1-2 qt./acre for optimum activity. Ammonium sulfate (spray grade) may be substituted for liquid fertilizer at the rate of 4 lbs. per acre.

PURSUIT PLUS EC may be applied up to 45 days prior to soybean planting. If other vegetation is present (not controlled by PURSUIT PLUS EC), apply paraquat, Roundup or 2,4-D in combination with PURSUIT PLUS EC. If 2,4-D is used, apply the following rates of 2,4-D in tank-mixture with PURSUIT PLUS EC:

<u>2,4-D Formulation</u>	<u>Rate (lb a.i./A)</u>	<u>Minimum Days Before Planting</u>
Ester	0.5	7
Amine	0.5	15
Ester or Amine	1.0	30

Use ground equipment only.

FALL APPLICATIONS OF PURSUIT PLUS EC (SOUTH DAKOTA ONLY)

Late fall applications of PURSUIT PLUS EC may be made for control of weeds in NO-TILL soybeans, rather than in the spring prior to planting soybeans. Apply PURSUIT PLUS EC after October 31, and prior to ground freeze-up in the winter. Fall and winter precipitation will activate the herbicide for control of most winter annual weeds and spring germinating weeds in no-till soybeans.

When no-till planting soybeans following small grain harvest, apply a burndown application of Gramoxone™ Extra, Roundup™ or Roundup™ Ultra to the small grain stubble within three weeks of harvest to control weeds present after harvest.

Application Rates

Apply PURSUIT PLUS EC at the rate of 2-1/2 pints/acre. If heavy grass pressure is expected, add PROWL® 3.3EC herbicide to the spray mixture at the rate of 1.25 pint/acre.

**WEEDS CONTROLLED
(PURSUIT PLUS EC ALONE)**

When applied as directed, PURSUIT PLUS EC will control or reduce competition from the weeds listed below.

NOTE: C = Control, R = Reduced Competition

BROADLEAF WEEDS

Weeds Controlled	Preplant Incorporated	Preemergence
Anoda, spurred	C	C
Buffalobur	C*	C
Carpetweed	C	C
Cocklebur, common	R	
Devilsclaw	C	
Galinsoga	C	C
Jimsonweed	C*	
Kochia	C	C
Lambquarters, common	C*	C
Mallow, Venice	C	
Morningglory		
entireleaf	R	
ivyleaf	R	
pitted	R	
smallflower	C	C
tall	R	
Mustard species	C	C
Nightshade		
black	C	C
Eastern black	C	C
hairy	C	C
Pigweed		
palmer	C	C
redroot	C	C
smooth	C	C
spiny	C	C
Puncturevine	C	C
Purslane, common	C	C
Pusley, Florida	C	C
Ragweed,		
common	R	R
giant	R	R
Sida, prickly (teaweed)	C*	
Smartweed		
ladysthumb	C	C
Pennsylvania	C	C
Spurge		
prostrate	C	C
spotted	C	C
Sunflower	C*	
Velvetleaf	C	
Waterhemp, tall**	C	C

- * Cultivation and/or a postemergence herbicide may be required for season-long control.
- ** If a heavy infestation of waterhemp species is anticipated a tank mixture of PURSUIT PLUS EC plus additional PROWL 3.3EC is required for control. Add PROWL 3.3EC to the PURSUIT PLUS EC mixture at the following rates, depending on soil type.

Coarse textured soils: add PROWL 3.3EC at 0.6 pints/acre.
 Medium texture soils: add PROWL 3.3EC at 1.2 to 1.8 pints/acre.
 Fine texture soils: add PROWL 3.3EC at 1.8 pints/acre.

Refer to the PROWL 3.3EC label for specific use rates, application methods and application timings based on soil texture and soil organic matter content. A postemergence herbicide such as Blazer™, Cobra™ or Reflex™ may be needed to control waterhemp species escapes. Refer to individual product labels for specific uses and recommendations.

GRASS WEEDS

Weeds Controlled	Preplant Incorporated	Preemergence
Barnyardgrass	C	C
Crabgrass		
large	C	C
smooth	C	C
Crowfootgrass	C	C
Cupgrass, woolly	C	
Foxtail		
giant	C	C
green	C	C
yellow	C	C
Goosegrass	C	C
Itchgrass	R	
Johnsongrass		
seedling	C	C
rhizome	R	
Millet, wild proso	R	R
Panicum		
browntop	C	C
fall	C	C
Texas	C	C
Sandbur, field	C	C
Shattercane	C	
Signalgrass, broadleaf	C	C
Sorghum alnum	C	R
Witchgrass	C	C

SEDGES

Nutsedge		
yellow	R	
purple	R	

ROTATIONAL CROP GUIDELINES

The following rotational crops may be planted after applying PURSUIT PLUS EC at the recommended rate:

1. Anytime

Lima beans	Soybeans
Southern peas	Peanuts

2. Four months after PURSUIT PLUS EC application:
 - Wheat
 - Edible beans and peas (other than lima beans and Southern peas)

3. Eight and one-half months after PURSUIT PLUS EC application:
 - Field corn
 - Field corn grown for seed

4. Nine and one-half months after PURSUIT PLUS EC application:
 - Alfalfa
 - Barley
 - Rye
 - Tobacco

5. Eighteen months after PURSUIT PLUS EC application:

Cotton	Safflower
Lettuce	Sorghum
Oats	Sunflower
Popcorn	Sweet corn

6. Twenty six months after PURSUIT PLUS EC application.
 - Potatoes

7. Forty months after PURSUIT PLUS EC application.*
 - All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.

*Following forty months after a PURSUIT PLUS EC application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP GUIDELINE, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP GUIDELINE, apply the lime at least 12 months prior to planting the rotational crop.

Use of PURSUIT PLUS EC herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

EXCEPTION TO ROTATIONAL CROP GUIDELINE

Corn inbred lines: Corn inbred seed lines may be planted the year following an application of PURSUIT PLUS EC. Several seed companies have tested a wide range of inbreds for sensitivity to PURSUIT PLUS EC soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, American Cyanamid has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT PLUS EC the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of American Cyanamid Company, all risks and consequences associated with planting seed corn inbreds into fields treated previously with PURSUIT PLUS EC shall be assumed by the user.

PRECAUTIONS

General

Only rotational crops harvested at maturity may be used for feed or food.

In the event of a crop loss due to weather, IMI-CORN, soybeans, lima beans, southern peas or peanuts can be replanted. DO NOT work the soil deeper than 2 inches.

IMI-CORN

There should be an interval of at least 45 days between an application of PURSUIT PLUS EC and corn harvest (silage, fodder or grain).

DO NOT graze or feed treated corn forage, silage, fodder or grain to livestock for at least 45 days after an application of PURSUIT PLUS EC.

If field corn is furrow irrigated, till the soil prior to planting winter wheat or barley. The beds should be broken up and the soil mixed with tillage equipment set to cut 4-6 inches deep.

All soil insecticides, including labeled banded or in-furrow applications, may be used in combination with Pioneer imidazolinone resistant (IR) corn hybrids.

Imidazolinone tolerant hybrids from other seed companies may occasionally exhibit injury symptoms when soil insecticides are used in combination with PURSUIT PLUS EC. DO NOT USE COUNTER® 15G systemic insecticide-nematicide in-furrow with imidazolinone tolerant corn hybrids. Other registered organophosphate insecticides such as banded applications of COUNTER 15G, COUNTER® CR® or THIMET® soil and systemic insecticide, or in-furrow applications of COUNTER CR or other registered carbamate or pyrethroid insecticides may be used in combination with PURSUIT PLUS EC applications. American Cyanamid has not tested all hybrids in which the imidazolinone tolerance trait is claimed and cannot be responsible for factors which are beyond its control, such as growing conditions, environmental conditions, grower practices and the specific genetics of each hybrid tolerance to PURSUIT PLUS EC and insecticide applications.

Soybean

If soybeans are furrow irrigated, till the soil prior to planting winter wheat or barley. The beds should be broken up and the soil mixed with tillage equipment set to cut 4-6 inches deep.

There should be an interval of at least 85 days between an application of PURSUIT PLUS EC and soybean harvest.

DO NOT graze or feed treated soybean forage, hay or straw to livestock.

Except where stated on a supplemental label or in this label, DO NOT apply products containing chlorimuron ethyl (Classic™, Canopy™, Gemini™, Lorox Plus™, Preview™, etc.) or imazaquin (SCEPTER®, SQUADRON®, TRI-SCEPT®, SCEPTER® O.T.®, SCEPTER® 70DG or DETAIL™ herbicides), or products containing imazethapyr (PURSUIT®, PURSUIT® DG, PASSPORT®, CONTOUR® or RESOLVE™ herbicides the same year as PURSUIT PLUS EC or injury to follow crops may occur.

™Bullet, Harness Plus, Lariat, Lasso, Permit and Roundup are trademarks of Monsanto Agricultural Products Company.

™Bicep, Bicep Lite, Dual and Exceed are a trademarks of Ciba Geigy Corporation.

™Buctril is a trademark of Rhone Poulenc Ag Products Company.

™Banvel, Clarity, Frontier and Marksman are a trademarks of Sandoz Crop Protection, Inc.

™Accent, Basis, Classic, Concert, Canopy, Extrazine II, Gemini, Lexone, Lorox, Lorox Plus, Pinnacle, and Staple are trademarks of E.I. duPont deNemours and Company.

™Sencor DF and Sencor 3L are trademarks of Mobay Corporation.

™Linex is a trademark of Griffin Ag Products Co., Inc.

™Broadstrike, TREFLAN and Sonolan are trademarks of DowElanco.

™Basagran, STATUS and PRESTIGE are trademarks of BASF Corporation.

™Eradicane Extra, Surpass, Sutan + and Gramoxone are trademarks of Zeneca, Inc.

™IMI-CORN is a trademark of American Cyanamid Company.

®Registered trademarks of American Cyanamid Company.

For additional information regarding the use of PURSUIT PLUS EC herbicide, call telephone no. 800-942-0500.

American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1996