



# Supplemental Labeling

EPA Reg. No. 241-331

**FOR USE IN ENGLISH PEAS, LIMA BEANS, NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE, PINTO, CRANBERRY, AND SMALL WHITE TYPE DRY BEANS ONLY**

**USE ONLY IN ILLINOIS, IOWA, MICHIGAN, MINNESOTA, NORTH DAKOTA, SOUTH DAKOTA AND WISCONSIN**

**OBSERVE ALL PRECAUTIONARY STATEMENTS AND SPRAYING AND APPLICATION INFORMATION IN THE PURSUIT® PLUS EC HERBICIDE LEAFLET LABEL BEFORE USING. APPLY WITH GROUND EQUIPMENT ONLY.**

### GENERAL INFORMATION

Reduced crop growth, quality, yield and/or delayed maturity may result from a **PURSUIT® PLUS EC herbicide** application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly. **DO NOT** apply **PURSUIT PLUS EC** if planting is delayed and chance of frost prior to maturity is likely.

**DO NOT** apply **PURSUIT PLUS EC** if cold and/or wet conditions are present or predicted to occur within one week of application.

**USE PURSUIT PLUS EC ONLY** if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans.

Use of **PURSUIT PLUS EC** in accordance with labeling directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This labeling must be in the possession of the user at the time of herbicide application.

**DO NOT** apply by air.

**DO NOT** apply **PURSUIT PLUS EC** in any manner not specifically described in this label.

Sensitive crops such as sugar beets, canola, and leafy vegetables may be injured by spray drift or other indirect contact with **PURSUIT PLUS EC**.

Spray equipment used for **PURSUIT PLUS EC** application must be drained and thoroughly cleaned with water before being used to apply other products.

**DO NOT** make more than one application of **PURSUIT PLUS EC** per year.

**DO NOT** apply to Domino variety black turtle beans.

**DO NOT** apply **PURSUIT PLUS EC** after June 30.

Allow at least 60 days between application and harvest of navy, great northern, red kidney, cranberry, black turtle, pinto and small white dry beans. Allow at least 30 days between application and harvest of lima beans and English peas.

ACCEPTED  
7-18-05  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.  
241-331

**Preplant Incorporated Applications.** Apply **PURSUIT® PLUS EC herbicide** within 1 week before planting. Applied preplant incorporated, **PURSUIT PLUS EC** may be tank-mixed with other registered herbicides. Refer to the following table for use rate:

Soil Type	PURSUIT PLUS EC Use Rate (oz/acre)
Sands, loamy sands in Michigan	<b>DO NOT USE</b>
Sandy loams and all other soils with ≤ 2% O.M. in Michigan, Minnesota (north of Highway 210), North Dakota, and South Dakota	20 (1.25 pts)
All other soils with > 2% O.M. in Michigan and any soil type in Iowa, Illinois, Minnesota (south of Highway 210), and Wisconsin. <b>Not for use in North Dakota at this rate.</b>	30

For control of the annual grass weed species listed on the **PROWL® 3.3 EC herbicide** label, add 0.75 to 1 pt/acre of **PROWL 3.3 EC** to the 20 oz/acre rate of

**PURSUIT PLUS EC**, and 0.25 to 0.5 pt/acre of **PROWL 3.3 EC** to the 30 oz/acre rate of **PURSUIT PLUS EC**.

**WEEDS CONTROLLED**

**PURSUIT PLUS EC** applied to low organic matter soils at the broadcast rate of 20 ounces per acre preplant incorporated will control light to moderate infestations of:

- Mustard, wild
- Nightshade, black
- Nightshade, Eastern black
- Pigweed, redroot

**PURSUIT PLUS EC** applied at the broadcast rate of 30 ounces per acre preplant incorporated will control:

- Mustard, wild
- Nightshade, black
- Nightshade, Eastern black
- Nightshade, hairy
- Pigweed, redroot

**ROTATIONAL CROP GUIDELINE**

The following crops may be planted after applying **PURSUIT PLUS EC** as recommended in this label:

1. Anytime:
  - Soybeans
  - Peanuts
  - Edible bean and pea types listed on this label (Refer to heading on page 1)
2. Two months after **PURSUIT PLUS EC** application:
  - Snap Beans
3. Four months after **PURSUIT PLUS EC** application:
  - Barley
  - Edible bean and pea types not listed on this label (Refer to heading on page 1)
  - Wheat
4. Eight and one-half months after **PURSUIT PLUS EC** application:
  - Field corn
  - Field corn grown for seed\*
5. Nine and one-half months after **PURSUIT PLUS EC** application:
  - Alfalfa
  - Tobacco
6. Eighteen months after **PURSUIT PLUS EC** application:
 

Cotton	Popcorn	Sunflower
Lettuce	Safflower	Sweet Corn
Oats	Sorghum	
7. Twenty-six months after **PURSUIT PLUS EC** application:
  - Potatoes

8. Forty months after **PURSUIT® PLUS EC herbicide** application:

All crops not listed elsewhere in this **ROTATIONAL CROP GUIDELINE**. Following forty months after a **PURSUIT PLUS EC** application, and before planting any crop not listed elsewhere in the **ROTATIONAL CROP GUIDELINE**, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip should include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

If the field is limed to adjust pH prior to planting rotational crops not listed in the **ROTATIONAL CROP GUIDELINE**, apply the lime at least 12 months prior to planting the rotational crop.

In the event of a crop loss, the bean or pea crop may be replanted.

**DO NOT** feed treated forage, hay, or straw to livestock.

\* **Several seed companies have tested a wide range of inbreeds for sensitivity to PURSUIT PLUS EC soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, BASF Corporation has not been given access to the inbred data. Growers are directed to contact the seed company for information and recommendations regarding the planting of corn grown for seed in fields treated with PURSUIT PLUS EC the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of BASF Corporation, all risks and consequences associated with planting seed corn inbreeds into fields treated previously with PURSUIT PLUS EC shall be assumed by the user.**

**CONSULT THE PURSUIT PLUS EC LEAFLET LABEL FOR OTHER PRECAUTIONS.**

#### **EXCEPTIONS TO ROTATIONAL CROP GUIDELINE**

**Sweet corn and popcorn varieties: (States of Iowa, Illinois, Michigan, Minnesota, and Wisconsin only).**

Sweet corn and popcorn varieties may be planted the year following an application of **PURSUIT® PLUS EC herbicide**. Some sweet corn and popcorn varieties may be injured when planted at less than 18 months following an application of **PURSUIT PLUS EC herbicide**. Before planting sweet corn for processing, contact the processor company for information and recommendations regarding the tolerance of sweet corn varieties planned for fields treated with **PURSUIT PLUS EC** the previous year. **DO NOT** plant fresh market sweet corn varieties prior to 18 months after **PURSUIT PLUS EC** use. Before planting popcorn, contact the popcorn company for information and recommendations regarding the tolerance of popcorn varieties. Planned for fields treated with **PURSUIT PLUS EC** the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of BASF Corporation, **ALL RISKS AND CONSEQUENCES ASSOCIATED WITH PLANTING SWEET CORN OR POPCORN VARIETIES INTO FIELDS TREATED PREVIOUSLY WITH PURSUIT PLUS EC SHALL BE ASSUMED BY THE USER.**

Stunting and maturity delay or other adverse effects may result when sweet corn or popcorn are planted following **PURSUIT PLUS EC** use.

**Conditions of Sale and Warranty**

The Directions For Use of this product reflects the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. All such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT PERMITTED BY LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

*Pursuit and Prowl are registered trademarks of BASF*

©2005 BASF Corporation  
All rights reserved

000241-00331.20050518b.NVA 2005-04-131-0225  
Supercedes NVA 2005-04-131-0092

BASF Corporation  
Agricultural Products  
26 Davis Drive  
Research Triangle Park, NC 27709

  
The Chemical Company