

**FOR USE IN COTTON
TO AID IN THE CONTROL, OR REDUCE COMPETITION FROM,
LANCELEAF SAGE (MINTWEED)**

OBSERVE ALL PRECAUTIONARY STATEMENTS, MIXING INSTRUCTIONS, AND ROTATIONAL CROP RESTRICTIONS IN THE PROWL 3.3 EC LABEL BEFORE USING.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

NOTE: Existing living vegetation must be controlled at or before the application of PROWL 3.3 EC. This may be accomplished by applying a registered contact herbicide, or by using PROWL 3.3 EC or PROWL 3.3 EC tank mixtures with such a contact herbicide. Consult the contact herbicide label for all directions, precautions, and restrictions.

Apply PROWL 3.3 EC or TRI-4[®] HF (trifluralin) herbicide preplant incorporated at labeled use rates. In order to suppress lanceleaf sage (mintweed; *Salvia reflexa*), follow standard PPI treatments with an application of PROWL preemergence (split shot) at planting or up to 2 days after planting. Apply PROWL 3.3 EC preemergence at 1.8 pts/A on coarse soils, or 2.4 pts/A on medium and fine soils. Refer to the PROWL 3.3 EC leaflet label for soil type classifications. Cotton seed must be covered with soil to prevent direct contact from the preemergence PROWL 3.3 EC application or crop injury may occur. **DO NOT** apply more than 4.8 pts of PROWL 3.3 EC per season.

DO NOT use planter seed covering mechanisms to incorporate preemergence PROWL 3.3 EC treatments. Excessive, or deep cultivation can reduce mintweed control by moving untreated soil into the rows. Rainfall or irrigation is required within 7 days of application for maximum mintweed activity. If application is delayed and rainfall is received before preemergence application is made, reduced activity on mintweed may result on the first flush of mintweed. PROWL 3.3 EC will not control actively growing mintweed.

PROWL 3.3 EC may be applied in combination with other registered herbicides. Consult the PROWL 3.3 EC label for tank mix directions. All applicable directions, restrictions, and precautions on the EPA registered label(s) are to be followed. Refer to the PROWL 3.3 EC leaflet label for specific directions regarding mixing, spraying, and liquid fertilizer compatibility determinations.

DO NOT apply more than 4.8 pts of PROWL 3.3 EC per season.

This label must be in the possession of the user at the time of herbicide application.

[®]Registered Trademark of American Cyanamid Company
PE-_____ 10/6/92 SD



BEST AVAILABLE COPY

