

## FOR USE IN CONSERVATION TILLAGE COTTON IN TEXAS

**OBSERVE ALL PRECAUTIONARY STATEMENTS, MIXING INSTRUCTIONS, AND ROTATIONAL CROP RESTRICTIONS IN THE PROWL 3.3 EC LABEL BEFORE USING.**

### PROWL 3.3 EC AND PROWL 3.3 EC TANK MIXTURES FOR PREPLANT OR PREEMERGENCE APPLICATIONS IN CONSERVATION TILLAGE COTTON PRODUCTION SYSTEMS

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**NOTE:** Existing living vegetation must be controlled at or before the application of PROWL 3.3 EC. This may be accomplished by applying a contact herbicide registered for use in cotton, or by using PROWL 3.3 EC or PROWL 3.3 EC tank mixtures with such a contact herbicide.


Uniformly apply PROWL 3.3 EC, either alone or in combination with other herbicides registered for use in cotton, in 20 to 60 gallons of water per acre using ground equipment. The higher gallonage should be used for fields with a history of heavy weed infestations or if the field has excessive crop residues. Consult the application rate chart on the reverse side of this label for the broadcast rate per acre of PROWL 3.3 EC herbicide for use in conservation tillage cotton. In addition, PROWL 3.3 EC may be applied through chemigation; refer to the PROWL 3.3 EC label for chemigation instructions.

PROWL 3.3 EC may be applied immediately after planting, or up to 30 days before planting. **DO NOT** apply PROWL 3.3 EC postemergence since cotton exposed at the time of application may be killed. PROWL 3.3 EC is most effective in controlling weeds when adequate rainfall or irrigation is received within 7 days after application, otherwise a rotary hoeing, or use of a registered postemergence grass herbicide treatment may be required.

Refer to the PROWL 3.3 EC leaflet label for specific directions regarding mixing, spraying, and liquid fertilizer compatibility determinations.

All applicable directions, restrictions, and precautions on the EPA registered label(s), including tank mix partners are to be followed.

\*Registered Trademark of American Cyanamid Company



(over)

**BEST AVAILABLE COPY**

	Organic Matter	Organic Matter
COARSE	1.8 TO 3.6 pts.	3.6 pts.
MEDIUM	2.4 TO 4.8 pts.	4.8 pts.
FINE	3.6 TO 4.8 pts.	DO NOT USE


\*PROWL 3.3 EC is not recommended for use on peat or muck soils.

**NOTE: DO NOT** feed forage or graze livestock on treated cotton fields. In the event of crop loss due to weather conditions, cotton, or any crop labelled for preplant incorporated treatments of PROWL 3.3 EC may be replanted. If cotton replanting is necessary, **DO NOT** work the soil. Observe all precautions and limitations on tank mix partner labels.

Consult the PROWL 3.3 EC leaflet label for the list of weeds controlled. In order to control weeds other than those listed in the PROWL 3.3 EC leaflet label, a registered tank mix partner may be used. PROWL 3.3 EC plus a registered tank mix partner will control all weeds listed on the partner's label in addition to the weeds listed in the PROWL 3.3 EC leaflet label.

**NOTE: DO NOT** plant winter wheat or winter barley in the fall following PROWL 3.3 EC application at rates above 3.6 pints per acre. **DO NOT** plant no-till wheat the fall following a no-till application of PROWL 3.3 EC in cotton.

This label must be in the possession of the user at the time of herbicide application.



PE- 10/6/92 SD

 **CYANAMID**  
 American Cyanamid Company  
 Agricultural Division  
 Crop Protection Chemicals Department  
 Wayne, NJ 07470  
 ©1992