


**EPA Reg. No. 241-337
FOR USE IN SUGARCANE
(DO NOT USE IN HAWAII)**



OBSERVE ALL PRECAUTIONARY STATEMENTS, MIXING, SPRAYING AND APPLICATION INSTRUCTIONS, AND FOLLOW CROP RESTRICTIONS IN THE PROWL 3.3EC LEAFLET LABEL BEFORE USING. WHEN TANK MIXING, ALWAYS FOLLOW THE MOST RESTRICTIVE LABEL.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PROWL 3.3EC may be applied to plant or ratoon cane preemergence through layby.

APPLICATION INSTRUCTIONS

PROWL 3.3EC herbicide controls most annual grasses and certain broadleaf weeds as they germinate, but will not control established weeds. PROWL 3.3EC should be applied to rows free of weeds and trash.

PROWL 3.3EC may be applied twice per season, preemergence through layby, in plant or ratoon sugarcane. Applications may be band or broadcast. Although there may be adequate crop tolerance for postemergence applications at layby, the spray must be directed under the sugarcane canopy in order to obtain effective weed control.

DO NOT apply more than 14.4 pints per acre of PROWL 3.3EC during one growing season.

Refer to PROWL 3.3EC leaflet label on the PROWL 3.3EC container for complete mixing instructions.

See below for spraying instructions. **DO NOT** apply this product through any type of irrigation system.

PROWL 3.3EC is not recommended for use on peat or muck soils.

BROADCAST RATES OF PROWL 3.3EC HERBICIDE

Apply 4.8 to 7.2 pints of PROWL 3.3EC but **DO NOT** exceed 14.4 pints per acre in one growing season. See spraying instructions below for calculation of band treatment rate.

Use the 7.2 pint rate:

- a. for dark clay soils;
- b. if no mechanical incorporation is planned;
- c. if heavy weed populations are anticipated;
- d. if itchgrass infestation is anticipated;
- e. or if no shaving is planned.

SPRAYING INSTRUCTIONS

GROUND APPLICATIONS

BROADCAST TREATMENT - Apply PROWL 3.3EC in 10 or more gallons of water or in 20 or more gallons of liquid fertilizer per acre. (Refer to the PROWL 3.3EC leaflet label for liquid fertilizer compatibility determinations.) Use a properly calibrated low-pressure (20 to 40 psi) sprayer equipped with appropriate spray nozzles to achieve uniform spray distribution and minimize drift. Keep the by-pass line on or near the bottom of the tank to minimize foaming. Nozzle screens must be no finer than 50 mesh.

BAND TREATMENT - Apply the broadcast equivalent rate and volume per acre. To determine these:

$$\begin{array}{r} \text{Band width} \\ \text{in inches} \end{array} \times \begin{array}{r} \text{Broadcast} \\ \text{RATE} \\ \text{per acre} \end{array} = \begin{array}{r} \text{Band RATE} \\ \text{per acre} \end{array}$$

$$\begin{array}{r} \text{Band width} \\ \text{in inches} \end{array} \times \begin{array}{r} \text{Broadcast} \\ \text{VOLUME} \\ \text{per acre} \end{array} = \begin{array}{r} \text{Band VOLUME} \\ \text{per acre} \end{array}$$

AERIAL APPLICATIONS

Uniformly apply in 5 or more gallons of water per acre. To minimize drift, **DO NOT** apply during periods of gusty winds or when wind conditions favor drifting. It is recommended that a flagman or an automatic mechanical flagging unit on the aircraft be used to avoid overlapping and possible crop injury. **DO NOT** make aerial applications at layby because complete and uniform coverage cannot be obtained.

*Registered Trademark of American Cyanamid Company

(See other side)

BEST AVAILABLE COPY

Mechanical Incorporation:

PROWL 3.3EC should be applied to loosened beds and incorporated into the top 1 to 2 inches of soil within 7 days after application. Ratoon sugarcane must be lightly shaved in early spring to remove the old stubble before incorporation over the line of sugarcane is possible. Carefully adjust equipment to incorporate without causing excessive damage to emerging shoots. Mechanical incorporation can be achieved by the following:

- a. Rolling cultivator (Lilliston type - Lely Roterra¹) set to cut 2 or 3 inches deep and operated two times at 6 to 8 mph. This technique may be used in the spring or at layby.
- b. Rolling disc cultivator (Hipper) set to cut 2 to 3 inches deep and operated two times at 6 to 8 mph. This technique may be used to incorporate between sugarcane lines (rows) at layby only.

WEED SPECIES CONTROLLED

When applied as directed, PROWL 3.3EC will control the following grass and broadleaf weeds:

GRASSES

Barnyardgrass	(<i>Echinochloa crus-galli</i>)
Browntop panicum	(<i>Panicum fasciculatum</i>)
Crabgrass	(<i>Digitaria</i> spp.)
Fall panicum	(<i>Panicum dichotomiflorum</i>)
Foxtail (giant)	(<i>Setaria faberi</i>)
Foxtail (green)	(<i>Setaria viridis</i>)
Foxtail (yellow)	(<i>Setaria lutescens</i>)
Goosegrass	(<i>Eleusine indica</i>)
Itchgrass (Raoulgrass)	(<i>Rottboellia exaltata</i>)*
Johnsongrass (from seed)	(<i>Sorghum halepense</i>)
Signalgrass	(<i>Brachiaria texanum</i>)
Texas Panicum	(<i>Panicum capillare</i>)

BROADLEAF WEEDS

Carpetweed	(<i>Mollugo verticillata</i>)
Florida pusley	(<i>Richardia scabra</i>)
Lambsquarters	(<i>Chenopodium album</i>)
Pigweed	(<i>Amaranthus</i> spp.)
Purslane	(<i>Portulaca oleracea</i>)
Spurge, annuals	(<i>Euphorbia</i> spp.)

*PROWL 3.3EC must be applied at the 7.2 pint/acre broadcast rates for itchgrass control. Surface application (no mechanical incorporation) will provide partial itchgrass control. An additional application of 4.8 to 7.2 pints/acre may be made at layby.

For additional broadleaf weed control, use PROWL 3.3EC in combination with any registered herbicide. Always check tank-mix compatibility with tank-mix partners prior to filling spray tank. Consult partner labels for correct use rates and label restrictions.

NOTE: DO NOT graze treated fields or feed treated forage or fodder to livestock.

This label must be in the possession of the user at the time of herbicide application.

¹Lely Roterra is a trademark of Lely.
PE 4/6/93 SD



CYANAMID

American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1993

BEST AVAILABLE COPY

EPA Reg. No. 241-337

FOR USE IN FIELD CORN GROWN IN NO-TILL SYSTEMS

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PROWL 3.3 EC LEAFLET LABEL BEFORE USING. SEE THE PROWL 3.3 EC LEAFLET LABEL FOR PRECAUTIONS, GENERAL INFORMATION, MIXING AND APPLICATION AND SPRAYING INSTRUCTIONS, USE RATES, USES WITH OTHER PRODUCTS, WEEDS CONTROLLED, AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This label must be in the possession of the user at the time of herbicide application.

CHEMIGATION: Refer to the PROWL 3.3 EC leaflet label for use directions for chemigation. **DO NOT** apply this product through any type of irrigation system unless the label instructions on chemigation are followed.

DO NOT use in California.

PROWL 3.3 EC herbicide controls labeled weeds as they germinate, but it will not control established weeds. Destroy existing weeds before applying PROWL 3.3 EC (except as recommended in specific postemergence combination treatments).

DO NOT apply tank-mix combinations unless each product is labeled for use in no-till systems. If other herbicides are mixed with PROWL 3.3 EC, the label restrictions for these herbicides must also be followed. Observe the most restrictive instructions on any product label.

PROWL 3.3 EC alone and in tank-mix combinations may be applied in field corn grown in no-till systems. **PLANT CORN AT LEAST 1 1/2 INCHES DEEP. CORN SEED MUST BE COMPLETELY COVERED WITH SOIL.** The use of no-till planters under conditions which do not allow good soil coverage of the corn seed can result in reduced crop stand or injury if PROWL 3.3 EC contacts germinating corn seed. Check equipment to ensure good seed coverage. Apply in 20 or more gallons of spray solution per acre.

For additional information regarding the use of PROWL 3.3 EC herbicide, call telephone number (800) 942-0500.

*Registered Trademark of American Cyanamid Company
PE 3/16/93 SD

