

**FOR USE IN SUGARCANE****(DO NOT USE IN HAWAII)**

OBSERVE ALL PRECAUTIONARY STATEMENTS, MIXING, SPRAYING AND APPLICATION INSTRUCTIONS, AND FOLLOW CROP RESTRICTIONS IN THE PROWL 3.3EC LEAFLET LABEL BEFORE USING. WHEN TANK MIXING, ALWAYS FOLLOW THE MOST RESTRICTIVE LABEL.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PROWL 3.3EC may be applied to newly planted or ratoon sugarcane preemergence through layby and again in late summer or early fall to the newly planted sugarcane.

**APPLICATION INSTRUCTIONS**

PROWL 3.3EC herbicide controls most annual grasses and certain broadleaf weeds as they germinate, but will not control established weeds. PROWL 3.3EC should be applied to rows free of weeds and trash.

PROWL 3.3EC may be applied preemergence through layby, to newly planted or ratoon sugarcane, and again in late summer or early fall to the newly planted sugarcane. Applications may be band or broadcast. Although there may be adequate crop tolerance for postemergence applications at layby, the spray must be directed under the sugarcane canopy in order to obtain effective weed control.

**DO NOT** apply more than 14.4 pints per acre of PROWL 3.3EC during one growing season.

**DO NOT** apply within 90 days of harvest.

Refer to PROWL 3.3EC leaflet label on the PROWL 3.3EC container for complete mixing instructions.

See below for spraying instructions. **DO NOT** apply this product through any type of irrigation system.

PROWL 3.3EC is not recommended for use on peat or muck soils.

**BROADCAST RATES OF PROWL 3.3EC HERBICIDE**

Apply 4.8 to 7.2 pints of PROWL 3.3EC but **DO NOT** exceed 14.4 pints per acre in one growing season. See spraying instructions below for calculation of band treatment rate.

**BEST AVAILABLE COPY**

Use the 7.2 pint rate:

- for heavy clay soils;
- if no mechanical incorporation is planned;
- if heavy weed populations are anticipated;
- if itchgrass infestation is anticipated;
- or if no shaving is planned.

**SPRAYING INSTRUCTIONS****GROUND APPLICATIONS**

**BROADCAST TREATMENT** - Apply PROWL 3.3EC in 10 or more gallons of water or in 20 or more gallons of liquid fertilizer per acre. (Refer to the PROWL 3.3EC leaflet label for liquid fertilizer compatibility determinations.) Use a properly calibrated low-pressure (20 to 40 psi) sprayer equipped with appropriate spray nozzles to achieve uniform spray distribution and minimize drift. Keep the by-pass line on or near the bottom of the tank to minimize foaming. Nozzle screens must be no finer than 50 mesh.

**BAND TREATMENT** - Apply the broadcast equivalent rate and volume per acre. To determine these:

Band width in inches		Broadcast RATE	=	Band RATE
_____	X	per acre		per acre
Row width in inches				

Band width in inches		Broadcast VOLUME	=	Band VOLUME
_____	X	per acre		per acre
Row width in inches				

**ACCEPTED**

•Registered Trademark of American Cyanamid Company

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the purpose  
of registration under  
EPA Reg. No. 241-337

(See other side)

aircraft be used to avoid overlapping and possible crop injury. DO NOT make aerial applications at layby because complete and uniform coverage cannot be obtained.

## INCORPORATION INSTRUCTIONS

PROWL 3.3EC must be thoroughly and uniformly incorporated into the soil with either (a) mechanical incorporation equipment as outlined below, or (b) with rainfall or irrigation, if rainfall or irrigation is adequate for good crop and weed emergence and received within 7 days after application. If rainfall or irrigation is not obtained, PROWL 3.3EC should be mechanically incorporated.

### Mechanical Incorporation:

PROWL 3.3EC should be applied to loosened beds and incorporated into the top 1 to 2 inches of soil within 7 days after application. Ratoon sugarcane must be lightly shaved in early spring to remove the old stubble before incorporation over the line of sugarcane is possible. Carefully adjust equipment to incorporate without causing excessive damage to emerging shoots. Mechanical incorporation can be achieved by the following:

- Rolling cultivator (Lilliston type - Lely Roterra<sup>1</sup>) set to cut 2 or 3 inches deep and operated two times at 6 to 8 mph. This technique may be used with all application timings.
- Rolling disc cultivator (Hipper) set to cut 2 to 3 inches deep and operated two times at 6 to 8 mph. This technique may be used to incorporate between sugarcane lines (rows) at layby only.

## WEED SPECIES CONTROLLED

When applied as directed, PROWL 3.3EC will control the following grass and broadleaf weeds:

### GRASSES

Barnyardgrass	( <i>Echinochloa crus-galli</i> )
Browntop panicum	( <i>Panicum fasciculatum</i> )
Crabgrass	( <i>Digitaria</i> spp.)
Fall panicum	( <i>Panicum dichotomiflorum</i> )
Foxtail (giant)	( <i>Setaria faberi</i> )
Foxtail (green)	( <i>Setaria viridis</i> )
Foxtail (yellow)	( <i>Setaria lutescens</i> )
Goosegrass	( <i>Eleusine indica</i> )
Itchgrass (Raoulgrass)	( <i>Rottboellia exaltata</i> )*
Johnsongrass (from seed)	( <i>Sorghum halepense</i> )
Junglerice	( <i>Echinochloa colona</i> )
Signalgrass	( <i>Brachiaria texanum</i> )
Texas Panicum	( <i>Panicum capillare</i> )

### BROADLEAF WEEDS

Carpetweed	( <i>Mollugo verticillata</i> )
Florida pusley	( <i>Richardia scabra</i> )
Lambsquarters	( <i>Chenopodium album</i> )
Pigweed	( <i>Amaranthus</i> spp.)
Purslane	( <i>Portulaca oleracea</i> )
Spurge, annuals	( <i>Euphorbia</i> spp.)

\*PROWL 3.3EC must be applied at the 7.2 pint/acre broadcast rates for itchgrass control. Surface application (no mechanical incorporation) will provide partial itchgrass control. An additional application of 4.8 to 7.2 pints/acre may be made at layby.

For additional broadleaf weed control, use PROWL 3.3EC in combination with any registered herbicide. Always check tank-mix compatibility with tank-mix partners prior to filling spray tank. Consult partner labels for correct use rates and label restrictions.

**NOTE: DO NOT** graze treated fields or feed treated forage or fodder to livestock.

This label must be in the possession of the user at the time of herbicide application.

<sup>1</sup>Lely Roterra is a trademark of Lely  
PE 8506 Rev 4 4/5/94



**CYANAMID**

American Cyanamid Company  
Agricultural Products Division  
Crop Protection Chemicals Department  
Wayne, NJ 07470 ©1994

**BEST AVAILABLE COPY**

EPA REG. NO. 241-337

3 of 3

**FALL APPLICATIONS OF PROWL 3.3 EC  
FOLLOWED BY EARLY POST APPLICATIONS OF PURSUIT®  
HERBICIDE FOR SOYBEANS GROWN NORTH OF I-80**

OBSERVE ALL PRECAUTIONS AND LIMITATIONS ON THE PROWL 3.3 EC LEAFLET LABEL. SEE THE PROWL 3.3 EC LEAFLET FOR APPLICATION INFORMATION AND WEED SPECIES CONTROLLED.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Refer to the PROWL 3.3 EC leaflet label for Chemigation and impregnation on dry bulk fertilizer instructions. **DO NOT** apply PURSUIT through any type of irrigation system.

PROWL 3.3 EC may be surface applied or incorporated for soybean weed control in the fall when followed by a postemergence program using PURSUIT herbicide the following spring. Apply PROWL between October 1 and December 31 for soybeans grown north of Interstate Highway 80 (I-80) at the following rates:

**Broadcast Rate Per Acre of PROWL 3.3 EC**

Soil Texture	PROWL 3.3 EC (pints)
Coarse	1.8
Medium	2.4
Fine	3.6

For situations where incorporation is desired, consult the PROWL 3.3 EC leaflet label for incorporation instructions. Consult the PURSUIT label for PURSUIT herbicide application instructions and precautions.

Roundup<sup>1</sup>, 2, 4-D, Butyrac<sup>2</sup> 200 (2,4-DB), or paraquat should be used to kill existing vegetation, if present before planting. Fall application of PROWL 3.3 EC is not recommended on fields that are poorly drained or are subject to periods of flooding.

OBSERVE ALL PRECAUTIONS, LIMITATIONS, AND CROPPING RESTRICTIONS ON THE PRODUCT LEAFLET LABELS. ALWAYS FOLLOW THE MOST RESTRICTIVE LABEL.

This label must be in the possession of the user at the time of herbicide application.

®Registered Trademark of American Cyanamid Company

<sup>1</sup>Roundup is a trademark of Monsanto Agricultural Products Company

<sup>2</sup>Butyrac 200 is a trademark of Rohm and Haas Agricultural Company

PF- 6/16/94 SD

**BEST AVAILABLE COPY**



**CYANAMID**

American Cyanamid Company  
Agricultural Products Division  
Specialty Products Department  
Wayne, NJ 07470

