

FOR USE IN SUGARCANE GROWN ON MUCK SOILS IN FLORIDA

OBSERVE ALL PRECAUTIONARY STATEMENTS, MIXING, SPRAYING AND APPLICATION INSTRUCTIONS, AND FOLLOW CROP RESTRICTIONS IN THE PROWL 3.3EC LEAFLET LABEL BEFORE USING. WHEN TANK-MIXING, ALWAYS FOLLOW THE MOST RESTRICTIVE LABEL.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PROWL 3.3EC may be applied to plant or ratoon (stubble) cane preemergence through layby.

APPLICATION INSTRUCTIONS

PROWL 3.3EC herbicide controls most annual grasses and certain broadleaf weeds as they germinate, but will not control established weeds. PROWL 3.3EC should be applied to rows free as possible of weeds and trash

PROWL 3.3EC may be applied twice per season, preemergence through layby, in plant or ratoon (stubble) sugarcane. Applications may be band or broadcast. Although there may be adequate crop tolerance for postemergence applications at layby, the spray must be directed under the sugarcane canopy in order to obtain effective weed control.

DO NOT apply within 90 days of harvest

DO NOT apply more than 14.4 pints per acre of PROWL 3.3EC during one growing season

Refer to PROWL 3.3EC leaflet label on the PROWL 3.3EC container for complete mixing instructions

See below for spraying instructions. **DO NOT** apply this product through any type of irrigation system.

BROADCAST RATES OF PROWL 3.3EC HERBICIDE

Apply 4.8 to 7.2 pints of PROWL 3.3EC but **DO NOT** exceed 14.4 pints per acre in one growing season. See spraying instructions below for calculation of band treatment rate. Use the 7.2 pint rate if heavy weed populations are anticipated.

SPRAYING INSTRUCTIONS


GROUND APPLICATIONS

BROADCAST TREATMENT - Apply PROWL 3.3EC in 10 or more gallons of water or in 20 or more gallons of liquid fertilizer per acre. (Refer to the PROWL 3.3EC leaflet label for liquid fertilizer compatibility determinations.) Use a properly calibrated low-pressure (20 to 40 psi) sprayer equipped with appropriate spray nozzles to achieve uniform spray distribution and minimize drift. Keep the by-pass line on or near the bottom of the tank to minimize foaming. Nozzle screens must be no finer than 50 mesh

BAND TREATMENT - Apply the broadcast equivalent rate and volume per acre. To determine these:

Band width in inches		Broadcast RATE	Band RATE
_____	X	per acre	per acre
Row width in inches			

Band width in inches		Broadcast RATE	Band VOLUME
_____	X	per acre	per acre
Row width in inches			



because complete and uniform coverage cannot be obtained.

INCORPORATION INSTRUCTIONS

PROWL 3.3EC MUST be thoroughly and uniformly incorporated into the soil with either (a) rainfall or irrigation, if rainfall or irrigation is adequate for good crop and weed emergence and received within 7 days after application, or (b) mechanical incorporation equipment as outlined below.

Mechanical Incorporation:

PROWL 3.3EC should be incorporated into the top 1 to 2 inches of soil within 7 days after application. Carefully adjust equipment to incorporate without causing excessive damage to emerging shoots. Mechanical incorporation can be achieved by the following:

- a. Rolling cultivator (Lilliston type - Lely Roterra¹) set to cut 2 or 3 inches deep and operated at 6 to 8 mph. This technique may be used with all application timings.
- b. Rolling disc cultivator (Hipper) set to cut 2 to 3 inches deep and operated at 6 to 8 mph. This technique may be used to incorporate between sugarcane lines (rows) at layby only.

WEED SPECIES CONTROLLED

When applied as directed, PROWL 3.3EC will control the following grass weeds:

Barnyardgrass	(<i>Echinochloa crus-galli</i>)	Goosegrass	(<i>Eleusine indica</i>)
Browntop panicum	(<i>Panicum fasciculatum</i>)	Johnsongrass (from seed)	(<i>Sorghum halepense</i>)
Crabgrass	(<i>Digitaria spp.</i>)	Junglerice	(<i>Echinochloa colona</i>)
Fall panicum	(<i>Panicum dichotomiflorum</i>)	Signalgrass	(<i>Brachiaria texanum</i>)
Foxtail (giant)	(<i>Setaria faberi</i>)	Texas Panicum	(<i>Panicum capillare</i>)
Foxtail (green)	(<i>Setaria viridis</i>)		
Foxtail (yellow)	(<i>Setaria litescens</i>)		

PROWL 3.3EC will reduce the competition of certain broadleaf weeds including carpetweed, Florida pusley, lambsquarters, pigweed, purslane and annual spurge.

For additional broadleaf weed control, use PROWL 3.3EC in combination with any registered herbicide. Always check tank-mix compatibility with tank mix partners prior to filling spray tank. Consult partner labels for correct use rates and label restrictions.

NOTE: DO NOT graze treated fields or feed treated forage or fodder to livestock.

This label must be in the possession of the user at the time of herbicide application.

¹Lely Roterra is a trademark of Lely.
PR 2012-01-01