



000241-00337-072998

Printed: 13:25:23 Friday, 21 May, 1999 # 113 / 2979

Systems Integration Group, Inc.

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

<input type="checkbox"/>	Registration
<input checked="" type="checkbox"/>	Amendment
<input type="checkbox"/>	Other

OPP Identifier Number
230748

Application for Pesticide - Section I

1. Company/Product Number 241-337	2. EPA Product Manager J. Tompkins	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) PROWL® 3.3 EC herbicide	PM# 25	
5. Name and Address of Applicant (Include ZIP Code) American Cyanamid Company Agricultural Research Division P.O. Box 400 Princeton, NJ 08543-0400 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response Agency letter dated _____	NOTIFICATION JUL 29 1998
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	
<input type="checkbox"/> Notification - Explain below.	<input checked="" type="checkbox"/> Other - Explain below.	

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Request for Approval of Two New Supplemental Labels Allowing Fall Applications in Soybeans Grown North of I-80 and in the State of Iowa and a New Supplemental Label for the Control of Russian Thistle in Cotton Grown in Arizona.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____		
<i>Verification must submitted</i>		If "Yes" Unit Packaging wgt. No. per container			
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Zareen Ahmed	Title Product Registrations Manager	Telephone No. (Include Area Code) 609-716-2321
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamp)
2. Signature 	3. Title Product Registrations Manager	
4. Typed Name Zareen Ahmed	5. Date 7/14/98	

PROWL[®]

3.3 EC herbicide

Page 2 of 5

Supplemental Labeling

EPA REG. NO. 241-337

**FALL APPLICATIONS OF PROWL 3.3 EC
FOLLOWED BY EARLY POST APPLICATIONS OF RAPTOR[®] HERBICIDE,
FOR SOYBEANS GROWN NORTH OF
INTERSTATE 80 (I-80) AND IN THE ENTIRE STATE OF IOWA**

OBSERVE ALL PRECAUTIONARY STATEMENTS ON THE PROWL 3.3 EC LEAFLET LABEL BEFORE USING. OBSERVE ALL USE DIRECTIONS INCLUDING PRECAUTIONS, RESTRICTIONS AND LIMITATIONS, GENERAL INFORMATION, MIXING AND APPLICATION INSTRUCTIONS, USES WITH OTHER PRODUCTS, WEEDS CONTROLLED AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Refer to the PROWL 3.3 EC leaflet label for impregnation on dry bulk fertilizer instructions. **DO NOT** apply RAPTOR through any type of irrigation system.

For soybean weed control, PROWL 3.3 EC may be surface applied or incorporated in the fall when followed by a postemergence program using RAPTOR herbicide the following spring. Apply PROWL 3.3 EC between October 1 and December 31 at the following rates:

Broadcast Rate Per Acre of PROWL 3.3 EC

Soil Texture	PROWL 3.3 EC (pints)
Coarse	2.4
Medium	3.0 to 3.6
Fine	3.6

For situations where incorporation is desired, consult the PROWL 3.3 EC leaflet label for incorporation instructions. Minimize furrow and ridge formation in the tillage operation. Consult the RAPTOR label for RAPTOR herbicide application instructions and precautions.

Roundup¹, 2, 4-D, Butyrac² 200 (2,4-DB), or paraquat should be used to kill existing vegetation, if present before planting. Fall application of PROWL 3.3 EC is not recommended on fields that are poorly drained or are subject to periods of flooding.

OBSERVE ALL PRECAUTIONS, LIMITATIONS, AND CROPPING RESTRICTIONS ON THE PRODUCT LEAFLET LABELS. ALWAYS FOLLOW THE MOST RESTRICTIVE LABEL.

This label must be in the possession of the user at the time of herbicide application.

[®]Registered Trademarks of American Cyanamid Company

¹Roundup is a trademark of Monsanto Agricultural Products Company


²Butyrac 200 is a trademark of Rohm and Haas Agricultural Company

PE-XXXX 7/14/98

NOTIFICATION

1111 29 1998

 **CYANAMID**
American Cyanamid Company
North America Agricultural Products Division
Crop Protection Products Department
One Campus Drive, Parsippany, NJ 07054 ©1998



PROWL[®]

3.3 EC herbicide

Supplemental Labeling

EPA Reg. No. 241-337

FOR USE IN COTTON IN THE STATE OF ARIZONA ONLY TO AID IN THE CONTROL OF RUSSIAN THISTLE

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PROWL 3.3 EC AND TANK MIX PARTNER LEAFLET LABELS BEFORE USING. SEE THE PROWL 3.3 EC AND TANK MIX PARTNER LEAFLET LABELS FOR PRECAUTIONS, GENERAL INFORMATION, MIXING AND APPLICATION AND SPRAYING INSTRUCTIONS, USE RATES, USES WITH OTHER PRODUCTS, WEEDS CONTROLLED, ROTATIONAL CROP RESTRICTIONS AND DISCLAIMERS. OBSERVE THE MOST RESTRICTIVE INSTRUCTIONS ON ANY PRODUCT LABEL.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This label must be in the possession of the user at the time of herbicide application.

In order to suppress Russian thistle (*Salsola iberica*), apply PROWL 3.3 EC herbicide preplant incorporated. Apply PROWL 3.3 EC up to 60 days prior to planting and incorporate within 7 days of application. Refer to the PROWL 3.3 EC leaflet label for soil type classifications.

Broadcast Rate Per Acre of PROWL 3.3 EC

Soil Type	PROWL 3.3 EC (pints per Acre)
Coarse	1.2 to 2.4 pts
Medium	1.8 to 2.4 pts
Fine	2.4 to 3.6 pts

The high rates for each soil texture above should be used if heavy weed populations are anticipated, or extensive crop residues are present prior to seed bed preparation.

Apply PROWL 3.3 EC tank mixes and sequential programs as specified under the tank mix section of the leaflet label. When PROWL 3.3 EC is tank mixed with other herbicides, follow all cropping restrictions, precautionary statements and limitations in the tank mix partner leaflet label. Always follow the most restrictive label recommendations. Refer to specific companion product label(s) to determine specific use rates and weeds controlled. In addition, follow all precautions and restrictions including State and local use restrictions that may apply.

NOTIFICATION

(see reverse side)

JUL 29 1998

PROWL[®]

3.3 EC herbicide

page 5 of 5

Supplemental Labeling

EPA REG. NO. 241-337

FALL APPLICATIONS OF PROWL 3.3 EC FOR EARLY SPRING SEASON GRASS AND SMALL SEEDED-BROADLEAF WEED CONTROL FOR SOYBEANS GROWN NORTH OF INTERSTATE 80 (I-80) AND IN THE ENTIRE STATE OF IOWA

OBSERVE ALL PRECAUTIONARY STATEMENTS ON THE PROWL 3.3 EC LEAFLET LABEL BEFORE USING. OBSERVE ALL USE DIRECTIONS INCLUDING PRECAUTIONS, RESTRICTIONS AND LIMITATIONS, GENERAL INFORMATION, MIXING AND APPLICATION INSTRUCTIONS, USES WITH OTHER PRODUCTS, WEEDS CONTROLLED AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Refer to the PROWL 3.3 EC leaflet label for impregnation on dry bulk fertilizer instructions.

For soybean weed control, PROWL 3.3 EC may be surface applied or incorporated in the fall. Fall applications of PROWL 3.3 EC will not provide season long weed control. For season long weed control, use a labeled early postemergence herbicide such as PURSUIT[®] herbicide or RAPTOR[®] herbicide. Apply PROWL 3.3 EC between October 1 and December 31 at the following rates:

Broadcast Rate Per Acre of PROWL 3.3 EC

Soil Texture	PROWL 3.3 EC (pints)
Coarse	2.4
Medium	3.0 to 3.6
Fine	3.6

For situations where incorporation is desired, consult the PROWL 3.3 EC leaflet label for incorporation instructions. Minimize furrow and ridge formation in the tillage operation.

Roundup¹, 2, 4-D, Butyrac² 200 (2,4-DB), or paraquat should be used to kill existing vegetation, if present before planting. Fall application of PROWL 3.3 EC is not recommended on fields that are poorly drained or are subject to periods of flooding.

OBSERVE ALL PRECAUTIONS, LIMITATIONS, AND CROPPING RESTRICTIONS ON THE PRODUCT LEAFLET LABELS. ALWAYS FOLLOW THE MOST RESTRICTIVE LABEL.

This label must be in the possession of the user at the time of herbicide application.

[®]Registered Trademarks of American Cyanamid Company

¹Roundup is a trademark of Monsanto Agricultural Products Company

²Butyrac 200 is a trademark of Rohm and Haas Agricultural Company

PE- 7/14/98

NOTIFICATION

JUL 29 1998



CYANAMID

American Cyanamid Company
North America Agricultural Products Division
Crop Protection Products Department
One Campus Drive, Parsippany, NJ 07054 ©1998