



000241-00337-100798

Printed: 13:24:53 Friday, 21 May, 1999 # 111 / 2977

Systems Integration Group, Inc.

PROWL[®] 3.3 EC herbicide

Supplemental Labeling

EPA REG. NO. 241-337

FALL APPLICATIONS OF PROWL 3.3 EC FOR EARLY SPRING SEASON GRASS AND SMALL SEEDED-BROADLEAF WEED CONTROL FOR SOYBEANS GROWN NORTH OF INTERSTATE 80 (I-80) AND IN THE ENTIRE STATES OF IOWA, OKLAHOMA AND TEXAS

OBSERVE ALL PRECAUTIONARY STATEMENTS ON THE PROWL 3.3 EC LEAFLET LABEL BEFORE USING. OBSERVE ALL USE DIRECTIONS INCLUDING PRECAUTIONS, RESTRICTIONS AND LIMITATIONS, GENERAL INFORMATION, MIXING AND APPLICATION INSTRUCTIONS, USES WITH OTHER PRODUCTS, WEEDS CONTROLLED AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Refer to the PROWL 3.3 EC leaflet label for impregnation on dry bulk fertilizer instructions.

For soybean weed control, PROWL 3.3 EC may be surface applied or incorporated in the fall. Fall applications of PROWL 3.3 EC will not provide season long weed control. For season long weed control, use a labeled early postemergence herbicide such as PURSUIT[®] herbicide or RAPTOR[®] herbicide. Apply PROWL 3.3 EC between October 1 and December 31 at the following rates:

Broadcast Rate Per Acre of PROWL 3.3 EC

Soil Texture	PROWL 3.3 EC (pints)
Coarse	2.4
Medium	3.0 to 3.6
Fine	3.6

For situations where incorporation is desired, consult the PROWL 3.3 EC leaflet label for incorporation instructions. Minimize furrow and ridge formation in the tillage operation.

Roundup¹, 2, 4-D, Butyrac² 200 (2,4-DB), or paraquat should be used to kill existing vegetation, if present before planting. Fall application of PROWL 3.3 EC is not recommended on fields that are poorly drained or are subject to periods of flooding.

OBSERVE ALL PRECAUTIONS, LIMITATIONS, AND CROPPING RESTRICTIONS ON THE PRODUCT LEAFLET LABELS. ALWAYS FOLLOW THE MOST RESTRICTIVE LABEL.


This label must be in the possession of the user at the time of herbicide application.

[®]Registered Trademarks of American Cyanamid Company

¹Roundup is a trademark of Monsanto Agricultural Products Company

²Butyrac 200 is a trademark of Rohm and Haas Agricultural Company

PE- 9/4/98



241-

PROWL[®]

3.3 EC herbicide

Supplemental Labeling

EPA Reg. No. 241-337

FOR USE IN COTTON IN THE STATE OF ARIZONA ONLY TO AID IN THE CONTROL OF RUSSIAN THISTLE

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PROWL 3.3 EC AND TANK MIX PARTNER LEAFLET LABELS BEFORE USING. SEE THE PROWL 3.3 EC AND TANK MIX PARTNER LEAFLET LABELS FOR PRECAUTIONS, GENERAL INFORMATION, MIXING AND APPLICATION AND SPRAYING INSTRUCTIONS, USE RATES, USES WITH OTHER PRODUCTS, WEEDS CONTROLLED, ROTATIONAL CROP RESTRICTIONS AND DISCLAIMERS. OBSERVE THE MOST RESTRICTIVE INSTRUCTIONS ON ANY PRODUCT LABEL.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This label must be in the possession of the user at the time of herbicide application.

In order to suppress Russian thistle (*Salsola iberica*), apply PROWL 3.3 EC herbicide preplant incorporated. Apply PROWL 3.3 EC up to 60 days prior to planting and incorporate within 7 days of application. Refer to the PROWL 3.3 EC leaflet label for soil type classifications.

Broadcast Rate Per Acre of PROWL 3.3 EC

Soil Type	PROWL 3.3 EC (pints per Acre)
Coarse	1.2 to 2.4 pts
Medium	1.8 to 2.4 pts
Fine	2.4 to 3.6 pts

The high rates for each soil texture above should be used if heavy weed populations are anticipated, or extensive crop residues are present prior to seed bed preparation.

Apply PROWL 3.3 EC tank mixes and sequential programs as specified under the tank mix section of the leaflet label. When PROWL 3.3 EC is tank mixed with other herbicides, follow all cropping restrictions, precautionary statements and limitations in the tank mix partner leaflet label. Always follow the most restrictive label recommendations. Refer to specific companion product label(s) to determine specific use rates and weeds controlled. In addition, follow all precautions and restrictions including State and local use restrictions that may apply.

ACCEPTED

OCT 7 1998

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 241-337

PROWL 3.3 EC will not control established weeds. Destroy existing weeds before or during application of PROWL 3.3 EC. Unusually cold, excessively wet, or hot dry conditions that delay germination or extend germination over a long period of time may reduce weed control.

NOTE: If loss of crop occurs due to adverse weather conditions, cotton or any crop registered for PROWL 3.3 EC preplant incorporated use can be replanted the same year into PROWL 3.3 EC treated soils without adverse effects. If replanting is necessary, **DO NOT** rework the soil deeper than the herbicide treated zone.

See the PROWL 3.3 EC leaflet label for additional weeds controlled and rotational crop restrictions.



®Registered Trademark of American Cyanamid Company

7/6/98

PE-XXXX