

241-337

01/27/2000

1/2

PROWL[®]

3.3 EC herbicide

Supplemental Labeling

FOR DISTRIBUTION AND USE ONLY WITHIN LOUISIANA

EPA Reg. No. 241-337

FOR USE ON FALLOW GROUND AND NON-IRRIGATION NON-CROPPED WATER DRAINAGE AREAS (DITCHBANKS) IN SUGARCANE

OBSERVE ALL PRECAUTIONARY STATEMENTS ON THE PROWL 3.3 EC LEAFLET LABEL BEFORE USING. SEE THE PROWL 3.3 EC LABEL FOR MIXING AND APPLICATION INFORMATION, WEED SPECIES CONTROLLED, AND FOLLOW CROP RESTRICTIONS. THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF HERBICIDE APPLICATION.

DIRECTIONS FOR USE

FALLOW GROUND APPLICATION

PROWL 3.3 EC applied according to this labeling, prior to weed emergence, will control annual grasses such as itchgrass (*Raouigrass*), seedling Johnsongrass and *Panicum* spp. in preplant fallow ground sugarcane. If necessary, control weeds that have emerged prior to application of PROWL 3.3 EC with postemergence herbicides and/or mechanical cultivation.


After cultivation and the beds have been formed in the spring, apply 3 quarts of PROWL 3.3 EC per acre in 10 or more gallons per acre of spray using ground equipment. Sugarcane beds should be free of trash or clods at the time of application.

If sufficient rainfall (1/2 to 3/4 inch) has not occurred within 7 days of application, shallow incorporation (1 to 2 inches) with an additional pass of a Lilliston-type-Lely Roterra¹ cultivator set to cut 2 or 3 inches deep is recommended.

A minimum interval of 60 days between PROWL 3.3 EC application and planting of sugarcane is required or crop injury may occur.

After planting, use PROWL 3.3 EC according to the leaflet label directions for sugarcane and DO NOT exceed 14.4 pints per acre during one growing season.

ACCEPTED
JAN 27 2000
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 241-337



(See other side)

NON-IRRIGATION NON-CROPPED WATER DRAINAGE AREAS (DITCHBANKS) APPLICATION

Apply 3 to 4 quarts of PROWL 3.3 EC per acre in 20 or more gallons per acre of spray to non-cropped ditchbanks adjacent to sugarcane fields. Use ground equipment and apply prior to weed emergence. If necessary, control weeds that have emerged prior to application of PROWL 3.3 EC with postemergence herbicides.

DO NOT apply PROWL 3.3 EC when water is present in the drainage area (ditchbank).

DO NOT apply PROWL 3.3 EC below the high water mark.

DO NOT exceed 14.4 pints of PROWL 3.3 EC per acre during one growing season.

When tank mixing with PROWL 3.3 EC, refer to the tank-mix partner product label for application rates and timing by soil types, weed species, and weed or crop growth stage; restrictions, precautions, adjuvant recommendations, weeds controlled and other information. Follow all precautions and restrictions including state and local use restrictions that may apply to specific products.

When tank-mixing herbicides, always read and follow the most restrictive label recommendations.

Refer to the PROWL 3.3 EC leaflet label for additional information regarding WEEDS CONTROLLED, MIXING AND SPRAYING INSTRUCTIONS



® Registered Trademark of American Cyanamid Company.
¹ Trademark of Lely

